

**Green algae in soil:
assessing their biodiversity and biogeography
with molecular-phylogenetic methods
based on cultures**

Dissertation

zur Erlangung des mathematisch-naturwissenschaftlichen Doktorgrades

"Doctor rerum naturalium"

der Georg-August-Universität Göttingen

im Promotionsprogramm Biologie

der Georg-August University School of Science (GAUSS)

vorgelegt von

Ladislav Hodač

aus Mladá Boleslav (Tschechien)

Göttingen, 2015

Betreuungsausschuss

Prof. Dr. Thomas Friedl, Experimentelle Phykologie und Sammlung von Algenkulturen, Albrecht-von-Haller-Institut für Pflanzenwissenschaften

Prof. Dr. Rolf Daniel, Genomische und Angewandte Mikrobiologie und Göttinger Genomlabor, Institut für Mikrobiologie und Genetik

Mitglieder der Prüfungskommission

Referent: Prof. Dr. Thomas Friedl, Experimentelle Phykologie und Sammlung von Algenkulturen, Albrecht-von-Haller-Institut für Pflanzenwissenschaften

Korreferent: Prof. Dr. Rolf Daniel, Genomische und Angewandte Mikrobiologie und Göttinger Genomlabor, Institut für Mikrobiologie und Genetik

Weitere Mitglieder der Prüfungskommission:

Prof. Dr. Elvira Hörandl, Systematik, Biodiversität und Evolution von Pflanzen (mit Herbarium), Albrecht-von-Haller-Institut für Pflanzenwissenschaften

PD Dr. Gernot Arp, Abteilung Geobiologie, Geowissenschaftliches Zentrum

PD Dr. Michael Hoppert, Abteilung für Allgemeine Mikrobiologie, Institut für Mikrobiologie und Genetik

Prof. Dr. Hermann Behling, Abteilung Palynologie und Klimadynamik, Albrecht-von-Haller-Institut für Pflanzenwissenschaften

Tag der mündlichen Prüfung: 13.01.2016

Contents

| | |
|-----------------------------------|------------|
| General Introduction | 1 |
| Chapter 1..... | 11 |
| Chapter 2..... | 45 |
| Chapter 3..... | 68 |
| Chapter 4..... | 90 |
| General Summary | 103 |
| References | 105 |
| Appendix Chapter 1 | 133 |
| Appendix Chapter 2 | 159 |
| Appendix Chapter 3 | 164 |
| Appendix Chapter 4 | 179 |
| Publications..... | 182 |
| Danksagung..... | 185 |

General Introduction

Importance of soil microalgae

Terrestrial eukaryotic microalgae were recognized as a part of soil microflora in the late 19th century and were introduced into soil ecology during the second half of the 20th century (Gollerbach and Shtina 1969; Shtina and Gollerbach 1976; Metting 1981; Starks *et al.* 1981). As photoautotrophs, microalgae constitute the basis of soil food webs and play a key role in soil development (Zenova and Shtina 1990; Zenova *et al.* 1995; Hu *et al.* 2002; Hu *et al.* 2003b; Rahmonov and Piątek 2007; Langhans *et al.* 2009; Viles 2012). The role of terrestrial eukaryotic microalgae in geomorphic processes was discussed by Viles (2012). Changes in algal abundance and diversity enable biological monitoring of soil recovery after anthropogenic pollution and contamination (Megharaj *et al.* 1986; Dahlin *et al.* 1997; Megharaj *et al.* 2000; Bérard *et al.* 2004; Krivorotov and Bukareva 2005; Nagy *et al.* 2005; Novakovskaya and Patova 2007; Kalinowska and Pawlik-Skowrońska 2010; Temraleeva *et al.* 2011; Hastings *et al.* 2014), or during postpyroge soil development (Scherbina *et al.* 2014). Some authors recognized the importance of algal diversity in quality assessment of agricultural ecosystems (Shields and Durrell 1964; Gollerbach and Shtina 1969; Kabirov 1993; Hastings *et al.* 2014) and of land usage (King and Ward 1977). Particular species of soil microalgae are considered to be biological indicators (Balezina 1974; Shtina 1990; Megharaj *et al.* 2000; Bérard *et al.* 2005; Krivorotov and Bukareva 2005; Zancan *et al.* 2006; Dotsenko 2008), classified as a part of ecologically meaningful algocenoses (Gollerbach and Shtina 1969; Zenova *et al.* 1995; Kuzyakhmetov 1998; Şalaru *et al.* 2008; Kabirov *et al.* 2010). The interest in soil microalgae resulted in a series of comprehensive reviews on algal taxonomy, diversity, ecology and geographic distribution (Reisigl 1964; Gollerbach and Shtina 1969; Shtina and Gollerbach 1976; Metting 1981; Starks *et al.* 1981; Shtina 1990; Ettl and Gärtner 1995; Ettl and Gärtner 2014).

Distribution of soil microalgae: a microscopy-based approach

Our knowledge of algal species diversity in soils is still based mostly upon morphological observations accomplished via light microscopy of mixed or monoclonal algal cultures. Since the major fraction of algal species is invisible by direct microscopy of soil particles, algal growth should be first triggered via enrichment culturing techniques (Starks *et al.* 1981; Ettl and Gärtner 1995). The growth of particular species of microalgae is differentially favored according to the chemical composition of culturing media, or whether algae are cultured in liquid or agarized

media. The most widely used medium for culturing of terrestrial microalgae, the Bold's Basal Medium (BBM; Bischoff and Bold 1963), is selective in favor of easy-growing (generalist) rather than rare species (Broady 1996). Our picture of the algal diversity in soils is incomplete, not only due to selective culturing techniques but also because of the demanding morphological determination of similarly looking species.

Research on diversity of edaphic microalgae has a long tradition particularly in Europe, beginning in the 19th century (Kützing 1843; Hansgirg 1888). In fact, most of the early recognized microscopic algae are also known from soils, e.g., *Stichococcus* (Nägeli 1849), *Chlorella* (Beijerinck 1893), *Hormidium/Klebsormidium* (Kützing 1843), *Chlamydomonas* (Ehrenberg 1833), *Coccomyxa* (Schmidle 1901), *Scenedesmus* (Meyen 1829) or *Pleurastrum/Leptosira* (Borzi 1895). Apart from several studies from British Isles (Bristol 1920; 1927; Lund 1945; Lund 1947), the most biodiversity research on soil microalgae was conducted in continental Europe. Considering terrestrial ecosystems of Central and Western Europe, species lists of edaphic microalgae were revealed from forests (Trenkwalder 1975; Komáromy 1983; Kostikov *et al.* 2001a; Hoffmann *et al.* 2007; Škaloud 2009), grasslands (Komáromy 1976; Komáromy 1983; Neustupa 2001), agricultural fields (Lukešová 1993; Zancan *et al.* 2006), urban regions (Perútková 2014), contaminated (e.g., post-mining) areas (Lukešová and Komárek 1987; Lukešová and Hoffmann 1996; Frouz *et al.* 2001; Lukešová 2001; Neustupa and Škaloud 2004; Kalinowska *et al.* 2008; Trzcińska and Pawlik-Skowrońska 2008), subalpine soils (Rosa and Lhotský 1971; Lukešová *et al.* 2010), alpine soils (Reisigl 1958; 1964; Türk and Gärtner 2001; Peer *et al.* 2010) and xeric sand soils (Hoppert *et al.* 2004; Rahmonov and Piątek 2007; Langhans *et al.* 2009). Particular attention was paid to the edaphic microalgae in Eastern Europe and Russia, focusing on characteristic habitats including steppe grasslands (Musabaeva 2009), steppe forests (Kostikov *et al.* 2001b; Maltseva 2005b; 2005a; 2007; Maltsev 2013; Scherbina *et al.* 2014), Carpathian and other forests (Chornevych *et al.* 2008; Iljushenko 2008; Posrednikova *et al.* 2009; Nikorych and Chornevych 2012; Maltsev and Negrulja 2013), Russian tundra (Novakovskaya and Patova 2008; Patova and Dorokhova 2008), agricultural ecosystems (Asfandijarova 2008; Șalaru *et al.* 2008; Maltseva *et al.* 2009), disturbed soils (e.g., post-mining) (Novakovskaya and Patova 2007; Bachura and Khramchenkova 2008), urban areas (Soare and Dobrescu 2010; Shekhovtseva 2014; Dorokhova *et al.* 2015), or islands in the Black Sea (Vinogradova and Darienko 2008; Darienko 2012).

The best sampled region outside Europe is North America, where the investigators focused on algal communities of mountainous regions (Bischoff and Bold 1963; Khaybullina *et al.* 2010), deserts (Flechtner *et al.* 1998; Flechtner *et al.* 2005; Flechtner *et al.* 2008), xeric sandy soils

(Smith *et al.* 2004), contaminated soils (Maxwell 1991; Nagy *et al.* 2005), serpentine soils (Terlizzi and Karlander 1979) and soils of agricultural landscapes (Fairchild and Willson 1967; Hunt *et al.* 1979; Metting and Rayburn 1979; Metting 1981). Microalgae of desert soil crusts were further studied in Africa (Büdel *et al.* 2009; Mansour and Shaaban 2010) and Asia (Islam 1960; Friedmann *et al.* 1967; Novichkova-Ivanova 1972; Chunxiang and Yongding 2003; Hu *et al.* 2003b; Al-Fredan and Fathi 2007; Bhatnagar *et al.* 2008). A lot of attention was paid to the edaphic microflora of the extremely cold and dry polar regions including both the Arctic (Androsova 1964; Broady 1978; Elster *et al.* 1999; Kaštovská *et al.* 2005; Stibal *et al.* 2006; Kaštovská *et al.* 2007; Matula *et al.* 2007; Kim *et al.* 2008; Patova and Dorokhova 2008) and the Antarctic (Broady 1979a; Broady 1984; Broady 1989; 1996; Mataloni *et al.* 2000; Cavacini 2001; Fermani *et al.* 2007). The least known is the algal diversity in humid tropical soils and so far we are aware of investigations from South and Central America (Archibald 1972; Archibald and Bold 1975; Tell 1976; Büdel 2003; Spitzer *et al.* 2014), Asia (MacEntee *et al.* 1977; Watanabe 1983; Reynaud and Lumpkin 1988; Ray and Thomas 2012; Lin *et al.* 2013), Australia and Oceania (Arvik and Willson 1974; Carson and Brown 1976; MacEntee *et al.* 1977; Carson and Brown 1978; Watanabe 1983). The edaphic microflora of small and isolated islands is almost unknown, except for e.g. St. Paul's Rocks in the equatorial Atlantic Ocean (Smith *et al.* 1974).

Phylogenetic diversity of soil microalgae

Considering the modern system of Eukaryotes, soil microalgae comprise photoautotrophic microorganisms that are phylogenetically nested within five kingdoms (Archeplastida, Excavata, Stramenopila, Alveolata and Cryptophyceae; **Fig. 1**). Archaeplastida (Adl *et al.* 2005)—mainly Chlorophyta and Streptophyta (Friedl and Rybalka 2012; Leliaert *et al.* 2012)—represents a dominant group of the most soil types (Broady 1979b; Metting 1981; Hoffmann 1989; Megharaj *et al.* 2000; Lukešová 2001; Zancan *et al.* 2006; Trzcińska and Pawlik-Skowrońska 2008; Freeman *et al.* 2009; Lin and Wu 2014). Stramenopila (Patterson 1989) are as well abundant in soils, represented by the classes Bacillariophyceae (Lukešová and Komárek 1987; Fermani *et al.* 2007; Darienko 2012; Stanek-Tarkowska and Noga 2012), Xanthophyceae (Broady 1979b; Maltseva 2005b; 2005a; Posrednikova *et al.* 2009; Novis *et al.* 2015) and Eustigmatophyceae (Neustupa 2001; Khaybullina *et al.* 2010; Darienko 2012; Scherbina *et al.* 2014). Three further groups were sporadically recorded from soils: Excavata-Euglenophyta (Ashley *et al.* 1985; Maltseva 2005a; Şalaru *et al.* 2008; Lukešová *et al.* 2010; Lang *et al.* 2011), Alveolata-Dinophyta (Ettl and Gärtner 1995; Kutovaya *et al.* 2012) and Cryptophyceae-incertae sedis (Paulsen *et al.* 1992; Ettl and Gärtner 1995). Across the temperate zone, particularly the Green algae

(Chlorophyta lineage) are highly abundant in soils (Sumbali and Mehrotra 2009), whereas in tropics Cyanobacteria dominate (Roger and Reynaud 1982; Lin *et al.* 2013). It is widely accepted that variation of key physico-chemical parameters in soils (pH, organic carbon, nitrogen) affect green algae less than other microalgae (Metting 1981; Starks *et al.* 1981; Hoffmann 1989). Examples from temperate forests and grasslands show, that alkaline and nutrient rich soils favor higher group diversity comprising Cyanobacteria, Diatoms, Xanthophytes and Green algae, whereas rather acidic and nutrient poor soils are often inhabited only by Green algae (Sumbali and Mehrotra 2009; Nikorych and Chornevych 2012). Compared to different terrestrial microhabitats (hard natural or artificial substrates, tree bark, deadwood), the diversity of microalgae in soils is much higher and rarely dominated by a single group (Hallmann *et al.* 2011b; Kulichová *et al.* 2014). In soils of temperate regions, usually several groups of microalgae co-dominate.




Figure 1. Schematic tree of Eukaryotes and their major lineages. Red branches represent algal lineages which include species living in soils. Adopted and modified after del Campo *et al.* (2014).

Molecular diversity of soil Green microalgae

Considering modern systematics of Green algae, we recognize 20 lineages, which comprise terrestrial species (Rindi *et al.* 2009; Friedl and Rybalka 2012; Leliaert *et al.* 2012), **Fig. 2**. The most terrestrial green microalgae belong to the Chlorophyta lineage, which has an ancestor in common with the Streptophyta (the estimated split of both lineages is in Neoproterozoic; Becker 2013). Whereas the Streptophyta (= primarily freshwater lineage) colonized terrestrial habitats at first, Chlorophyta (= primarily marine lineage) colonized land later on (Becker and Marin 2009). The molecular evidence of green algal diversity in terrestrial habitats is still scant. The most molecular data were so far retrieved from soils, e.g., Alpine glacier forefields (Frey *et al.* 2013), Siberian permafrost (Vishnivetskaya 2009), Himalaya (Schmidt *et al.* 2011; Schmidt and Darcy




Figure 2. Schematic tree of Chlorophyta, Streptophyta and their major lineages. Red letters represent algal lineages which include soil dwelling species. Adopted and modified after Leliaert *et al.* (2012).

2014), North-American desert crusts (Lewis and Flechtner 2002; Lewis and Lewis 2005; Flechtner *et al.* 2013) and mountains (Freeman *et al.* 2009), Atacama desert volcano (Costello *et al.* 2009), Ecuadorean rainforest (Faßhauer *et al.* 2011; Spitzer *et al.* 2014) and Antarctica (Lawley *et al.* 2004; Schmidt *et al.* 2011). A few molecular diversity studies recovered green microalgae in different aero-terrestrial habitats, e.g., European tree barks (Kulichová *et al.* 2014; Hallmann *et al.* in prep.-a) and epilithic biofilms (Horath and Bachofen 2009; Hallmann *et al.* 2011b; Ragon *et al.* 2012; Hallmann *et al.* 2013a), Ecuadorean rainforest tree barks and tree leaves (Faßhauer *et al.* 2011; Spitzer *et al.* 2014) or South and Central American sloth hairs (Suutari *et al.* 2010). Central European soils were intensively studied by microscopy-based approach, however, no molecular diversity studies were published so far. Consequently, molecular taxonomy studies on soil Green algae are scant as well. A few genera of Green

algae, which underwent taxonomic revisions, comprise species confirmed in European soils, e.g., *Bracteacoccus* (Fučíková *et al.* 2012), *Chlorella* (Luo *et al.* 2010; Bock *et al.* 2011; Treves *et al.* 2013), *Coccomyxa* (Darienکو *et al.* 2015), *Chloropyrula* (Gaysina *et al.* 2013), *Chlamydomonas* (Pröschold *et al.* 2001), *Chloroidium* (Darienکو *et al.* 2010), *Chromochloris* (Fučíková and Lewis 2012), *Dictyococcus* (Fučíková *et al.* 2011a), *Interfilum* (Mikhailyuk *et al.* 2008), *Klebsormidium* (Škaloud *et al.* 2014a), *Neocystis* (Eliáš *et al.* 2013), *Pseudomuriella* (Fučíková *et al.* 2011a; Fučíková *et al.* 2011b) and *Scenedesmus*-relatives (Lewis and Flechtner 2004; Hegewald *et al.* 2013; Kaufnerová and Eliáš 2013). Some terrestrial microalgae were described elsewhere and re-detected in European soils, but are still in need of taxonomic revisions, e.g., *Stichococcus* (Neustupa *et al.* 2007; Hodač *et al.* subm.), *Jenufa* (Němcová *et al.* 2011; Hodač *et al.* 2012), *Xylochloris* (Neustupa *et al.* 2011; Hodač *et al.* 2012). Unique Green microalgae were so far recorded from North American desert soils, e.g., *Bracteamorpha*, *Rotundella*, *Tumidella* (Fučíková *et al.* 2013), *Desertella*, *Eremochloris*, *Xerochlorella* (Fučíková *et al.* 2014), *Flechtneria* (Sciuto *et al.* 2015) and *Tetraflagellochloris* (Barsanti *et al.* 2013). Green algal species descriptions from African or polar soils are still extremely rare and include *Desmochloris* from Namib Desert (Darienکو *et al.* 2009) or *Pabia* from the Antarctic (Friedl and O'Kelly 2002).

Biogeography of soil Green microalgae

In Green microalgae, unambiguous recognition of the most species is not yet possible, due to a lack of morphological features and because molecular taxonomy do not follow any consensual species concept. Microalgae in terrestrial habitats exhibit morphological convergence (López-Bautista *et al.* 2007), i.e., a tendency to evolve simple globular or filamentous forms throughout unrelated phylogenetic groups (**Fig. 3**). Demanding morphological determination of similarly looking species lead to general underestimation of Green algal diversity and to putative misinterpretations of their distribution patterns (Hoffmann 1989; Rindi *et al.* 2009; Sharma and Rai 2010; Rindi *et al.* 2011; Guiry 2012; Škaloud *et al.* 2015). Whereas in aquatic protists biogeography is widely accepted (Finlay and Clarke 1999; Šlapeta *et al.* 2006; Řezáčová and Neustupa 2007; De Wever *et al.* 2009; Naselli-Flores and Padisák 2015), biogeography of terrestrial Green microalgae is still far from well explored (Lawley *et al.* 2004; Sharma *et al.* 2007; Rindi *et al.* 2009; Leliaert *et al.* 2012; Bates *et al.* 2013; Ryšánek *et al.* 2014). Terrestrial Green microalgae can be encountered nearly everywhere around the globe. Particularly soil species are capable of survival after long periods of desiccation (Trainor 1970; Buzer *et al.* 1985; Trainor 1985; Trainor and Gladych 1995) and can photosynthesize even in cave darkness (Kol 1967). Their resting cysts may easily survive transportation over long distances in the air being

resistant to drought, high as well as low light intensities and high UV radiation, e.g., due to the presence of thickened cell walls or light protection pigments (Řezanka *et al.* 2004; Karsten *et al.* 2005; Häubner *et al.* 2006; Gustavs *et al.* 2010). Early morphology-based attempts comparing geographic distributions of soil microalgae, e.g. Feher (1948), as well as later studies concordantly suggested that a couple of soil algae might be cosmopolitan (Metting 1981; Starks *et al.* 1981; Hoffmann 1989), **Fig. 3**. The putative cosmopolitanism of some microalgae was in accordance with the *neutral dispersal model* (Finlay and Clarke 1999; Finlay 2002; Fenchel and Finlay 2004), which claimed that protists generally do not have biogeography. This view changed rapidly with the era of molecular phylogenies (Sharma and Rai 2010). Molecular markers enabled description of a plethora of morphologically indistinguishable species (Boenigk *et al.* 2005; Rindi *et al.* 2008; Dal Grande *et al.* 2014). As a result, microalgal biogeography seemed to be supported by cryptic species mainly observed within a restricted area. The biogeographic evidence comprises multiple algal lineages such as (1) Stramenopila, e.g., *Spumella* (Boenigk *et al.* 2005), *Xanthonema* (Rybalka *et al.* 2009; Rybalka *et al.* 2013), *Synura* (Škaloud *et al.* 2014b), *Frustulia*




Figure 3. Bar chart summary of the most frequently recorded Green algal morphospecies/morphotypes. Species occurrences were derived from 46 floristic studies (shown in the world map and mentioned on pages 2-3).

(Veselá *et al.* 2012); (2) Streptophyta, e.g., *Micrasterias* (Jurdíková *et al.* 2014), *Klebsormidium*, (Ryšánek *et al.* 2014; Škaloud *et al.* 2014a) and (3) Chlorophyta, e.g., *Asterochloris* (Řídká *et al.* 2014; Škaloud *et al.* 2015). However, more extensive samplings of molecular data from geographically remote regions already confirmed that at least some species of terrestrial Green microalgae are cosmopolitan. A wide geographic distribution might apply for free living species of *Coccomyxa* (Darienkov *et al.* 2015) and *Klebsormidium* (Ryšánek *et al.* 2014), as well as for lichenized species of *Dictyochloropsis* (Dal Grande *et al.* 2014) and *Diplosphaera* (Fontaine *et al.* 2013). In conclusion, biogeography studies on Green microalgae still have to cope with immense lack of molecular diversity data, since only

a few regions have been sampled so far (Novis *et al.* 2008; De Wever *et al.* 2009; Kulichová *et al.* 2014). Species richness of eukaryotic microorganisms is generally poorly understood and diversities revealed from environmental samples significantly differ when using culture-dependent or culture-independent approaches (Moon-van der Staay *et al.* 2006; Doherty *et al.* 2007; Lara *et al.* 2007; Hallmann *et al.* 2009; Hallmann *et al.* 2013a; Spitzer 2013). Particularly in Green microalgae, the link between culture-independent and culture-dependent diversities has been rarely studied. To establish a link between the both approaches, more monoclonal cultures/strains need to be sequenced and phylogenetically characterized. Although there is no universal barcoding marker for identification of microalgae, plastid and mitochondrial markers are widely used, along with the nuclear SSU (e.g. V4/V9 hypervariable regions) and the internal transcribed spacers ITS1-5.8S-ITS2 (Hall *et al.* 2010; Bock *et al.* 2011; Buchheim *et al.* 2011; Fučíková *et al.* 2011b; Krienitz and Bock 2012; Wolf *et al.* 2013; Darienko *et al.* 2015; Darienko and Pröschold 2015; Heeg and Wolf 2015; Lutz *et al.* 2015; Sciuto *et al.* 2015).

General aims

(1) Biodiversity of terrestrial Green microalgae | Chapters 1-2

Traditional morphospecies-based studies revealed a considerable diversity of Green microalgae in terrestrial habitats of Central Europe. Terrestrial Green algae in temperate climate zones, however, are still poorly explored using molecular techniques, in contrast to those from less favorable environments, e.g. hot desert soils. We expect molecular methods to reveal high algal diversities in moist soils of temperate grasslands and forests as well as in submerged algal biofilms. By focusing on cultures of Green microalgae from Central European soils and semi-terrestrial biofilms, we aim to investigate genetic diversity of microscopically hardly distinguishable morphospecies.

(2) Biogeography of terrestrial Green microalgae | Chapters 3-4

Terrestrial Green microalgae are able of long-distance dispersal by resisting harsh environmental stresses. Such 'airborne' microalgae might dwell in soils and terrestrial biofilms across distant geographic regions and climate zones. However, there is little existing molecular evidence to address the question of microbial biogeography in terrestrial Green algae. With focusing on a few exemplar morphogenera, we aim to investigate the widely accepted idea of cosmopolitan distribution of terrestrial Green microalgae.

Summary of the Chapters

Within the frame of this thesis, terrestrial samplings from two main sources were examined. Soil samples originating from 27 grassland and 30 forest sites of the German Biodiversity Exploratories were investigated in **Chapters 1, 3-4**. Semi-terrestrial biofilm samples originating from two karstic streams in Germany were studied in **Chapter 2**. Culturing and sequencing approach was applied consistently throughout the Chapters. Environmental samples were cultivated in standard enrichment media favoring the growth of terrestrial Green algae. After isolating the algae into monoclonal cultures, they were subjected to combined microscopical and molecular phylogenetic analyses. New sequence data were complemented by accessions from GenBank and additional algal strains were kindly provided by the SAG Culture Collection of Algae and by further projects of the Department of Experimental Phycology (Lepka 2007; Spitzer 2013).

Chapter 1 | *Molecular diversity of microscopic Green algae isolated from German soils*

Here we examined the diversity of culturable Green microalgae dwelling in forest and grassland soils. We asked whether our cultured strains genetically match SSU-amplicons (clones, pyrotag reads) from published culture-independent diversity surveys. Our approach enables to link operational taxonomic units from soil metagenomes with physical organisms. We contribute an extensive dataset of sequenced isolates from Central Europe (57 sampling sites, 188 sequenced monoclonal cultures, 61 monophyletic species of Green microalgae). We further inform about new species of soil microalgae and one novel lineage within the class Trebouxiophyceae. By putting geographic data for identical ITS2-ribotypes on the world map, we suggest long-distance dispersal of *Stichococcus*, *Chlorella* and *Klebsormidium*.

Chapter 2 | *Diversity of microscopic Green algae (Chlorophyta) in calcifying biofilms of two karstic streams in Germany*

Green algal biofilms cover periodically desiccating stromatolites in karstic streams in Germany. We investigated this unusual semi-terrestrial habitat for the first time using a combined morphological and molecular approach. Thirty-four species were inferred from 18S rDNA sequence analyses. Among the closest relatives are species known from aquatic and terrestrial habitats. We isolated filamentous Green algae morphologically resembling *Gongrosira* Kützinger, often reported from freshwater tufa-stromatolites. Molecular analysis suggests that the taxon

represents a collective morphotype, which encompasses several genera phylogenetically nested within the Ulvophyceae (Chlorophyta).

Chapter 3 | *Phylogenetic analysis of polar Chlorella and Stichococcus suggests biogeography of airborne microalgae*

Here we aim to uncover biogeographic patterns in cosmopolitan morphospecies *Chlorella* and *Stichococcus*. Already published 18S rDNA sequences from terrestrial and aquatic habitats worldwide, provided a basis for biogeographic analyses. We expanded existing data by new sequences from terrestrial habitats of Germany, Ecuador, the Arctic and Antarctica. We found out that particular *Stichococcus* clades exhibit either temperate-polar or temperate-tropical distribution, but not both. The clades of *Chlorella*-like microalgae which include polar and hot desert strains (e.g., *C. vulgaris* and *Muriella terrestris*) were so far never uncovered in the tropics. Our data provide evidence that airborne microalgae might exhibit biogeography.

Chapter 4 | *Molecular evidence for the wide distribution of two lineages of terrestrial Green algae (Chlorophyta) from tropics to temperate zone*

Terrestrial microalgae *Jenufa* and *Xylochloris* were originally described from the Southeast Asia. Our discovery of similar 18S rDNA sequences in terrestrial samples originating from temperate Europe and South and Central America suggests temperate-pantropic distribution of *Jenufa* and *Xylochloris*. Both lineages occur in soils, but differ in subaerial growth; whereas *Jenufa* grows on rocks and artificial hard substrates, *Xylochloris* was so far detected on organic substrates (tree barks).

Chapter 1

Molecular diversity of microscopic Green algae isolated from German soils

Ladislav Hodač, Christine Hallmann, Karolin Spitzer and Thomas Friedl

University of Göttingen, Experimental Phycology and Culture Collection of Algae (SAG), Göttingen, Germany

(Manuscript draft)

Abstract

Microscopic Green algae are highly diversified in soils of temperate forests and grasslands. The morphology-based literature counts hundreds of (morpho)species, however, only a minor fraction of this diversity has been verified via DNA sequencing of monoclonal cultures. Therefore, SSU-based operational taxonomic units in soil algal metagenomes can hardly be linked with physical organisms. Here we extended the sampling of sequenced algal isolates and cultivated soils from 57 sampling sites in Central European forests and grasslands. We isolated and sequenced 188 monoclonal cultures and retrieved 61 species of Green microalgae. More than a half of them match conspecific cultures isolated elsewhere in Europe, predominantly from terrestrial habitats, but also from freshwaters. Furthermore, 90% of our new cultures match environmental clones retrieved from the same sampling sites and from different terrestrial and aquatic habitats. We contribute new ITS2-based evidence of the long-distance dispersal of *Stichococcus*, *Chlorella* and *Klebsormidium*. On the other hand, at least seven novel species were detected, suggesting that soils in Central Europe might veil algal diversity hardly accessible without culturing.

Keywords: soil biodiversity; Chlorophyta; monoclonal cultures, phylogeny, biogeography, morphology

Abbreviations: MB, Bayesian inference; ML, maximum likelihood; ITS2, internal transcribed spacer 2; OTU, operational taxonomic unit; GBE, German Biodiversity Exploratories; HAI, Hainich; ALB, Schwäbische Alb; SCH, Schorfheide-Chorin

Introduction

Molecular diversity of soil microorganisms is far from well understood (Daniel 2005; Mocali and Benedetti 2010; Shade *et al.* 2012). Nonetheless, investigations of soil prokaryotes successfully target ecological questions (Lauber *et al.* 2009; Nacke *et al.* 2011; Leff *et al.* 2015; Prober *et al.* 2015), accelerated by modern diversity acquisition techniques. In contrast, less attention was paid to the molecular diversity of soil eukaryotic photoautotrophs—microalgae abundant in the topsoil horizon (Starks *et al.* 1981; Hoffmann *et al.* 2007; Wojtuń *et al.* 2013) and generally recognized as primary producers and soil stabilizers (Lukešová 2001; Hu *et al.* 2002; Hu *et al.* 2003a; Hu *et al.* 2003b). Unlike in the case of bacteria (Preston-Mafham *et al.* 2002; Prosser 2002) and fungi (Bailly *et al.* 2007; Prober *et al.* 2015) functional diversity of eukaryotic microalgae in soils remains poorly understood. The prevalent morphology-based concept of ecologically meaningful algocenoses (Gollerbach and Shtina 1969; Zenova *et al.* 1995; Kuzyakhmetov 1998; Şalaru *et al.* 2008; Kabirov *et al.* 2010) urgently needs molecular re-evaluation. Similar to prokaryotes and non-photoautotrophic protists (Boenigk *et al.* 2012; Caron *et al.* 2009; Couradeau *et al.* 2011), molecular diversity of microalgae is underestimated (Guiry 2012; Škaloud *et al.* 2015). Despite considerable sequencing efforts focused on microalgae and other protists in aquatic ecosystems (Moon-van der Staay *et al.* 2001; Chariton *et al.* 2010; Cheung *et al.* 2010; Medinger *et al.* 2010), molecular diversity of soil microalgae remains poorly explored (Lawley *et al.* 2004; Bates *et al.* 2013). Estimates of species richness in soils significantly differ between culture-dependent and culture-independent approaches. This applies for prokaryotes (Rondon *et al.* 2000; Daniel 2005; Shade *et al.* 2012), non-photoautotrophic protists (Moon-van der Staay *et al.* 2006; Doherty *et al.* 2007; Lara *et al.* 2007) and terrestrial microalgae (Hallmann *et al.* 2009; Hallmann *et al.* 2013a; Spitzer 2013). Guiry (2012) estimates that there are approximately 72 500 algal species, 13 000 of these species represent Chlorophyta with 8000 described (cultured) species and 5000 species to be discovered. Even though this estimate relies on Algaebase—the largest online database of information on algae (Guiry and Guiry 2015)—species were verified through molecular methods in a minority of cases.

18S rDNA (SSU) marker once changed our view of microbial diversity (Moreira and Lopez-Garcia 2002), was substantially criticized (Forney *et al.* 2004; Richards and Bass 2005; Škaloud *et al.* 2015), but still holds its significance due to application universality (Lesaulnier *et al.* 2008; Bråte *et al.* 2010; Cheung *et al.* 2010; Bates *et al.* 2013; Lutz *et al.* 2015). However, culture-independent diversity studies often miss a deeper taxonomic resolution and treat Green microalgae as undifferentiated group—Chlorophyta (Lawley *et al.* 2004; Brown *et al.* 2009;

Bates *et al.* 2013; Lutz *et al.* 2015). Number of detected species (or operational taxonomic units, OTUs) directly depends from sequence-similarity threshold-value and sequence length (Caron *et al.* 2009; Grattepanche *et al.* 2014). Eukaryote-wide next generation sequencing (NGS) studies preferred SSU hypervariable regions SSU-V4 (Cheung *et al.* 2010; Taib *et al.* 2013) and SSU-V9 (Amaral-Zettler *et al.* 2009; Taib *et al.* 2013). Considering SSU-V4, 97% sequence similarity is a widely accepted minimum cutoff-value, applied in e.g. Ion Torrent sequencing of snow Chlorophyta (Lutz *et al.* 2015), 454-pyrosequencing of Chlorophyta in an estuary reservoir (Sun *et al.* 2014) or from saltern ponds (Filker *et al.* 2015). The maximum cutoff-value is usually 99%, used in e.g. 454-pyrosequencing of aquatic Chlorophyta (Zhan *et al.* 2013). In biodiversity acquirements, next generation sequencing outperform any traditional approach by means of achieved species/OTU quantity and detection of extremely rare and (so far) uncultivable species (Shi *et al.* 2009; Santoferrara *et al.* 2014; Xiao *et al.* 2014). On the other hand, Xiao *et al.* (2014) noticed limits of NGS for microalgal species resolution and affirmed importance of microscopic species identification. The same recognized Bazin *et al.* (2014) applying 98% cutoff and Viprey *et al.* (2008), who proposed a more stringent threshold for species identification (98.5-99.5%), both cloning marine Green microalgae.

In the taxonomy of Green microalgae, cultures are crucial for precise species characterization (Lukešová 1993; Ettl and Gärtner 1995; Lukešová and Hoffmann 1996; Lukešová 2001). Fine scale identification of microorganisms is essential for understanding their role in ecosystems (Rastogi and Sani 2011) and urgently required for reproducible comparisons of biodiversity shifts between environments (Richards and Bass 2005). Cultures of microalgae are not only valuable for original species characterization, but also for preservation of genetic material (Müller *et al.* 2007a; Day *et al.* 2010). This applies particularly in times of accelerated biodiversity loss (Pimm and Raven 2000). Monitoring of microbial/protistan biodiversity shifts correlated with climate change will hardly be possible without preservation of the current status, e.g., by maintaining cultures or environmental samples (Cary and Fierer 2014; Fierer and Cary 2014). Leading morphology-based identification key for terrestrial algae counts 785 species of (mostly cultured) Green microalgae (Ettl and Gärtner 1995; Ettl and Gärtner 2014). It is not clear how the (morpho)species diversity correlates with OTU-based diversities reported from terrestrial ecosystems via culture-independent techniques. This is due to scarcity of biodiversity surveys combining morphological observations and molecular techniques. Such comparative studies so far focused mainly on phytoplankton (Krienitz and Bock 2012), semi-terrestrial biofilms (Brinkmann *et al.* 2015; Hodač *et al.* 2015) and aerophytic biofilms (Lin *et al.* 2012; Kulichová *et al.* 2014). Regarding soils—the most species-rich terrestrial habitats—polyphasic data are

available so far for arid lands (Lin and Wu 2014; Flechtner et al. 2013; Fučíková et al. 2014) or subtropical farmlands (Lin *et al.* 2013).

In the present study, for the first time we investigate molecular diversity of culturable Green microalgae in soils from Central Europe. The soils of this region remain almost untouched by culture-dependent molecular diversity surveys targeting Green microalgae, despite the fact that significant fraction of them were described from here (Ettl and Gärtner 1995). Expecting differences in cultured and uncultured microbial diversities, we hypothesize that extensive sampling and culturing would retrieve new species, not yet submitted to GenBank from culture-independent surveys. We investigate 188 new monoclonal cultures of soil Green microalgae sampled from 57 sites covering deciduous and needle forests and meadows and pastures in Germany (Fischer *et al.* 2010). To achieve comparability with culture-independent studies, we sequenced SSU marker and conducted species identification following monophyletic species concept (Johansen and Casamatta 2005; Mallet 2005; Vyverman *et al.* 2010; Škaloud and Rindi 2013; Leliaert *et al.* 2014). Allowing minimal sequence divergence—using high sequence similarity threshold of $\geq 99.90\%$ —we aim to evaluate how our data match already published cultured species. In a second step we test whether our cultured species match OTU-based diversities acquired by cloning or NGS techniques, hereby applying a similarity threshold of 97% for pyrotags (Sun *et al.* 2014; Lutz *et al.* 2015) and 99% for clones (Viprey *et al.* 2008; Zhan *et al.* 2013). Since soil is a semi-aquatic habitat, it presumably hosts an assemblage of terrestrial and aquatic species. Though morphological literature suggest cosmopolitanism of microalgae in temperate soils (Ettl and Gärtner 1995; Lukešová 2001), the molecular evidence is missing. This we aim to prove by comparing our isolates with cultures sampled elsewhere, analyzing the highly variable ITS2 marker (Müller *et al.* 2007b; Coleman 2009; Buchheim *et al.* 2011; Keller 2011; Caisová *et al.* 2013; Wolf *et al.* 2013).

Material and Methods

Sampling sites

Soil samples were collected from defined plots within large-scale and long-term research sites termed Biodiversity Exploratories (Hainich-Dün, Abbr. HAI; Schorfheide-Chorin, Abbr. SCH; Schwäbische Alb; Abbr. ALB; **Fig. 1a**). The sampling covers plots in forests (**Figs. 1b, 1c, 1d**) and grasslands (**Figs. 1e, 1f**), which represent a gradient of land-use intensities, described in detail by Fischer *et al.* (2010). We analyzed samples from two soil horizons: 1) drill core (A-horizon) samples were collected in HAI, SCH, ALB in April 2008 from 27 grassland and 30 forest plots

and multiple samples from within each plot were pooled, 2) soil surface (O-horizon, upper 3 cm) samples were taken in autumn (September 2010) and spring (March 2011, June 2011) from two forest and three grassland plots within the HAI exploratory (using a sterile spoon and transported in sterile Petri dishes. In the latter case, three different sampling points were defined within each plot (middle, north-east corner, south-west corner; **Fig. S5e**). These subplot samples were analyzed in two ways, i.e., separately and pooled together.




Figure 1. Soil sampling in the German Biodiversity Exploratories. (a) Localization of the three sampling areas (highlighted in red): ALB=Schwäbische Alb, HAI=Hainich-Dün, SCH=Schorfheide-Chorin). (b)-(c) Examples of sampling sites in extensively managed (b) and intensively managed (c) forests (Hainich). (d) Soil sample taken by a spoon from topsoil. (e)-(f) Examples of sampling sites in extensively managed (e) and intensively managed (f) grasslands (Hainich). (g)-(i) Culturing techniques—establishment of mixed cultures. (g) Algal colonies on agar plate with spread soil suspension; (h) Algal growth in soil suspended in liquid BBM medium (Bischoff and Bold 1963); (i) Culturing of microalgae on coverslips put on the soil surface.

Soil culturing, isolation and light microscopy of monoclonal cultures

Two grams of each soil sample were suspended and cultured in liquid media and spread on agarized media (**Fig. 1g**). The composition of liquid (in 100 ml Erlenmeyer flasks; **Fig. 1h**) and 1.5% agarized (Petri dishes) media was the same: 3NBBM+V and BG11 (EPSAG; <https://www.uni-goettingen.de/de/list-of-media-and-recipes/186449.html>). Both liquid and

agarized cultures were kept for at least six weeks at 18 °C under a light : dark regime of 14 : 10 hours at a light intensity of about 25 μ E from a white fluorescent bulb. After three, four and six weeks of growth in liquid media, algal suspension was inspected using a light microscope. The algal colonies growing on the agarized plates were transferred with sterile inoculating loops onto new fresh plates with the same medium and final monoclonal cultures were kept on agar slants (1.5%) under the same growth conditions. The soil surface samples were treated in the same way as described above. The only modification concerns the BG11 medium which was replaced by Z medium. In addition to the culturing techniques on media, cover slips method was used (Ettl and Gärtner 1995; Lukešová 2001). Sterile glass cover slips were put directly on wet soil samples (stored in sterile Petri dishes; **Fig. 1i**) and after 2-3 weeks of algal growth on the contact surface, the covers slips were analyzed by light microscopy. The light microscopy of the mixed raw cultures and unialgal isolates was accomplished using an Olympus BX60 microscope (Tokyo, Japan) with Nomarski DIC optics with an attached ColorView III camera (Soft Imaging System, Münster, Germany). The observations were documented by micrographs, processed using the Cell[^]D image software (Soft Imaging System, Münster, Germany). Cell size measurements (diameter of spherical cells; length/width of elongated cells) were conducted were conducted in program ImageJ (Schneider *et al.* 2012) and were based on 100 cells per culture. and the summary statistics including maximum and minimum values were computed in PAST 2.12 (Hammer *et al.* 2001).

DNA isolation, PCR and sequencing

DNA was extracted from 188 monoclonal isolates (**Table S1**) using the Invisorb Spin Plant Kit (Strattec molecular, Berlin, Germany) as recommended by the manufacturer. 18S rDNA marker was amplified using primers NS1 and 18L (Hamby *et al.* 1988). For selected isolates 18S and ITS2 rDNA gene region was amplified using primers NS1 and LR1850 (Friedl 1996). Conditions for PCR and sequencing reactions and the standard set of sequencing primers were described previously by Mikhailyuk *et al.* (2008). The newly determined sequences will be deposited in GenBank. Two additional isolates from polar regions (Shukla *et al.* 2011), and two from Ecuador (Spitzer 2013; Faßhauer in prep.) were sequenced for 18S rDNA.

Phylogenetic analysis

The newly obtained 80 full to almost full 18S rDNA sequences (~1600-1800 nucleotides in length; **Tables S2a, S2b**) were compared to NCBI database using BLAST queries (Altschul *et al.*

1997) in order to download all closest relatives available from GenBank (Benson *et al.* 2012). All sequences were checked for chimeras by Bellerophon (Huber *et al.* 2004). Two alignments were established, one for Green algae (Chlorophyta + Streptophyta) and another for Stramenopiles including Xanthophyceae and Eustigmatophyceae. Both alignments were further processed in the same way. Alignments were computed by the program MAFFT-Q (Kato and Toh 2008) and inspected in BioEdit (Hall 1999) for possibly misaligned nucleotide positions. The alignment of Green algae consisted of 169 sequences with 1817 positions (711/564 variable/parsimony informative). The alignment of Stramenopiles consisted of 72 sequences with 1895 positions (613/434 variable/parsimony informative). In both cases, the GTR+ Γ +I model was selected as the best fitting substitution model using AIC criterion in jModelTest 0.1.1 (Posada 2008). Phylogenetic trees were calculated using maximum likelihood method in the program RAxML 7.0.4 (Stamatakis *et al.* 2008). The Bayesian phylogenetic trees were calculated using MrBayes 3.2.1 x64 (Ronquist *et al.* 2012). For the latter, two MCMC runs for three million generations each were employed with one cold and three heated chains with trees sampled every 100 generations. Internal edge support values were inferred from the rapid bootstrapping algorithm (100 replicates) as implemented in RAxML (Stamatakis *et al.* 2008) and from Bayesian posterior probabilities using MrBayes 3.2.1 x64 (Ronquist *et al.* 2012). Pairwise sequence similarities were derived from p-distances based on Kimura-2-parameter model computed in the program MEGA5 (Tamura *et al.* 2011). Pairwise similarities among our isolates and their closest GenBank relatives are listed in **Table S3**. The same method was used to identify species represented by partial 18S rDNA sequences (**Tables S4a, S4b**).

Delimitation of morphotypes and statistical analysis of their distribution

The morphological diversity of monoclonal cultures is summarized in **Table S5**. Observations on morphotypes occurring in mixed cultures (agar plates, liquid media and cover slips) were documented as binary presence/absence matrices (**Table S6a; Table S6b**) and summarized in **Table S7**. Delimitation of morphotypes was accomplished by identification literature (Reisigl 1964; Trenkwalder 1975; Metting 1981; Huber-Pestalozzi *et al.* 1983; Ettl and Gärtner 1995; Andreyeva 1998; Komárek and Anagnostidis 1998; Komárek and Anagnostidis 2005; Hoffmann *et al.* 2007; Škaloud 2009; Komárek 2013; Ettl and Gärtner 2014). Morphotype representatives are documented as drawings of monoclonal cultures and microphotographs of monoclonal cultures and mixed cultures. Where it was possible, taxa were morphologically determined at the genus level. Otherwise, assignation at higher taxonomic level was preferred. Taxonomically

ambiguous taxa predominantly occurring as similar living forms were combined into collective morphotypes. For example, sarcinoid forms traditionally named *Chlorosarcina*, *Chlorosarcinopsis* and *Desmotetra* represented one collective morphotype and named Chlorosarcinaceae. The resulting binary presence/absence matrix was further used for multivariate statistical analysis. Prior to the analysis, we pooled the presence/absence data of each three replicates representing the same land-use intensity, e.g., AEG1+AEG2+AEG3=AEG_i (intensively managed grassland plots within ALB). Redundancy analysis (RDA) (Hill 1979; Jaarsma *et al.* 2006) was conducted to test for correlations between important soil physico-chemical parameters, i.e., pH, organic carbon/C_{org}, total nitrogen/N_{tot} (Metting 1981; Marschner *et al.* 2003) and the composition of algal morphotypes at the sampling sites. The original measurement data were obtained from Nadine Herold in the frame of the German Biodiversity Exploratories project. Because the diversity data were pooled within each land-use category (Fischer *et al.* 2010), in the ordination analysis we used corresponding arithmetic means of each physico-chemical parameter. For statistical support the Monte Carlo test with 2000 permutations under reduced model was used (Hope 1968). The morphotype composition of the soil surface samples was analyzed using Non-metric Multidimensional Scaling (NMDS) using Jaccard distance measure. All statistical analyses were performed in PAST version 2.12 (Hammer *et al.* 2001) and CANOCO version 4.5 (ter Braak and Šmilauer 2002). Hence Green algae (Chlorophyta, Streptophyta) and stramenopiles (Xanthophyceae, Eustigmatophyceae) were present in almost all samples and many taxa are relatively easy to cultivate (compared to e.g. Diatoms or Cyanobacteria) we focused on them in further molecular studies.

Results

Molecular diversity and morphology of the monoclonal isolates

We analyzed a total of 188 18S sequences; 146 sequences were longer (1600-1800 bp) and 42 shorter (300-1600 bp). Within this dataset we distinguished different 73 phylotypes corresponding to species at $\geq 99.90\%$ sequence similarity level. The species belong mainly to Green algae: Chlorophyta (59 spp.) and Streptophyta (2 spp.) and to Stramenopiles: Xanthophyceae (11 spp.) and Eustigmatophyceae (1 sp.).

Green algae

We detected 61 species assigned of 21 lineages from four classes of Chlorophyta (Chloro-, Trebouxioid-, Ulvo- and Prasinophyceae) and one class of Streptophyta (Klebsormidiophyceae).

The Chlorophyceae are represented by ten lineages: *Oogamochlamydia*, *Chloromonadinia*, *Tatrensinia*, *Reinhardtina*, *Desmotetra*, *Stephanosphaerina*, *Jenufa*, *Scenedesmaceae*, *Pseudomuriellaceae*, *Bracteacoccaceae*). The Trebouxiophyceae are represented by eight lineages: *Prasiola/Stichococcus*, *Neocystis*, *Botryococcus*, *Watanabea*, *Lobosphaera*, *Xylochloris/Dictyochloropsis*, '*Navichloris*', *Chlorellaceae*. The Ulvophyceae are represented by Ulotrichales and Pedinophyceae by *Pedinomonas*. The Klebsormidiophyceae is represented by *Klebsormidium*.

Streptophyta; Klebsormidiophyceae

The Streptophyta were represented by two different species of *Klebsormidium* (**Fig. 2**), common terrestrial algae with unbranched filaments and a single band-shaped chloroplast (with a pyrenoid) per cell. (I) *Klebsormidium flaccidum* (e.g., isolate LH10HG2056; **Fig. 3a**) from grasslands (HAI, SCH) and forest (HAI) is identical to *K. flaccidum* SAG 7.91 (EU434019) from freshwater (**Fig. 2**). *Klebsormidium* cf. *dissectum/elegans* (isolate LH08HW9106; **Fig. 3b**) from grasslands (ALB) and forests (HAI) is identical to authentic strain *K. dissectum* SAG 2155 (from soil) and to authentic strain *K. elegans* SAG 7.96 (from tree bark). The whole clade including species of *Klebsormidium* and *Interfilum* is highly supported.

Chlorophyta; Pedinophyceae

Prasinophyte algae were represented by one isolate belonging to *Pedinomonas* (**Fig. 2**), exhibiting small *Chlamydomonas*-like cells containing one pyrenoid and basal stigma. *Pedinomonas* cf. *minor* (isolate LH08SG2033; **Fig. 3c**) from grassland (SCH) is closely related to *P. minor* SAG 1965-3 (HE610132) from freshwater (**Fig. 2**). The clade comprising both accessions is highly supported.

Chlorophyta; Ulvophyceae

The class Ulvophyceae is here represented by the Ulvales/Ulotrichales lineage (Watanabe and Nakayama 2007) and one genus *Pseudendocloniopsis* (**Fig. 2**), with spherical to ovoid (slightly acuminate) cells with a cup-shaped chloroplast and prominent pyrenoids. *Pseudendoclonium* cf. *botryoides* (isolate LH08HW9058; **Fig. 3d**) from forest (HAI) was closely related to authentic strain *P. botryoides* SAG 465-1 (AJ416103) from freshwater. The clade comprising both accessions is highly supported.

Chlorophyta; Chlorophyceae; Sphaeropleales.

Scenedesmaceae (Hegewald and Hanagata 2000) comprised three species (**Fig. 2**) with spherical to symmetrically elliptic cells with slightly acuminate poles. (I) *Coelastrella multistriata* (e.g., isolate LH10HG7083; **Fig. 3e**), identical to terrestrial *C. multistriata* Hanagata C6-2 (AB012846), was found in grasslands (HAI, ALB) and forest (ALB). (II) *Coelastrella* sp. (e.g., isolates LH10HG2098, LH10HG7018; **Fig. 3f**) occurs in grasslands (HAI) and is identical to *Scenedesmus* sp. KGU Y002 (AB742453). (III) *Acutodesmus rubescens* (e.g., isolate LH08SG8041; **Fig. 3g**) from grassland (SCH) is identical to '*Scenedesmus*' (*Acutodesmus*) *rubescens* CCAP 232/1 (X74002). The Scenedesmaceae clade is highly supported.

Members of Pseudomuriellaceae and Bracteacoccaceae (Fučíková *et al.* 2013) exhibit spherical cells with several to multiple parietal chloroplasts without pyrenoids. *Pseudomuriella aurantiaca* (isolates LH10HG9038, LH10HG2039; **Fig. 3h**) from grasslands (HAI) is identical to the authentic strain *P. aurantiaca* SAG 249-1 (X91268) isolated from soil. *Bracteacoccus cohaerens* (isolate LH10HG9034; **Fig. 3i**) from grassland (HAI) is identical to the authentic strain *B. cohaerens* UTEX 1272 (GQ985406) from soil (USA). Another isolate from grassland (SCH), *Bracteacoccus* cf. *cohaerens* (isolate LH08SG2015), is slightly different from the same GenBank accession but might be as well conspecific. Both *Pseudomuriella* and *Bracteacoccus* clades were highly supported.

Chlorophyta; Chlorophyceae; Chlamydomonadales

Isolates assigned to Chlorophyceae were nested in two major lineages Sphaeropleales and Chlamydomonadales, and *Jenufa* clade incertae sedis (**Fig. 2**). The Chlamydomonadales isolates were characteristic for monadoid/flagellate *Chlamydomonas*-like and/or coccal *Chlorococcum*-like morphologies (**Fig. 4a-k**). The Sphaeropleales were either coccal colonial *Scenedesmus*-like, or non-colonial *Bracteacoccus*-like (**Fig. 4a-e**).

Oogamochlamydia clade (Nakada *et al.* 2008). The clade comprises monadoid isolates belonging to two species of similar morphology differing in cell size. (I) *Oogamochlamys* sp.(I) (isolate LH08SG8047; **Fig. 4a**) from grassland (SCH) is less closely related to *O. ettliei* UTEX 2218 and *O. gigantea* SAG 21.72. (II) *Oogamochlamys* sp.(II) strain SAG 2476 (= isolate LH08AW1069); **Fig. 4b**) from forest (ALB) is less closely related to freshwater *Chlamydomonas* sp. CCAP 11/159. Both species are nested in well supported clade including *Oogamochlamys* and *Lobochlamys* accessions.

Chloromonadinia clade (Nakada *et al.* 2008). One monadoid species, *Chlamydomonas* cf. *gerloffii* (isolate LH08SW5031; **Fig. 4c**), found in forest (SCH), is closely related to freshwater

C. gerloffii CCAP 11/72 (FR865610). The species is nested in a highly supported clade including authentic strains *C. gerloffii* CCAP 11/72 and *Chloromonas rosae* UTEX 1337 accessions.

Reinhardtinia clade (Nakada *et al.* 2008). Monadoid isolates of *Chlamydomonas*-like morphology clustered in four different species. (I) *Chlamydomonas rapa* (isolates LH08SG1077, LH08SG9055) detected in grasslands (SCH), is identical to freshwater authentic strain *C. rapa* SAG 48.72 (U70790). (II) *Chlamydomonas* cf. *rapa* (isolate LH10HG1027; **Fig. 4d**) from grassland (HAI), is very closely related to *C. rapa* SAG 48.72 and might be conspecific as well. These three accessions create a highly supported clade, a sister group to *C. reinhardtii* (M32703). Two isolates from grasslands (ALB/SCH) were nested in another highly supported subclade within *Reinhardtinia* clade. (III) *Heterochlamydomonas* sp. (isolate LH08AG2004; **Fig. 4e**) closely related to freshwater *H. rugosa* SAG 45.86 (AF367859) and authentic strain *H. inaequalis* UTEX 1705 (AF367857). (IV) *Chlamydomonas* cf. *typica* (isolate LH08SG9022; **Fig. 4f**) is closely related to soil *C. typica* NIES-2246 (AB701557). The clade including the four species from Germany is highly supported.

Desmotetra clade (Nakada *et al.* 2008). *Desmotetra* Deason & Floyd detected in this study is characterized by pairs of widely elliptic non-motile cells remaining attached after division. *D. stigmatica* (isolate LH08SG2049; **Fig. 4g**) from grassland (SCH) is identical to authentic strain *D. stigmatica* UTEX B 962 (DQ009760) in a highly supported clade – a sister group of the *Reinhardtinia* clade. The species was further found in grassland (HAI; isolate LH10HG6P18).

Tatrensinia clade (Nakada *et al.* 2008). The clade comprises two *Chlorococcum*-like species from grasslands (HAI/SCH). (I) '*Tatrensinia*' ('*Chlorococcum*') sp.1 (LH08SW7115; **Fig. 4h**) is closely related to soil Haematococcaceae clone Amb_18S_582 (EF023273). (II) '*Tatrensinia*' ('*Chlorococcum*') sp.2 (isolates LH10HG7016, LH10HG9131; **Fig. 4i**) is closely related to '*Chlorococcum*' cf. *tatrense* (= *Chlamydocapsa* sp.) CCCryo 101-99 (AF514407) from Spitsbergen permafrost. Both species are nested in a highly supported clade.

Stephanosphaerinia clade (Nakada *et al.* 2008). Isolates exhibiting monadoid *Chlamydomonas*-like morphology clustered in four different species. (I) *Chlorococcum sphacosum* ('*Neospongiococcum*' *gelatinosum*) (isolate LH10HG3113; **Fig. 4j**) is identical to authentic strain *C. sphacosum* SAG 66.80 (JN968580) and authentic strain *N. gelatinosum* SAG 64.80 (JN968584), both isolated from soils. (II) *Chlamydropodium vacuolatum* (isolate LH10HG1013) identical to '*Chlorococcum*' *robustum* Kr 1986/30 (AY122332) from freshwater. (III) '*Stephanosphaerinia*' ('*Chlorococcum*') sp. (isolate LH10HG6108), is closely related to freshwater isolate *Chlorococcales* sp. VII3 (FJ946904) from Antarctica. The three species are nested in a highly supported clade. Another highly supported clade consisted of (IV)

Chlorococcum cf. *minutum* SAG 2479 (= LH08AW5056; **Fig. 4k**) and closely related *C. minutum* SAG 21.95 (JN968585) from soil. *Chlorococcum* cf. *minutum* was detected in grassland and forest (ALB).




Figure 2. 18S rDNA phylogeny of Green algae, part 1 including Chlorophyceae, Ulvophyceae, Pedinophyceae and Klebsormidiophyceae. New accessions are written in bold and grey underlined. Black dots mark references to microphotographs (**Fig. 3**, **Fig. 4**) sorted in top down order. Original drawings of representative morphologies are shown for each clade (scale bars = 5 μ m). Sequences of authentic strains are marked by a 'c' sign. Numbers next to branches indicate statistical support values (maximum-likelihood bootstraps/Bayesian posterior probabilities).

Chlorophyta; Chlorophyceae; *Jenufa* clade

Jenufa clade (Němcová *et al.* 2011) comprises coccid species with spherical cells with an extensively lobed parietal chloroplasts without pyrenoids. *Jenufa* sp. SAG 2383 (= LH08AW8035; **Fig. 4l**; JQ988933) and (identical) isolate LH08AW8098, found in forest (ALB), are related to authentic strain *J. minuta* CAUP H 8102 (HM563744) from tree bark in Singapore. The *Jenufa* clade, incertae sedis within the Chlorophyceae, consists of *J. minuta* CAUP H 8102, *J. perforata* CAUP H 8101 and *Jenufa* sp. SAG 2383 (= LH08AW8035) and is highly supported.




Figure 3. Microphotographs of monoclonal cultures of Green algae (Klebsormidiophyceae, Prasinophyceae, Ulvophyceae and Chlorophyceae/Sphaeropleales). (a) *Klebsormidium* cf. *flaccidum* LH10HG2056 (l=6.4-19.7 μ m, w=4.5-7.0 μ m); (b) *Klebsormidium* cf. *dissectum* LH08HW9106; (c) *Pedinomonas minor* LH08SG2033 (l=2.9-5.6 μ m, w=2.5-4.8 μ m); (d) *Pseudendocloniopsis botryoides* LH08HW9058 (\varnothing =4.7-9.5 μ m); (e) *Coelastrella multistriata* LH10HG7083 (\varnothing =3.5-7.0 μ m); (f) *Coelastrella* sp. LH10HG7018; (g) *Acutodesmus rubescens* LH08SG8041; (h) *Pseudomuriella aurantiaca* LH10HG2039 (\varnothing =3.6-8.2 μ m); (i) *Bracteacoccus cohaerens* LH10HG9034 (\varnothing =5.4-12.3 μ m). Scale bars = 10 μ m.




Figure 4. Microphotographs of monoclonal cultures of Green algae (Chlorophyceae/Chlamydomonadales). (a) *Oogamochlamys* sp.(I) LH08SG8047; (b) *Oogamochlamys* sp.(II) SAG 2476 (= LH08AW1069; l=9.1-14.5 μ m, w=5.1-10.0 μ m); (c) *Chlamydomonas* cf. *gerloffii* LH08SW5031 (l=10.9-16.4 μ m, w=4.3-10.1 μ m); (d) *Chlamydomonas* cf. *rapa* LH10HG1027 (l=6.1-9.9 μ m, w=2.7-5.1 μ m); (e) *Heterochlamydomonas* sp. LH08AG2004 (l=6.0-10.5 μ m, w=3.1-7.5 μ m); (f) *Chlamydomonas* cf. *typica* LH08SG9022 (l=6.7-12.2 μ m, w=4.0-8.0 μ m); (g) *Desmotetra stigmatica* LH08SG2049 (\varnothing =3.5-6.6 μ m); (h) *Tatrensinia* sp.(I) LH08SW7115 (l=8.3-17.8 μ m, w=5.2-13.3 μ m); (i) *Tatrensinia* sp.(II) LH10HG9131 (\varnothing =7.9-13.1 μ m); (j) *Chlorococcum sphacosum* LH10HG3113 (l=5.8-9.3 μ m, w=2.6-6.3 μ m); (k) *Chlorococcum* cf. *minutum* SAG 2479 (= LH08AW5056; l=5.6-9.5 μ m, w=5.6-9.5 μ m); (l) *Jenufa* sp. SAG 2383 (= LH08AW8035; \varnothing =2.9-7.3 μ m). Scale bars = 10 μ m.

Chlorophyta; Trebouxiophyceae

Isolates assigned to Trebouxiophyceae are phylogenetically nested in eight different clades (**Fig. 5**) and comprised rod-shaped, spherical and elliptic morphologies (**Fig. 6**).

Stichococcus-like rod-shaped isolates were spread across the *Prasiola/Stichococcus* clade (Sluiman and Guihal 2008) and we distinguished at least nine different species. (I) '*Stichococcus*' sp. (isolate LH08AW8025; **Fig. 6a**) from forest (ALB) is identical to '*Stichococcus*' sp. SAG 2059 (AY762604) from a building façade. (II) '*Stichococcus*' sp. (isolate LH08SG5057; **Fig. 6b**), occurring in grasslands (HAI, SCH, ALB), is identical to '*Stichococcus*' sp. SAG 2060 (AY762606) from a building façade. These species clustered near to authentic strain *Stichococcus deasonii* UTEX 1706 (DQ275460) from soil, yet the clade is not supported. (III) '*Stichococcus*' sp. (isolates LH08AG7010, LH10HG6110) from grasslands (HAI, ALB) is closely related to authentic strain *Stichococcus deasonii* UTEX 1706 (DQ275460). (IV) '*Stichococcus*' sp. (isolate LH08SG1073) from grassland (SCH) is identical to '*Stichococcus*' sp. SAG 2406 (= WB47; KF144240) from Westerhöfer creek. (V) '*Stichococcus*' sp. (isolate LH08SW1099) from forest (SCH) clustered near to the aforementioned species (IV) in an unsupported clade. (VI) '*Stichococcus*' sp. SAG 2482 (= LH08AW8023; **Fig. 6c**) from forest (ALB) is identical to '*Stichococcus*' sp. D4-2A (KF144238) from Deinschwanger creek. (VII) *Stichococcus* sp. SAG 2481 (= LH08AW8002) from forest (ALB) is identical to *Stichococcus* sp. K4-4 (AB055866) of unknown origin (probably isolated in Japan). *Pseudostichococcus* is represented by two species exhibiting rod-shaped *Stichococcus*-like morphology. (I) *Pseudostichococcus* cf. *monallantoides* (e.g., isolate LH10HG3045), detected in grasslands (HAI, ALB), is closely related to authentic strain *P. monallantoides* SAG 380-1 (JX185690). (II) *Pseudostichococcus* sp. (isolate LH08SW8044) from forest (SCH) is identical to '*Stichococcus*' *mirabilis* CCAP 379/3 (AJ311638). The clade comprising both *Pseudostichococcus* species and '*Stichococcus*' *mirabilis* CCAP 379/3 consistently occurred in both maximum-likelihood and Bayesian phylogenies, yet without statistical support. *Diplosphaera* is represented by two species exhibiting spherical to slightly cylindrical (shortly rod-shaped) cells which characteristically remain in doublets after cell division. (I) '*Chlorella*' cf. *sphaerica* (isolate LH08HW8075) from forest (HAI) is very closely related to authentic strain *C. sphaerica* SAG 11.88 (AJ416105) from New Zealand. Both specimens might be conspecific. Another isolate (LH08AG9089) from grassland (ALB) is as well closely related to the same authentic strain with a slightly higher genetic distance. The clade including both species (*Diplosphaera* clade) is well supported.

Elliptic morphotypes. Elliptic cell forms were characteristic for phylogenetically unrelated lineages *Neocystis* (Eliáš *et al.* 2013), *Coccomyxa* (Darienکو *et al.* 2015), *Chloroidium* (Darienکو

et al. 2010) and '*Navichloris*' (this study). The isolates from grasslands (HAI, SCH, ALB) and forests (HAI, ALB) belonging to *Neocystis brevis* (e.g., LH10HG9080; **Fig. 6d**) are identical to authentic strain *N. brevis* CAUP D 802 (JQ920360) from soil (the *Neocystis* clade was highly supported). *Coccomyxa* was represented by three species. (I) *C. simplex* (isolate LH08SG9051; **Fig. 6e**) from grassland (SCH) is identical to authentic strain *C. rayssiae* SAG 216-8 (= UTEX 273; HQ317304) from freshwater. The clade further includes authentic strain *C. chodatii* SAG 216-2 (FJ648512) and is highly supported. (II) *Coccomyxa viridis* SAG 2483 (= LH08AW8039; **Fig. 6f**), from forest (ALB), is identical to lichenized *C. glaronensis* CCALA 306 (AM167525). The clade comprises as well authentic strain *C. mucigena* SAG 216-4 (FJ648513) and is highly supported. (III) *Coccomyxa* sp. (isolate LH08AW1017) from forest (ALB) is very closely related to lichenized *Coccomyxa* sp. KN-2011-T3 (HE586515) from Indonesia. The clade additionally includes Uncultured Trebouxiophyceae RL75K2 (HE617184) from an acidic mining lake in Germany, and is highly supported. The strain SAG 2477 (= LH08AW3007; **Fig. 6g**; KP081399) detected in forest (ALB), exhibits symmetrically elliptic cells resembling *Coccomyxa* Schmidle. The species is neither phylogenetically related to *Coccomyxa* nor to other known genera and clades of Trebouxiophyceae. The closest BLAST-hit *Leptochlorella* sp. clone Qe17 (FJ790649) from China exhibits low similarity (= 95.54%). In both maximum-likelihood and bayesian phylogenies, SAG 2477 consistently clusters near to the *Leptochlorella* clade (including authentic strain *Leptochlorella corticola* CAUP H8401; HE984579), yet without statistical support. The isolates of *Chloroidium*, exhibiting widely elliptic cells with a band-shaped lobed chloroplast with a pyrenoid, were assigned to two species. (I) *C. saccharophilum* (e.g., isolate LH10HG7062; **Fig. 6h**), detected in grassland (HAI) and forest (ALB), is identical to authentic strain *C. saccharophilum* SAG 211-9a (FM946000) from tree sap. *Chloroidium* cf. *ellipsoideum* (e.g., isolate LH10HG9105; **Fig. 6i**), detected in grasslands (HAI, ALB), is closely related to authentic strain *C. ellipsoideum* SAG 3.95 (FM946012). The clade comprising both species is highly supported.

Coccoid isolates exhibiting predominantly spherical vegetative cells comprised phylogenetically divergent lineages *Lobosphaera* (Neustupa *et al.* 2011), *Xylochloris* (Neustupa *et al.* 2011), *Dictyochloropsis* (Dal Grande *et al.* 2014), *Prasiola* clade ('*Chlorella*' *mirabilis*) and members of the Chlorellaceae (Krienitz *et al.* 2004). The isolates assigned to *Lobosphaera* exhibited spherical cells with one parietal chloroplast without a pyrenoid and could be phylogenetically classified in two species. (I) *Lobosphaera* cf. *irregularis* (isolates LH08SW5063, LH08AW3064; KP081398) from forests (ALB, SCH) is identical to '*Myrmecia*' (= *Lobosphaera*) *irregularis* CCAP 221/8 (HQ902935). (II) *Lobosphaera* cf. *bisecta* (isolate

LH10HG3P15; **Fig. 6j**) from grassland (HAI) is identical to authentic strain *Myrmecia* (= *Lobosphaera*) *bisecta* SAG 2043 (Z47209) from soil (Italy). The *Lobosphaera* clade (including as well '*Parietochloris*' *cohaerens* UTEX 1707; EU878372) is highly supported. *Xylochloris* sp. SAG 2382 (= LH08AG7024; **Fig. 6k**; JQ988942) from grassland (ALB), with characteristic parietal star-like lobed chloroplasts, is less closely related (similarity = 96.47%) to authentic strain *X. irregularis* CAUP H7801 (EU105209) from tree bark (Singapore). The clade including both *Xylochloris* accessions is only partly supported and the new *Xylochloris* sp. SAG 2382 is likely to represent new taxon on at least species or even genus level. Closely related genus *Dictyochloropsis* comprises *D. splendida* (isolate LH08AW3050; **Fig. 6l**; JQ988930) from forest (ALB), identical to *D. splendida* CAUP H8601 (GU017662) from soil near to a fumarole (Czech Republic). *Dictyochloropsis* and *Xylochloris* species built a well-supported clade. *Chlorella*-like isolates from the *Prasiola* clade, phylogenetically assigned to '*Chlorella*' *mirabilis*, were characterized by spherical cells with one parietal chloroplast containing characteristic small starch granules over the pyrenoid circumference. The isolates from grasslands in ALB (isolate LH08AG9040; **Fig. 7a**) and HAI (isolate LH10HG6139) are closely related to '*Chlorella*' *mirabilis* Andreyeva 748-I (X74000) from soil and might represent two different species. The clade including both isolates and '*Chlorella*' *mirabilis* Andreyeva 748-I is highly supported.

Chlorellaceae were represented by four different morphological types, i.e., *Chlorella*-like (spherical to slightly elliptic cells with cup-shaped chloroplast with one pyrenoid), *Auxenochlorella*-like (spherical cells with facultatively degraded chloroplasts; heterotrophic), *Muriella*-like (bigger spherical cells with single parietal chloroplast without a pyrenoid), *Nannochloris*-like (smaller spherical cells with single parietal chloroplast without a pyrenoid). *Chlorella vulgaris* isolates were sampled in grasslands (HAI, SCH) and forests (HAI). For example, *C. vulgaris* isolates LH08SG3006 and LH08HG1081 (**Fig. 7b**) were identical (similarity = 99.94-100%) to authentic strain of *C. vulgaris* SAG 211-11b (FM205832) from freshwater (Netherlands). *Chlorella* cf. *vulgaris* isolate LH10HG2081 (**Fig. 7c**) is closely related to *C. vulgaris* SAG 211-11b and identical to isolates L1 and L4 from soil (King George Island, Antarctica). The Antarctic strains (and isolate LH10HG2081) constitute a well-supported clade and might represent a new *Chlorella* species. *Muriella* cf. *terrestris* isolates LH08SG3009 and LH10HG7118 (**Fig. 7d**) from grasslands (HAI, SCH) are almost identical (similarity = 99.94%) to *Muriella terrestris* ASIB V38 (AB012845) from soil (Italy), within a highly supported clade. *Nannochloris*-like isolates clustered as four different species. (I) *Nannochloris* sp. isolates from grasslands (e.g., LH08SG8030, **Fig. 7e**; LH10HG6095, **Fig. 7f**) are almost identical to *N. bacillaris* (AB080300) within a highly supported clade. The remaining *Nannochloris*-like isolates

clustered within the highly supported Unidentified Chlorellaceae clade, a sister group to *N. bacillaris* (AB080300) and comprises three different (yet undescribed) species. One species comprises isolates from grasslands of HAI and ALB (e.g., isolate LH08AG1034; **Fig. 7g**), identical to '*Marvania*' sp. WB67 (KF144207) from Westerhöfer creek within a highly supported clade. Second species from grasslands (HAI, SCH) (e.g., isolate LH10HG9020; **Fig. 7h**) is identical to *Nannochloris* sp. JL-4-6 (AY195983) from freshwater (USA). Third species from grassland (SCH) (e.g., LH08SG3093; LH08SG3078, **Fig. 7i**) is identical to *Nannochloris* sp. Ant-1 (EF440182) from permafrost (Antarctica). *Auxenochlorella*-like colorless isolates represent two different taxa on either species or even genus level. (I) *A. protothecoides* (e.g., isolate LH10HG6096; **Fig. 7k**) from grasslands (HAI) is identical to authentic strain *A. protothecoides* SAG 211-7a (X56101) isolated from tree sap (Germany). This species clusters together with *Chlorella* sp. CCAP 211/61 (AB206551) in a highly supported clade. (I) *Auxenochlorella* sp. SAG 2478 (= LH08AW4103; **Fig. 7l**; KP081390) from forest (ALB), is less closely (similarity = 97.46%) related to *A. protothecoides* SAG 211-7a, without any known closer relatives, probably representing either new species or genus. The whole clade including *Auxenochlorella* sp. and *A. protothecoides* is highly supported. Another highly supported clade, consisting of isolate LH10HG709K (**Fig. 7j**) from grassland soil (HAI) and clone HEW1B K3342 from forest soil (HAI), does not include other known relatives (not shown in phylogenetic trees in this Chapter, but in **Fig. 2 Chapter 3**). The species might represent a new genus with basal placement among the *Nannochloris*-like lineages within the Chlorellaceae.




Figure 5. 18S rDNA phylogeny of Green algae, part 2 including Trebouxiophyceae. New accessions are written in bold and grey underlined. Black dots mark references to microphotographs (Fig. 6, Fig. 7) sorted in top down order. Original drawings of representative morphologies are shown at each clade (scale bars = 5µm). Sequences of authentic strains are marked by '>'. Numbers next to branches indicate statistical support values (maximum-likelihood bootstraps/Bayesian posterior probabilities).




Figure 6. Microphotographs of monoclonal cultures of Green algae (Trebouxiophyceae). (a) *Stichococcus* sp.(I) LH08AW8025 (l=1.7-19.8 μm , w=1.4-3.6 μm); (b) *Stichococcus* sp.(II) LH08SG5057 (l=4.5-9.8 μm , w=1.9-3.3 μm); (c) *Stichococcus* sp.(VI) SAG 2482 (= LH08AW8023; l=3.8-7.2 μm , w=1.9-6.3 μm); (d) *Neocystis brevis* LH10HG9080 (l=4.8-8.7 μm , w=2.7-6.6 μm); (e) *Coccomyxa simplex* LH08SG9051 (l=4.4-7.8 μm , w=2.0-4.3 μm); (f) *Coccomyxa viridis* SAG 2483 (= LH08AW8039; l=4.7-8.4 μm , w=1.8-3.6 μm); (g) *Navicloris fusiformis* SAG 2477 (= LH08AW3007; l=8.3-15.3 μm , w=3.6-7.8 μm); (h) *Chloroidium saccharophilum* LH10HG7062 (l=5.6-9.1 μm , w=4.0-6.2 μm); (i) *Chloroidium* cf. *ellipsoideum* LH10HG9105 (l=4.8-7.8 μm , w=3.3-5.3 μm); (j) *Lobosphaera bisecta* LH10HG3P15 (\varnothing =6.2-9.6 μm); (k) *Xylochloris* sp. SAG 2382 (= LH08AG7024; l=4.7-12.9 μm , w=3.1-10.6 μm); (l) *Dictyochloropsis splendida* LH08AW3050 (\varnothing =7.7-39.4 μm). Scale bars = 10 μm .




Figure 7. Microphotographs of monoclonal cultures of Green algae (Trebouxiophyceae). (a) '*Chlorella*' cf. *mirabilis* LH08AG9040 ($\varnothing=3.0-6.4\ \mu\text{m}$); (c) *Chlorella* cf. *vulgaris* LH10HG2081 ($\varnothing=2.5-6.2\ \mu\text{m}$); (d) *Muriella terrestris* LH10HG7118; (e) *Nannochloris* sp. LH08SG8030 ($\varnothing=2.2-3.9\ \mu\text{m}$); (f) *Nannochloris* sp. LH10HG6095 ($\varnothing=2.1-3.9\ \mu\text{m}$); (g) Unidentified Chlorellaceae (I) LH08AG1034 ($\varnothing=2.4-4.5\ \mu\text{m}$); (h) Unidentified Chlorellaceae (II) LH10HG9020 ($\varnothing=2.4-4.8\ \mu\text{m}$); (i) Unidentified Chlorellaceae (III) LH08SG3078 ($\varnothing=2.1-3.7\ \mu\text{m}$); (j) Unidentified Chlorellaceae LH10HG709K ($l=3.1-5.1\ \mu\text{m}$, $w=2.0-3.7\ \mu\text{m}$); (k) *Auxenochlorella protothecoides* LH10HG6096 ($\varnothing=2.9-6.5\ \mu\text{m}$); (l) *Auxenochlorella* sp. SAG 2478 (= LH08AW4103; $\varnothing=2.2-5.2\ \mu\text{m}$). Scale bars = $10\ \mu\text{m}$.

Geographic dispersal and taxonomic 'novelty' of the isolated Green algae

In total, we identified 61 species of Green algae. Thirty-nine (53%) species were detected once and 34 (47%) species were detected more than once (e.g., *Chlorella vulgaris*; 19 times; **Table S4a**). Thirty (41%) species were retrieved from more than one plot and 18 (25%) species were retrieved from more than one exploratory (**Fig. 8**; **Table S4b**). There were only two species which we detected in all three exploratories, i.e., *Stichococcus* sp. (II) and *Neocystis brevis* (**Table S4b**). Almost all clades included both forest and grassland species, with the only exception of the Chlorellaceae with most species detected only in grasslands (**Fig. 8**). More than a half of all detected species (61%) exhibited > 99.90% sequence similarity to accessions known from previous studies. But only less than a half of these accessions belong to validly named (authentic) strains (**Fig. 9a**). Many next relatives to our species originate from Europe (54%), most of them from soil (30%) or various aquatic habitats (27%). Species less closely related to already published accessions (< 99.98% sequence similarity) mostly originate from grassland plots (62%; **Fig. 9a**). ITS2 sequence comparisons revealed occurrences of identical ribotypes separated by considerable distances, e.g., '*Chlorella*' *mirabilis*, *Chlorella vulgaris*, *Pseudostichococcus monallantoides* and *Klebsormidium dissectum* (**Fig. 9b**). Other notably high similarities were detected in *Chlamydomodium vacuolatum* (99%) and *Muriella terrestris* (98%) (**Fig. 9b**).




Figure 8. Diversity and distribution of all green microalgae isolated within this study. See further explanations on next page.

Figure 8. Diversity and distribution of all Green microalgae isolated within this study. The 18S phylogenetic tree gives an overview of all detected species (each species is represented by one sequence in the tree; due to a lack of space, names of reference sequences were removed from the figure; the comprehensive taxon samplings are shown in **Fig. 2**, **Fig. 5**). Numbers in brackets give percentage sequence similarity to the closest relatives. 'I' or 'II' indicate whether a species was found only once or multiple times within the Exploratories. Colored squares to the right of the tree summarize the distribution of each species across the three Exploratories (ALB, HAI, SCH) and across the two different habitats (grassland, forest). This evidence is based on sequenced isolates listed in **Table S4**. Further two squares indicate whether the closest GenBank-relative originate from terrestrial or aquatic habitats. Numbers in brackets behind isolate identifiers indicate (18S-based) sequence similarity to the closest cultured relatives. Algal clones, which are closely related to our isolates, are differentiated by colors due to their source (red: environmental clones obtained from the same sampling sites (Hallmann *et al.* in prep.-b); purple: GenBank accessions from culture-independent surveys; blue: GenBank accessions from culture-dependent studies).




Figure 9. Geographic distribution of Green algal species isolated from soils. (a) Pie charts derived from 18S rDNA sequence similarities illustrate (I) how many isolates could be identified at species level (and where their closest relatives originated from), (II) how many isolates represent undescribed species (and from which habitats these undescribed species originated from); (b) Map showing places where identical or highly similar ITS2 rDNA sequences were found. Evidence for 100%-identity: *Stichococcus* LH08SW8044 (DE) and SAG 379-4 (CA), *Chlorella* LH08SG3006 (DE) and FN298918 (US-MA), *Klebsormidium* LH08AG1113 (DE) and KS164CL6L (EC), *'Chlorella' mirabilis* LH08AG9040 (DE) and KFFB12-1 (EC); Evidence for high similarity [98-99%]: *Chlamydomodium* LH10HG1013 (DE) and FR865591 (US-AK), *Muriella* LH08SG3009 (DE) and JN653521 (AQ). CA=Canada, DE=Germany, US-AK=Alaska, US-MA= Massachusetts, EC=Ecuador. AQ=Antarctica.

Stramenopiles: Xanthophyceae

Phylogenetic analysis of the Stramenopiles (**Fig. 10a**), assigned isolates to the classes Xanthophyceae (six clades) and Eustigmatophyceae (one clade) (**Fig. 10b**).

Tribonematalean lineage (Maistro *et al.* 2009). Tribonematalean isolates with unbranched filaments could be assigned to *Xanthonema* (three species) and *Heterothrix* (two species). (I) *Xanthonema* sp. (e.g., isolates LH10HG7029, LH10HG9058; **Fig. 11a**), detected in grassland (HAI) and forest (HAI), is closely related to *X. exile* PAB 395 (AM491615). (I) *Xanthonema* cf. *exile* (isolate LH10HG7078; **Fig. 11b**) from grassland (HAI) is almost identical to *X. exile* PAB 395 (AM491615). (III) *Xanthonema* cf. *bristolianum* (isolates LH10HG6059, LH08HW9018;

Fig. 11c) from grassland (HAI) and forest (HAI), is identical to terrestrial *X. bristolianum* CCALA 516 (AM490819) isolated from snow (Slovakia). The *Xanthonema* clade is well supported. (IV) *Heterothrix* sp. (isolates LH10HG7061, **Fig. 11d**; LH08SG5052, **Fig. 11e**) from grasslands (HAI, SCH) is related to *Heterothrix* sp. ACOLA1 (AM491612) and might represent a new or undescribed species. (V) *H. sessile* (e.g., isolate LH10HG5079) from grassland (HAI) is identical to *H. sessile* IBSG-V28 (AM490818). The *Heterothrix* clade is highly supported.

Chlorellidialean lineage (Maistro *et al.* 2009). Chlorellidialean heterotrichal isolates clustered in three different species of *Heterococcus*. (I) *Heterococcus* sp. (isolates LH10HG2140, LH10HG9085; **Fig. 11f**) from grasslands (HAI) is related to freshwater Xanthophyceae sp. IX3 (FJ946906) from Antarctica and might represent a new species. (II) *Heterococcus* cf. *chodatii* (e.g., isolate LH10HG9111; **Fig. 11g, 11h**) from grasslands (HAI) is identical to authentic strain *H. chodatii* SAG 835-3 (AM490822) from a subaerial habitat (Switzerland). The clade including *H. chodatii* (and *H. pleurococcoides* PAB 380; AJ579335) is well supported. (III) *Heterococcus* cf. *caespitosus* (isolate LH08AG2020) from grassland (ALB) is almost identical to authentic strain *H. caespitosus* SAG 835-2a (AM490820) from soil (Germany) and at the same to *H. protonematoideus* SAG 835-9 (AJ579334) from soil (Switzerland). The clade including both species is highly supported, as well as the whole *Heterococcus* clade.

Botrydiopsalean lineage (Maistro *et al.* 2009). Botrydiopsalean isolates (two species) exhibited either cell packages or solitary spherical cells with numerous discoidal chloroplasts. (I) Botrydiopsalean sp. (e.g., isolate LH08AW1076; **Fig. 11i**) from grassland (SCH) and forest (ALB) is related to *Chlorellidium pyrenoidosum* PAB 785 (AJ579338) from soil (Antarctica). The clade including both species is not supported and the isolate LH08AW1076 might represent a new species. (II) *Botrydiopsis* sp. (isolate LH08AW4043; **Fig. 11j**) from forest (ALB) is related to *B. callosa* SAG 30.83 (AJ579340) from soil (Italy). The clade including both species is well supported and the isolate LH08AW4043 might represent a new species.

The thallosous genus *Asterosiphon* Dangeard is represented by *Asterosiphon* sp. (isolate LH10HG3064; **Fig. 11k**) from grassland (HAI), which is less closely related (similarity = 98.54%) to *A. dichotomus* UTEX LB 2066 (AM490829). *Asterosiphon* sp. LH10HG3064 exhibited filamentous and coccal stages in the culture. The *Asterosiphon* clade is well supported and the isolate LH10HG3064 probably represents a new species.

Stramenopiles: Eustigmatophyceae

The genus *Eustigmatos* Hibberd is represented by *Eustigmatos* sp. (isolates LH10HG5036, LH10HG9133; **Fig. 11l**) from grasslands (HAI) is closely related to *E. magna* CCMP 387

(U41051) from soil (New Zealand). Both isolates exhibit spherical cells with characteristic angular shaped 'false pyrenoid'. The clade including both isolates and *E. magna* CCMP 387 is supported.




Figure 10. Phylogeny of some Stramenopiles, including Xanthophyceae and Eustigmatophyceae. (a) Schematic multi-gene phylogeny (modified after Maistro *et al.* (2009)) of the same Stramenopile groups which were used for the 18S phylogeny. (b) 18S rDNA phylogenetic tree showing placement of the new isolates (grey underlined; black dots mark isolates which are shown on microphotographs (Fig. 11). References to the microphotographs (Fig. 11, listed below clade names) correspond to black dots in top down order. Sequenced authentic strains are marked by a '<' sign. Numbers next to branches indicate statistical support values (maximum-likelihood bootstraps/Bayesian posterior probabilities).




Figure 11. Microphotographs of monoclonal cultures (Trebouxiophyceae, Xanthophyceae, Eustigmatophyceae). (a) *Xanthonema* sp. LH10HG9058; (b) *Xanthonema exile* LH10HG7078; (c) *Xanthonema* cf. *bristolianum* LH08HW9018; (d) *Heterothrix* sp. LH10HG7061; (e) *Heterothrix* sp. LH08SG5052; (f) *Heterococcus* sp. LH10HG9085; (g) *Heterococcus* cf. *chodatii* LH10HG9111; (h) *Heterococcus* cf. *chodatii* LH10HG9111; (i) Botrydiopsalean sp. LH08AW1076; (j) *Botrydiopsis* cf. *callosa* LH08AW4043; (k) *Asterosiphon* sp. LH10HG3064; (l) *Eustigmatos* sp. LH10HG9133. Scale bars = 10 μ m.

Morphological diversity observed in cultured soil samples

Microscopic investigation of the cultured soil samples (liquid cultures, agar plates, cover slips; **Fig. 1g-i**) revealed 45 distinct morphotypes (**Figs. S1, S2, S3, S4**). We compared the diversities among the Exploratories and between the two soil horizons (**Fig. S5a-b**). The A-horizon counts lower morphotype diversity (22 morphotypes; 57 plots/57 samples) than the O-horizon (46 morphotypes; 5 plots/20 samples; **Fig. S5b**). The highest diversity exhibited Chlorophyta/Streptophyta (24), followed by Bacillariophyceae (8), Cyanobacteria (7), Xanthophyceae/Eustigmatophyceae (5), Rhodophyta (1) and Cryptophyta (1) (**Table S6, S7**). In general, the number of morphotypes is nearly the same in grasslands and forests (**Fig. S5b-c**), irrespective of the sampling technique (drill cores, soil surface samples). The A-horizon sampling revealed negligible differences in alpha-diversity between the three regions. In average, intermediately managed plots were slightly more diverse than extensively and intensively managed plots (**Fig. S5f**). The same applies for forests and grasslands, yet without statistical support. However, the morphotype composition differed between forest and grassland plots, explained through differences in physico-chemical parameters (**Fig. S5d**). The statistically significant ordination model ($P < 0.05$) based on Redundancy Analysis (RDA) revealed separation of forest and grassland replicate samples along the first ordination axis which is correlated with pH values ($R = 0.74$; $P < 0.05$). The contribution of total nitrogen ($R = 0.26$; $P < 0.05$) and organic carbon ($R = 0.11$; $P < 0.05$) is lower. The difference between soil communities in forests and grasslands can be explained by higher diversity of Cyanobacteria, Diatoms and Xanthophytes in grasslands. Considering the O-horizon (soil surface samples from HAI), multiple samples within a single plot exhibited different morphotype compositions (**Fig. S5e**). The ordination based on NMDS analysis (Shepard stress = 0.11) separates forest and grassland samples (axis 1) and points out higher heterogeneity of the forest plots (axis 2).

Discussion

Culturing of soil samples from 30 forest and 27 grassland sites retrieved 188 monoclonal cultures of Green microalgae. We inferred conspecificity of the sequenced isolates based on ≤ 0.001 SSU dissimilarity threshold. In multiple cases, our isolates were divergent from already sequenced authentic strains available from GenBank. Therefore, to denominate detected species, we relied on the phylogenetic species concept (Leliaert *et al.* 2009; Leliaert *et al.* 2014), not strictly following the taxonomical nomenclature (Cantino and de Queiroz 2007; McNeill *et al.* 2012). This approach distinguished 61 species covering five classes of Green algae: Chloro-,

Klebsormidio-, Pedino-, Trebouxio- and Ulvophyceae. In the following we focus mainly on two classes accounting for the most detected species - Trebouxioophyceae (36 spp.; 8 lineages) and Chlorophyceae (21 spp.; 10 lineages). We discuss whether our soil isolates can be recognized in already denominated algal strains deposited in public culture collections (e.g., SAG, UTEX, CCAP). Then we trace the distribution of our species by analysing records in GenBank and discuss their putative detectability in different environments by culture-independent sequencing.

New soil isolates and their cultured relatives

More than 50% of Green microalgae we isolated from soils are putatively conspecific with isolates from previous studies (**Fig. 8; Fig. 9a**). The most of these conspecific cultures originate from Central and Western Europe (54%) and from the USA (14%), corresponding to strains deposited in the Culture Collection of Algae in Göttingen (SAG) incl. their derivatives in UTEX or CCAP. Exotic counterpart cultures were few, e.g., *Coccomyxa* sp. (HE586515) from Indonesia or *Nannochloris* sp. from Antarctica (EF440182; Gilichinsky *et al.* 2007). Sequenced strains—conspecific with our isolates—originate from terrestrial (edaphic, epilithic, epixylic/lichenized) and aquatic environments. So far, there are no estimates of total diversity of microscopic Green algae in soils. However, the number of cultured species which match environmental clones, suggests that the uncultured fraction is not as extensive as in bacteria (99%; Hirsch *et al.* 2010). All 61 isolated species grow easily on agarized (solid) media, which are selective and might favor so called 'weedy' or generalists species (Škaloud *et al.* 2014a). Additional enrichment techniques (growth in liquid media or on coverslips), indeed, retrieved additional species, which we did not retrieved as monoclonal cultures for sequencing (e.g., *Macrochloris*, *Spongiochloris*, *Keratococcus*, *Podohedra*, *Scotinosphaera*, *Pseudendoclonium*, *Interfilum* or *Cosmarium*; **Fig. S1, S2**). The identification literature by Ettl and Gärtner (1995) covers about 500 morphospecies of terrestrial Green microalgae—partially documented as sequenced strains deposited in culture collections. We re-detect some of them in the present study (via SSU-comparisons) and could either support or reject their ecological preference (so far inferred via morphospecies-based approaches). Among such sequenced Green microalgae, which were documented by particular algal strains in (Ettl and Gärtner 1995), we confirmed characteristic grassland species: *Bracteacoccus cohaerens* UTEX 1272 (Bischoff and Bold 1963), '*Chlorococcum*' *gelatinosum* SAG 64.80 (Archibald and Bold 1970), *Chlamydomonas typica* NIES-2246 (Deason and Bold 1960), '*Chlorella*' *mirabilis* Andreyeva 748-I (Andreyeva 1973), *Desmotetra stigmatica* UTEX B 962 (Deason and Floyd 1987), *Muriella terrestris* ASIB V38 (Ettl and Gärtner 1995) and *Pseudomuriella aurantiaca* SAG 249-1 (Hanagata 1998). A few species were repeatedly verified

in forests, e.g., *Chloromonas gerloffii* and *C. rosae* (Ettl 1963) or in soils of both forests and grasslands, e.g., *Lobosphaera bisecta* SAG 2043 (Trenkwalder 1975), *Chlorococcum minutum* ASIB T50 (Trenkwalder 1975) and *Heterochlamydomonas inaequalis* UTEX 1705 (Cox and Deason 1969). Finally, in our soil samples we recognized well documented terrestrial generalists such as: *Chloroidium saccharophilum* SAG 211-9a (Darienkov *et al.* 2010), *Dictyochloropsis splendida* CAUP H8601 (Škaloud *et al.* 2005) and '*Chlorella*' *sphaerica* SAG 11.88 (Tschermaek-Woess 1988). Despite the sampling efforts in forests, we did not retrieve species characteristic for temperate tree-bark biofilms or lichens, e.g., *Apatococcus* (Hallmann *et al.* 2009; Hallmann *et al.* in prep.-a), *Elliptochloris* (Eliáš *et al.* 2008), *Kalinella* (Neustupa *et al.* 2009), *Leptochlorella* (Neustupa *et al.* 2013a), *Parachloroidium* (Neustupa *et al.* 2013b), *Trebouxia* (Friedl and Rokitta 1997; Kroken and Taylor 2000) or *Asterochloris* (Škaloud *et al.* 2015). This might be due to selectivity of the used culturing medium, reflecting nutrient demands of soil but not corticolous species.

New taxa detected in soils

Seven isolates are considerably divergent from all GenBank clones or cultures, presumably representing novel taxa. The novel Chlorophyceae are: *Jenufa* sp. SAG 2383, *Oogamochlamys* sp.(I) LH08SG8047, *Oogamochlamys* sp.(II) SAG 2476. The novel Trebouxiophyceae are: *Auxenochlorella* sp. SAG 2478, *Stichococcus* sp.(V) LH08SW1099, '*Navichloris fusiformis*' SAG 2477, *Xylochloris* sp. SAG 2382. *Jenufa* Němcová, Eliáš, Škaloud & Neustupa represents a coccoid genus and at the same time a lineage incertae sedis within the class Chlorophyceae. Two terrestrial species were described from Southeast Asia (Němcová *et al.* 2011) and multiple congeneric accessions were acquired also from Neotropics (Hodač *et al.* 2012), the Alps (Horath and Bachofen 2009) and finally in the frame of this study (*Jenufa* sp. SAG 2383). The only European isolate *Jenufa* sp. SAG 2383 (**Fig. 4I**) might represent commonly occurring terrestrial species detected in forest soils and epilithic biofilms (Hodač *et al.* 2012).

Oogamochlamys Pröschold, Marin, Schlösser & Melkonian is a monadoid genus within the *Oogamochlamydia* clade (Chlamydomonadales). The three species described by Pröschold *et al.* (2001) genetically differ from our two congeneric isolates provisionally denominated *Oogamochlamys* sp.(I) LH08SG8047 (**Fig. 4a**) and *O.* sp.(II) SAG 2476 (**Fig. 4b**). Distribution and ecology of both species remains unclear, as they were detected only once, *Oogamochlamys* sp.(I) in one forest site and *O.* sp.(II) in one grassland site. Remarkably, all other described *Oogamochlamys* species were isolated from soils in Africa and North America (Pröschold *et al.* 2001). And at least one another verifiably congeneric isolate—*Chlamydomonas* sp. CCAP

11/159—was detected in a lake in North America (GenBank acc. no. FR865591).

Auxenochlorella (Shihira & Krauss) Kalina & Punčochářová is a genus within the Chlorellaceae comprising three facultatively heterotrophic species (Darienkov and Pröschold 2015; Rodó and Molinari-Novoa 2015). The new facultatively heterotrophic isolate *Auxenochlorella* sp. SAG 2478 (**Fig. 7l**) is verifiably congeneric with all described *Auxenochlorella* species but is still considerably divergent from all accessions available in GenBank. *Auxenochlorella* sp. SAG 2478 is the first species of this genus which was isolated from soil. Other *Auxenochlorella* species were isolated from different terrestrial and aquatic habitats, e.g., tree sap and *Hydra viridis* endosymbiont (Darienkov and Pröschold 2015).

Stichococcus Nägeli is a common denomination of rod-shaped Green algae from the *Prasiola* clade, counting 50 described species (Karbovska and Kostikov 2012a). Only two *Stichococcus* species were verified by genetic data and the polyphyletic taxon awaits taxonomical revision. All seven *Stichococcus*-like species detected in the frame of this study represent new taxa, however, six of them are closely related with already existing cultures or clones. Except for the isolate *Stichococcus* sp. LH08SW1099 (**Fig. 8; Fig. 5i Chapter 3**).

Xylochloris Neustupa, Eliáš & Škaloud is a genus incertae sedis within the Trebouxiophyceae with one described terrestrial species *X. irregularis* isolated in Southeast Asia (Neustupa *et al.* 2011). Our soil isolate *Xylochloris* sp. SAG 2382 (**Fig. 6k**) is putatively congeneric with *X. irregularis*, together with clones from European soils and tree-barks in South America (Hodač *et al.* 2012; Spitzer *et al.* 2014). The genus is probably widely distributed in Europe, as recently reported from tree-bark in the Southern Europe (Kulichová *et al.* 2014).

Navichloris fusiformis L.Hodač *et al.*, ad interim (**Fig. 6g**) is a new species and genus of an undescribed lineage within the class Trebouxiophyceae. The existence of this novel lineage was already proposed by Lewis and Lewis (2005), who phylogenetically analyzed the strain *Chlorella* sp. BC4VF9 from Baja California, Mexico, without affiliation to any known trebouxiophycean clade. Costello *et al.* (2009) pointed out close relationship of *Chlorella* sp. BC4VF9 and two clones sampled from cold-fumarole soil on the Socompa volcano, Atacama, Chile. However, the obviously novel trebouxiophycean clade remained undescribed, since the only existing culture *Chlorella* sp. BC4VF9 died (Fučíková *et al.* 2014). Phylogenetic analysis (**Fig. 12**) show that the three desert accessions cluster with the strain SAG 2477 isolated in the frame of this study from a forest soil in Schwäbische Alb, Germany. The clade is statistically supported and represents a phylogenetically unique lineage of terrestrial trebouxiophycean microalgae. The strain SAG 2477 as well exhibits a unique fusiform/elliptic morphology, generally resembling *Coccomyxa* (Schmidle 1901). However, phylogenetically it is not related to any member of the recently

revisited genus *Coccomyxa* (Darienkov *et al.* 2015). The characteristic feature of the strain SAG 2477 are lens-like light zones in the chloroplast lobes (**Fig. 6g**), similar to *Elliptochloris incisiformis* Hoffmann & Kostikov described from forest soil in Belgium (comp. Fig. 27, Hoffmann *et al.* 2007), but molecular marker are not available. Notably, the fusiform morphology shared by SAG 2477 and *Chlorella* sp. BC4VF9 (<http://pediastrum.eeb.uconn.edu/isolates/detail/411>) further supports the relatedness of both species.




Figure 12. Schematic 18S rDNA phylogeny of the class Trebouxiophyceae. The newly proposed *Navichloris* lineage is highlighted in a violet box. The full 18S rDNA dataset used for the tree computation is identical as in Fučíková *et al.* (2014), Fig. 3. We complemented the dataset by adding three sequences, i.e., KP081399 (SAG 2477; retrieved in this study), FJ592371, FJ592503 (Costello *et al.* 2009); the resulting 18S rDNA dataset comprised 135 sequences. The phylogeny was inferred using RAxML (Stamatakis *et al.* 2008) and statistical support values for the most branches (maximum likelihood/posterior probabilities) were computed in the same program and in MrBayes (Ronquist *et al.* 2012), respectively). Additional statistical support values (maximum parsimony and bio-neighbor-joining) for the *Navichloris* lineage were computed in the program SeaView ver. 4 (Gouy *et al.* 2010).

Cultured Green microalgae re-detected in environmental clones

Almost 90% of our cultured species match clones amplified from environmental samples, when considering a ≤ 0.010 SSU-similarity threshold. Even though this similarity level might lack accuracy for species discrimination, it enables to detect species which are closely related; in comparison, Achtman and Wagner (2008) indicates a ≤ 0.013 SSU-threshold for conspecificity in eubacteria. Culture-independent cloning of microalgae from identical sampling sites, conducted by Hallmann *et al.* (in prep.-b), allows to recognize those species, which are re-detectable by both approaches. Due to their frequent occurrence in soils, they might reach high population densities, being key players of soil algal communities (Büdel *et al.* 2009). This applies for several trebouxiphytes (e.g., *Stichococcus* spp., *Coccomyxa* spp., *Chlorella vulgaris*, *Muriella*-like species) and chlorophytes (*Oogamochlamydia*, *Tatrensinia*, *Jenufa*, *Scenedesmaceae*). Even though comparable cloning studies on soil microalgae are rare, our isolates additionally matched clones from North American soils (*Neocystis*, *Chloroidium*, *Reinhardtina*, *Desmotetra*; Lesaulnier *et al.* 2008) and from Dutch historical soil samples (*Muriella*; Moon-van der Staay *et al.* 2006). In multiple cases, clones closely related to our isolates, were amplified from either much dryer or even aquatic habitats, such as *Stichococcus* on marble monuments (Hallmann *et al.* 2013b) and on rocks in the Alps (Horath and Bachofen 2009), *Diplosphaera* in Rio Tinto river (Aguilera *et al.* 2007); *Coccomyxa* (HE617184) in an acidic lake and purification plant in Japan (GenBank acc. no. AB721066), *Lobosphaera* in sulfidic water (Behnke *et al.* 2006) or *Stephanosphaerina* and *Pseudendocloniopsis* in glacier debris of Alaska (Schmidt and Darcy 2014). Other isolates matched clones amplified from cultured environmental material, such as building stone (Hallmann *et al.* 2013a) and even from aeroplankton in Greece (Genitsaris *et al.* 2011). Remaining cultures were more divergent from all available environmental clones; hence we queried them against selected soil metagenomes deposited in Sequence Read Archive (Wheeler *et al.* 2007). Here we achieved only a ≤ 0.03 match of *Chlorococcum* cf. *minutum* SAG 2479 and *Chlamydomonas* cf. *gerloffii* LH08SW5031 with SSU-V4 pyrotag reads from permafrost metagenome in Sweden (Mondav *et al.* 2014).

Conclusive remarks

Our results suggest that Green microalgae in soils of Central Europe are far from known, being less intensively studied than e.g. desert soil crusts (Lewis and Lewis 2005; Fučíková *et al.* 2014). Even the choice of standard culturing media—supposedly not favoring demanding species (Lukešová 2001; Hoffmann *et al.* 2007)—retrieved a plethora of (phylogenetic) species. The most of them are re-detectable in environmental clone libraries, but some would remain

undiscovered without culturing approach. Particularly high phylogenetic diversity hide in morphospecies known as *Stichococcus* (Nägeli 1849) or *Nannochloris*-like (Henley *et al.* 2004)—consisting of unrelated species which exhibit morphological convergence characteristic for terrestrial Green microalgae (Sharma and Rai 2010). Simplistic morphology of microscopic algae favor their fast dispersal (Sharma *et al.* 2007), however, cosmopolitanism of Green microalgae remains questionable (Lawley *et al.* 2004; De Wever *et al.* 2009). Traditional morphology-based surveys generally proclaim ubiquity of soil Green microalgae (Feher 1948; Starks *et al.* 1981; Ettl and Gärtner 1995). A summary of morphospecies inventory lists collected worldwide (**Fig. S6**) suggests that Green algal communities might share a general morphological composition (i.e., consist of *Stichococcus*-, *Chlorella*-, *Chlamydomonas*-, *Chlorococcum*- and *Klebsormidium*-like morphospecies). Our cultures contribute DNA-based hints indirectly supporting the above mentioned observations—we detected identical ITS2 signatures of *Stichococcus*, *Chlorella* and *Klebsormidium* originating from two different continents (**Fig. 9b**). The ITS2 data further supported the wide distribution of *Muriella* (Kochkina *et al.* 2014) and *Chlorococcum* (GenBank acc. no. FR865591). It remains a major challenge of metagenomics to explore phylogenetic structure and geographic distribution of microscopic algae (Marande *et al.* 2009; Raven 2012; Lie *et al.* 2014). However, not less important is to link OTUs from environmental metagenomes with physical cultures of well characterized species (Richards and Bass 2005; Richards *et al.* 2005; Dickie 2010).

Acknowledgements

We thank the managers of the three Exploratories, Kirsten Reichel-Jung, Swen Renner, Katrin Hartwich, Sonja Gockel, Kerstin Wiesner, and Martin Gorke for their work in maintaining the plot and project infrastructure; Christiane Fischer and Simone Pfeiffer for giving support through the central office, Michael Owonibi for managing the central data base, and Markus Fischer, Eduard Linsenmair, Dominik Hessenmöller, Jens Nieschulze, Daniel Prati, Ingo Schöning, François Buscot, Ernst-Detlef Schulze, Wolfgang W. Weisser and the late Elisabeth Kalko for their role in setting up the Biodiversity Exploratories project. The work has been (partly) funded by the DFG Priority Program 1374 "Infrastructure-Biodiversity-Exploratories" (DFG-FR 905/16-1 and FR 905/17-1). Field work permits were issued by the responsible state environmental offices of Baden-Württemberg, Thüringen, and Brandenburg (according to § 72 BbgNatSchG).

Chapter 2

Diversity of microscopic green algae (Chlorophyta) in calcifying biofilms of two karstic streams in Germany

Ladislav Hodač¹, Nicole Brinkmann², Kathrin I. Mohr³, Gernot Arp², Christine Hallmann¹, Jessica Ramm⁴, Karolin Spitzer¹, Thomas Friedl¹

¹University of Göttingen, Experimental Phycology and Culture Collection of Algae (SAG), Göttingen, Germany

²University of Göttingen, Geoscience Centre, Göttingen, Germany

³Helmholtz Centre for Infection Research, Braunschweig, Germany

⁴Brandenburg University of Technology, Department of Freshwater Conservation, Bad Saarow, Germany

(Published in Geomicrobiology Journal; doi:10.1080/01490451.2013.878418)

Abstract

For the first time the diversity of microscopic green algae (Chlorophyta) from calcified biofilms of karstic streams was analyzed using a combined approach based on pure cultures, i.e. 18S rRNA gene sequencing and microscopic analyses. Our study focused on two creeks in Germany. A considerable diversity of 34 species of green microalgae comprising three classes, the Trebouxiophyceae, Chlorophyceae and Ulvophyceae, was recovered. The biofilms of both streams were rather different in their species compositions which may reflect that they are exposed to differed hydrochemical conditions. The shallow Westerhöfer creek harbored predominantly Trebouxiophyceae and exhibited higher Mg^{2+} and SO_4^{2-} concentrations. In contrast, the deeper, longer and spatially more heterogeneous Deinschwanger creek harbored numerous species of Chlorophyceae. A lower number of species from the Ulvophyceae were spread on both studied streams. The closest relatives of the identified species were from other freshwater habitats, but mostly from phytoplankton. However, also several species we recovered from freshwater for the first time; so far they have been known from terrestrial habitats only. Less than half of the recovered species could be identified with names at the species level based on high sequence identities with available sequences from named reference strains. Most recovered species could not be identified with names to species level, i.e. they may represent species for which no 18S rRNA gene sequence have become available yet or which have been retrieved for

the first time. The genus *Gongrosira* Kützing, often reported from freshwater tufa-stromatolites, was found to represent most likely a collective morphotype formed by several genera nested within the Ulvophyceae.

Keywords: Chlorophyta; Trebouxiophyceae; Chlorophyceae; Ulvophyceae; 18S rRNA gene phylogeny; karst-water creeks; cultures

Introduction

Many karstic streams in Europe and elsewhere are characterized by calcium carbonate deposits which veneer macrophytes as well as biofilm-covered rock surfaces at the stream bed. These deposits are termed tufa stromatolites, defined as macroscopically laminated benthic microbial deposits produced by precipitation of minerals on organic tissue (Riding 1991). Biomineralization (biological processes) and inorganic precipitation may act together (Ford and Pedley 1996) or photosynthetic CO₂ assimilation by cyanobacteria, eukaryotic algae, and plants may be the primary cause for the carbonate precipitation (Pia 1926; Wallner 1934; Hepperle and Krienitz 1996). Indeed, microsensor studies have demonstrated a photosynthetic control of CaCO₃ precipitation for biofilm-covered surfaces, while inorganically driven precipitation prevails e.g. at moss surfaces (Shiraishi *et al.* 2008a; Shiraishi *et al.* 2008b). Microscopic studies revealed the dominance of filamentous cyanobacteria in the calcified biofilms of freshwater karst creeks (Freytet and Plet 1996; Garcia-Pichel *et al.* 2004; Brinkmann *et al.* 2015). But also diatoms, xanthophytes, red algae and other microscopic algae as well as bryophytes and microscopic fungi occur being associated with freshwater stromatolites (Winsborough and Golubić 1987; Heath *et al.* 1995; Freytet and Verrecchia 1998; Bilan and Usov 2001; Brinkmann *et al.* 2015).

In an ongoing larger study the possible roles of photosynthesis and respiration in calcification processes are being studied in detail at two exemplar karstic streams with prominent CO₂-degassing along their course, the Westerhöfer creek and the Deinschwanger creek in Germany. Both tufa-forming streams attain high calcite supersaturation during their course downstream. The Westerhöfer creek, located in Middle Germany in the westerly Harz-foreland (51°45'N, 10°5'E), is 325 m long and less than 2 m wide and has its source in limestones and evaporites of the Middle Triassic Muschelkalk Group. The Deinschwanger creek is located in southern Germany at the western rim of the Franconian Alb (49°23'N, 11°28'E). It is fed by three main springs and a number of side springs, most of them discharging from the Upper Jurassic

Weißjura-Group aquifer. Compared to the Deinschwanger creek the Westerhöfer creek is rich in Mg^{2+} and SO_4^{2-} . Microscopy of biofilm samples from both creeks revealed cyanobacteria and diatoms as the dominant algae (Brinkmann *et al.* 2015), but other micro-algae were found only rarely or not at all. Concurrently with a study on the biodiversity of the cyanobacteria and diatoms from both creeks (Brinkmann *et al.* 2015), cultures of green algae were developed. Interestingly, in the enrichment cultures an unexpected variety of green micro-algae appeared besides numerous cyanobacteria and diatoms. It is not known yet whether there are green algal taxa with strict or even any preference for calcifying biofilms. Their presence in calcifying biofilms may even be entirely given by accident. There is an expectation that the most green algae in calcifying biofilms could originate from soils and other aerial and subaerial habitats. Here we report about the phylogenetic and morphological diversity of these green microalgae.

Materials and Methods

Sampling, culturing and microscopy

Biofilms from both, the Westerhöfer and Deinschwanger creeks, were collected in the spring or early summer (May/June) in 2005–2007. Five sites of the Westerhöfer creek (abbreviated as WB; **Figure A**) and eight sites from Deinschwanger creek (abbreviated as DB) were selected for sampling of apparently algae-dominated biofilms (**Table 1**). For starting enrichment cultures all




Figure A. Freshwater stromatolites (Westerhöfer Bach)

samples from the WB were pooled together because it is rather short and only a short segment (about 350 m) of the creek was investigated, whereas for the DB the samples from different locations were analysed separately to better reflect its higher habitat heterogeneity. The biofilms were scratched off from stone surfaces using an ethanol-sterilized knife or spatula and transferred to 1.5 ml reaction tubes which were cooled until further processing in the laboratory the following day. A spatula-full of biofilm material was then transferred into different standard liquid growth media. Because our initial main focus was to establish cultures of cyanobacteria and diatoms, growth media

provoking the development of these algae were used, i.e. BG11, BG11 without citrate, Z, Z 45/4, and ES (<http://www.uni-goettingen.de/de/186449.html>). Apart from the expected growth of cyanobacteria and diatoms, also intensive green algal growth was observed after about 5–10 days of cultivation. Putative green algal colonies were transferred on 1.5% agar plates with 3N BBM+V medium (<http://www.uni-goettingen.de/de/186449.html>). After incubation for another 4–8 weeks, all green algal morphotypes were isolated into unialgal cultures by several rounds of streaking on fresh agar plates of 3N BBM+V medium using sterile platinum needles. Finally, the unialgal isolates were microscopically checked for purity and further maintained on agar slants at 18°C under a light/dark regime of 14-h:10-h and a photon flow rate of about 25 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ from white fluorescent bulbs. A total of 77 pure cultures was established, out of which 45 were studied in more detail (**Table S1**). Nineteen of them were made publicly available, i.e. accessioned by the SAG culture collection (Göttingen University, Germany; www.epsag.uni-goettingen.de). For microscopy an Olympus BX60 microscope (Tokyo, Japan) with Nomarski DIC optics and an attached ColorView III camera (Soft Imaging System, Münster, Germany) was used. Micrographs were processed using Cell[^]D image software (Soft Imaging System, Münster, Germany).

DNA extraction, PCR and sequencing

DNA was extracted from all unialgal isolates (**Table S1**) using the Invisorb Spin Plant Kit (Stratec, Berlin, Germany) as recommended by the manufacturer. Nuclear-encoded 18S rRNA genes were amplified using primers NS1 and 18L (Hamby *et al.* 1988). For some strains the 18S and ITS1-5.8S-ITS2 rRNA gene region was amplified using primers NS1 and LR1850 (Friedl 1996). If a culture was suspected to be contaminated by fungi or bacteria, PCR primer 1650R (3'-TCACCAGCACAYYCAAT-5'; pos. 1652-1636 of the 18S rRNA gene sequence of *Chlorella vulgaris* SAG 211-11b, FM205832) which preferentially binds to members of Chlorophyta, was used as the reverse PCR primer. Conditions for PCR and cycle sequencing reactions and the standard set of sequencing primers were as described earlier (Mikhailyuk *et al.* 2008). The newly determined sequences were deposited in GenBank under the accession numbers KF144164-KF144240 (**Table S1**). In addition also the 18S rRNA gene sequences for the following strains were determined as references: *Chlorococcum vacuolatum* Starr SAG 213-8 (KF144189), *Dilabifilum printzii* (Vischer) Bourelly SAG 467-1 (KF144198), *Scotinosphaera gibberosa* (Vodenicarov & Benderliev) Wujek & R.H.Thompson SAG 75.80 (KF144229) and *S. lemnae* (Punčochářová) Wujek & R.H.Thompson SAG 240-1 (KF144230).

Table 1. Distribution of the recovered green algal species at the sampling sites.

| Class | Species | DB1 | DB2 | DB3 | DB4 | DB5 | DB6 | DB9 | DBS | WB |
|------------------|---|------|------|------|------|------|------|------|------|-----------|
| | | 1.19 | 1.15 | 1.11 | 1.09 | 1.09 | 1.11 | 0.45 | 0.01 | 0.28-0.99 |
| | | 3 | 3 | 2 | 1 | 1 | 11 | 2 | 4 | 19 |
| Trebouxiophyceae | <i>Chlorella</i> sp.* | + | + | + | - | - | + | - | + | - |
| | <i>Coccomyxa</i> cf. <i>pringsheimii</i> | - | - | - | - | - | - | - | + | + |
| | <i>Coccomyxa</i> cf. <i>simplex</i> | - | - | - | - | - | - | - | - | + |
| | <i>Elliptochloris subsphaerica</i> * | - | - | - | - | - | - | - | - | + |
| | <i>Marvania</i> sp. | - | - | - | - | - | - | - | - | + |
| | <i>Muriella terrestris</i> ** | - | + | - | - | - | - | - | - | + |
| | <i>Neocystis</i> cf. <i>mucosa</i> | - | - | - | - | - | - | - | - | + |
| | <i>Stichococcus bacillaris</i> ** | - | - | - | - | - | - | - | - | + |
| | <i>Stichococcus</i> cf. <i>deasonii</i> | - | - | - | - | - | - | - | - | + |
| | <i>Stichococcus mirabilis</i> ** | - | - | - | - | - | - | - | - | + |
| | <i>Stichococcus</i> sp.1** | - | - | - | - | - | + | - | - | + |
| | <i>Stichococcus</i> sp.2 | - | - | - | + | - | - | - | - | - |
| | <i>Stichococcus</i> sp.3 | - | - | - | - | - | - | - | - | + |
| | <i>Stichococcus</i> sp.4 | - | - | - | - | - | - | - | - | + |
| Chlorophyceae | <i>Acutodesmus obliquus</i> ** | - | - | - | - | - | + | - | - | - |
| | <i>Bracteacoccus aerius</i> -relative | - | - | - | - | - | - | - | - | + |
| | <i>Bracteacoccus</i> sp. | - | - | - | - | - | - | + | - | - |
| | <i>Chlamydomonas</i> sp. | - | - | - | - | - | - | - | - | + |
| | <i>Chlamydomonium</i> sp. | - | - | - | - | - | + | - | - | - |
| | <i>Chlorococcum minutum</i> -relative | - | - | + | - | - | - | - | - | - |
| | <i>Chlorococcum sphacosum</i> * | + | - | - | - | - | - | - | - | - |
| | <i>Chlorococcum ellipsoideum</i> -relative1 | - | - | - | - | + | - | - | - | - |
| | <i>Chlorococcum ellipsoideum</i> -relative2 | - | - | - | - | - | + | - | - | + |
| | <i>Desmodesmus</i> cf. <i>armatus</i> | - | - | - | - | - | + | - | - | - |
| | <i>Monoraphidium terrestre</i> cf. <i>dybowskii</i> | - | - | - | - | - | + | - | - | - |
| | <i>Mychonastes</i> cf. <i>homosphaera</i> | - | - | - | - | - | + | - | - | - |
| | <i>Mychonastes</i> sp.* | - | - | - | - | - | + | - | - | - |
| | <i>Pseudomuriella</i> cf. <i>schumacherensis</i> | - | - | - | - | - | - | - | - | + |
| | <i>Scenedesmaceae</i> sp. | - | - | - | - | - | + | - | - | - |
| Ulvophyceae | <i>Desmochloris</i> cf. <i>halophila</i> | + | - | - | - | - | - | - | - | - |
| | <i>Dilabifilum printzii</i> ** | - | - | - | - | - | - | - | + | + |
| | <i>Hazenia mirabilis</i> * | - | - | - | - | - | - | + | - | - |
| | <i>Pseudendocloniopsis botryoides</i> * | - | + | - | - | - | + | - | + | + |
| | <i>Pseudendoclonium akinetum</i> * | - | - | - | - | - | - | - | - | + |

Legend to Table 1. The recovered species were distributed at eight sampling sites of the Deinschwanger creek (DB) and the pooled sample from the Westerhöfer creek (WB, from five sites). Numbers below sampling sites are the corresponding values [log IAP/KT] of calcite saturation index ($SI_{Calcite}$, for explanation see Arp et al. 2010) and the total number of species recovered per site. Two asterisks next to a species name indicate 100%, a single asterisk 99.9% sequence identity with a reference strain.

Phylogenetic analyses

To search for the closest neighboring relatives of our isolates their sequences were compared to those from reference strains at NCBI (<http://www.ncbi.nlm.nih.gov>) using BLASTn queries (Altschul *et al.* 1997). Only almost full neighboring 18S rRNA gene sequences were downloaded together with a selection of reference sequences to better represent the green algal classes Trebouxiophyceae, Chlorophyceae and Ulvophyceae as well as additional green algal lineages and aligned using MAFFT, ver. 6 (Kato and Toh 2008) available online at <http://mafft.cbrc.jp/alignment/server/index.html>. Three sequence data sets were constructed after

the alignments were manually refined using BioEdit (Hall 1999). The sequence data set of Trebouxioophyceae (**Fig. 1**) contained 162 sequences and was 1766 positions long with 781/562 variable/parsimony informative sites, that of Chlorophyceae (**Fig. 2, Fig. 3**) contained 238 sequences and was 1797 positions long with 938/653 variable/parsimony informative sites, and that of Ulvophyceae (**Fig. 4**) 88 sequences and was 1785 positions long with 953/813 variable/parsimony informative sites. The GTR+ Γ +I model was selected as the best fitting model of nucleotide substitution for all three sequence data sets as based on the AIC criterion using jModelTest 0.1.1 (Posada 2008). Phylogenetic trees were calculated using maximum likelihood with the program RAxML 7.0.4 (Stamatakis *et al.* 2008) and Bayesian phylogenetic inference with MrBayes 3.2.1 x64 (Ronquist *et al.* 2012). For the latter, two MCMC runs for three million generations each were employed with one cold and three heated chains with trees sampled every 100 generations. Confidence values for the obtained groups (internal edge support) were inferred from the rapid bootstrapping algorithm (100 replicates) as implemented in RAxML (Stamatakis *et al.* 2008) and from Bayesian posterior probabilities using MrBayes 3.2.1 x64 (Ronquist *et al.* 2012). Pairwise sequence similarities from p-distances were determined as an additional measure of the relatedness of our isolates to certain reference strains/sequences under the Kimura 2-parameter model with the program MEGA5 (Tamura *et al.* 2011).

Statistical analysis

The distribution of all major clades of green algae at studied sampling sites was investigated using multivariate ordination by Non-metric Multidimensional Scaling (NMDS); the analysis was conducted in PAST 2.06 (Hammer *et al.* 2001). The input data corresponded to a matrix similar as in **Tables 1 and 2** summarizing the presence/absence of isolates at particular sampling sites as identified by 18S rRNA gene sequence comparisons.

Table 2. Distribution of the recovered phylogenetic clades of Green algae.

| | | DB1 | DB2 | DB3 | DB4 | DB5 | DB6 | DB9 | DBS |
|-------------------|---------------------------------------|------|------|------|------|------|------|------|------|
| | | 1.19 | 1.15 | 1.11 | 1.09 | 1.09 | 1.11 | 0.45 | 0.01 |
| | | 2 | 2 | 2 | 2 | 2 | 5 | 2 | 4 |
| Trebouxioophyceae | Chlorellaceae clade | + | + | + | - | - | + | - | + |
| | <i>Choricystis/Botryococcus</i> clade | - | - | - | - | - | - | - | + |
| | Prasiolales | - | - | - | + | - | + | - | - |
| Chlorophyceae | Chlamydomonadales | + | - | + | - | + | + | - | - |
| | Sphaeropleales | - | - | - | - | - | + | + | - |
| Ulvophyceae | Ulotrichales | - | + | - | - | - | + | + | + |
| | Ulvaes | - | - | - | - | - | - | - | + |

Legend to Table 2. Seven phylogenetic clades of Chlorophyta were distributed at the eight sampling sites of the Deinschwanger creek (DB). Numbers below sampling sites are values [log IAP/KT] of calcite saturation index (SI_{Calcite}, for explanation see Arp *et al.* 2010) and the total number of clades recovered per site ("phylogenetic diversity").

Results

A total of 77 isolates have been obtained from both creeks (**Table S1**). Phylogenetic analyses of 18S rRNA gene sequences determined for each isolate distinguished 34 distinct lineages supposedly corresponding to species. They were distributed on three classes of green algae, the Trebouxiophyceae (14 species; **Fig. 1**), the Chlorophyceae (15 species; **Fig. 2**, **Fig. 3**) and the Ulvophyceae (5 species; **Fig. 4**). Forty-five isolates have been selected for phylogenetic and microscopic analyses in this paper (**Figs. 1-7**, **Table S1**, **Table S2**). The other isolates shared identical (partial) sequences and, therefore, were suspected to represent the same species as the selected 45 isolates. Morphological features of the 34 species (77 isolates) were investigated and are summarized in **Table S2**.

Diversity of Trebouxiophyceae

The recovered Trebouxiophyceae isolates were distributed on four major clades of the class (**Fig. 1**). Those isolates with a *Stichococcus*-like rod shaped morphology (**Fig. 5a**, **Fig. 5b**) were distributed in seven distinct lineages within the *Prasiola* clade. Based on high sequence similarities (99.9 or 100 % as determined from p-distances) and short genetic distances with named reference sequences in the phylogenetic analyses (**Fig. 1**) two closely related isolates, WB13 and WB74 (**Fig. 5a**), were identified as *S. bacillaris* and isolate WB69 as *S. mirabilis* (**Table 1**, **Table S3**). For the other *Stichococcus*-like isolates sequence similarities with named references were lower or no closer named references were available at all. Isolate WB38 was named *S. cf. deasonii* as its next named reference was *S. deasonii*. For six isolates which were distributed on four independent lineages of the *Prasiola* clade (**Fig. 1**) no close named relatives were available and, therefore, they were named *Stichococcus* sp. 1 (isolates DB6-27, WB65, and SAG 2407), *S. sp. 2* (isolate D4-2A), *S. sp. 3* (isolate SAG 2408) and *S. sp. 4* (isolate SAG 2406; **Fig. 5b**). Based on morphology all seven lineages may be assigned to *Stichococcus*, but in the 18S rRNA gene phylogenies the genus was paraphyletic with many other coccoid and filamentous members of the *Prasiola* clade (**Fig. 1**). The recovered seven lineages of *Stichococcus*-like Trebouxiophyceae may correspond to seven distinct species, but their assignment to a single genus *Stichococcus* is not adequate. The genus as currently circumscribed may in fact encompass several distinct genera.

The unicellular spherical trebouxiophytes (**Figs. 5c-5f**) represented three distinct lineages within the Chlorellaceae clade (**Fig. 1**). The two isolates RK52 (**Fig. 5c**) and D11-2 (**Fig. 5d**) exhibited relatively large rounded cells with cup-shaped chloroplast and a single pyrenoid with

starch grains attached to it, i.e. the *Chlorella* morphotype (**Table S2**). Both isolates had sequences highly similar to each other and their closest neighbors (99.9 % sequence similarity) were several unidentified *Chlorella* sp. strains. Both isolates represented the most frequently recovered green alga in our study, i.e. there was a total of 12 strains with almost identical 18S rRNA sequences and identical morphology representing the species (**Table S1**, **Table S2**). Resolution within the clade corresponding to the Chlorellales in the 18S rRNA phylogeny was low (**Fig. 1**). Both *Chlorella*-like isolates were distinct from the authentic (type) strain of the genus, *C. vulgaris* SAG 211-11b, by a relatively large genetic distance (**Fig. 1**), but a closer assignment to any genus of Chlorellales was impossible. *Hindakia* was the closest relative (**Fig. 1**), but with a relatively low sequence similarity of 99.7 %.

Two other trebouxiophyte species with spherical cells differed from the *Chlorella* morphotype by their smaller cells with a single band-shaped chloroplast without pyrenoid (**Table S2**; **Fig. 5e**, **Fig. 5f**). They were distributed in two distinct lineages outside of Chlorellales, but within the clade representing the Chlorellales (**Fig. 1**). Two isolates (D6-DB2 and SAG 2390) were assigned to *Muriella terrestris* (**Fig. 5e**) due to low p-distances/high sequence similarities (99.9 and 100 %) with strain ASIB V38 (acc. no. AB012845) which has been isolated from soil (Gärtner 1996). Both our isolates also had the same high sequence similarity with an unidentified strain of *Muriella* AS2-4 from freshwater phytoplankton, acc. no. AY195969 (Fawley *et al.* 2004). Therefore, we assume that both our isolates as well as strains ASIB V38 and AS2-4 represent the same species, *M. terrestris*. The other species, represented by isolate WB67 (**Fig. 5f**), shared high sequence similarities with unidentified *Nannochloris*-like strains which together shared a well-supported monophyletic origin with *Marvania geminata*. Because of the relatively low sequence similarity (97.8 %) with the reference sequence of *M. geminata* our isolate probably represents another yet still undescribed species of *Marvania*.

Other isolated trebouxiophytes exhibited reniform (*Neocystis*-like; **Fig. 5g**), elliptic to nearly spherical (*Elliptochloris*-like; **Fig. 5h**) or elongated to elliptic (*Coccomyxa*-like; **Fig. 5i**) cell shapes. The next relative to isolate SAG 2405, characterized by its mucilaginous colonies and reniform cell shape (**Fig. 5g**), was *Neocystis mucosa* strain KR 1989/14 (acc. no. HM565928), therefore it was termed *Neocystis* cf. *mucosa*. Another *Neocystis* strain, *Neocystis brevis* CAUP D802 (acc. no. HQ287929), was slightly more distant to SAG 2405 and the whole clade of *Neocystis*-like trebouxiophytes was highly supported. Isolate WB5-D1e (**Fig. 5h**) was identified as *Elliptochloris subsphaerica* based on its high sequence similarity of 99.9% with *E. subsphaerica* strain SAG 2202 (acc. no. FJ648518). The same short p-distance/high sequence similarity was with unidentified *Elliptochloris* strain SAG 2117 (acc. no. FJ648515). The whole

clade comprising the three *Elliptochloris* sequences was well supported (**Fig. 1**). Two more species exhibited a typical *Coccomyxa*-like morphology (**Fig. 5i**) and were nested within a well-supported clade consisting of several *Coccomyxa* strains. The one species, represented by the single isolate WB40, was most closely related to strain *Pseudococcomyxa simplex* UTEX 274 (acc. no. FJ648514). The other species was represented by four isolates, retrieved from both creeks (**Table S1**). Out of them for isolate WB28 an almost full 18S rRNA gene sequence was determined and strain *Coccomyxa pringsheimii* SAG 69.80 (acc. no. AY762603) represented its closest neighboring sequence.

Diversity of Chlorophyceae

Isolates assigned to the class Chlorophyceae were distributed in two clades representing the orders Chlamydomonadales and Sphaeropleales (**Fig. 2**, **Fig. 3**). The isolated members of Chlamydomonadales exhibited two morphological types, i.e. *Chlorococcum*-like large spherical cells with a single large chloroplast (**Figs. 6a-6e**), *Chlamydomonas*-like monadoid biflagellated cells (**Fig. 6f**), those of Sphaeropleales four morphological types, i.e. *Scenedesmus*-like elongated cells with acute ends (**Figs. 6h-6i**), *Bracteacoccus*-like spherical cells with numerous discoidal parietal chloroplasts without a pyrenoid (**Figs. 7a, 7b**), *Mychonastes*- or *Nannochloris* like small spherical cells with a single chloroplast without a pyrenoid (**Figs. 7c, 7d**), and *Monoraphidium*-like fusiform cells.

Chlamydomonadales

The *Chlorococcum*-like isolates were distributed in four distinct clades within the larger and highly supported *Stephanosphaeria* clade in 18S rRNA gene phylogenies (**Fig. 2**). One clade was formed by isolate RK50 and strain SAG 2402 and their named relative *Chlamydropodium vacuolatum* (acc. no. M63001) and, therefore, both were assigned to the genus *Chlamydropodium*. Another sequence, but from an unidentified strain, *Chlamydomonadaceae* sp. KMMCC:EC-34, was even closer with the former two strains (**Fig. 2**). A second well supported clade included the isolates GRK6-DB5, GRK6-DB6, SAG 2400 and SAG 2401. Except for the latter two, which had almost identical sequences, they were all distant to each other and had sequences from unidentified strains as their next closest relatives. The next named closest relative was *Chlorococcum ellipsoideum* strain UTEX 972 (acc. no. U70586) and, therefore, we named our isolates *Chlorococcum ellipsoideum*-relatives (**Fig. 3**, **Figs. 6c-e**). *C. ellipsoideum*-relative1, isolate GRK6-DB5, was closer related to *C. ellipsoideum* UTEX 972 than the other three isolates of the clade. A third clade contained a single isolate, SAG 2398, which was within a well-

supported clade together with several named closest relatives (**Fig. 2**). Except for *Chlorococcum sphacosum* SAG 66.80 (which corresponds to the authentic strain of the type of the species), our isolate was morphologically rather distinct from them and, therefore, we assigned isolate SAG 2398 to *C. sphacosum* (**Fig. 6a**; **Table 1**, **Table S2**). Both strains, SAG 2398 and SAG 66.80 (acc. no. JN968580), also shared 99.9% sequence similarity (**Table 1**, **Table S3**). Finally, strain SAG 2399 (**Fig. 6b**) was nested within another distinct and highly supported clade with *C. vacuolatum* SAG 213-8 (acc. no. KF144189) and *C. minutum* ASIB T50 (acc. no. JN968585) as the closest next named relatives (**Fig. 2**). Because it appeared more similar in its morphology to *C. minutum* (**Table S2**) we named strain SAG 2399 *Chlorococcum minutum*-relative. The *Chlamydomonas*-like monadoid isolates RK68 (**Fig. 6f**) and DB6-shared identical (partial) sequences and isolate RK68 was within the well-supported *Reinhardtinia* clade in the phylogenetic analyses (**Fig. 2**). Its closest named next relatives were three species of *Chlamydomonas* and *Volvox carteri*. Because it exhibited solitary cells with a typical *Chlamydomonas* morphology we assigned it to that genus, but could not identify it to species level.

Sphaeropleales

The *Scenedesmus*-like isolates were distributed in three independent lineages (species) of the *Scenedesmaceae* clade (Sphaeropleales; **Fig. 3**). *Desmodesmus* cf. *armatus* was with six isolates among the most frequently recovered green algal species of our study (**Table S1**). Both isolates, RK43 (**Fig. 6g**) and strain SAG 2395, for which almost full 18S rRNA gene sequences were determined, formed a highly supported clade together with named reference strain, *Desmodesmus armatus* CCAP 276/4A (acc. no. FR865727) and, therefore, were assigned the isolates to this species (**Fig. 3**). Isolate D22-6-2B (**Fig. 6h**) shared an identical sequence with reference strain *Acutodesmus obliquus* CCAP 276/49 (acc. no. FR865726) and was assigned to this species. Finally, isolate RK49 (**Fig. 6i**) could not be assigned to a certain genus of the *Scenedesmaceae* clade because it was distinct from all named reference strains of that clade, albeit there was weak support for a closer relationship with species of *Acutodesmus* (**Fig. 3**). The three *Bracteacoccus*-like isolates were distributed in two independent clades of the Sphaeropleales representing the genera *Bracteacoccus* and *Pseudomuriella* (**Fig. 3**). The next closest named reference for isolate RK3 (**Fig. 7a**) was *Pseudomuriella schumacherensis* SAG 2137 (acc. no. HQ292768) but with a sequence similarity < 99.9% and, therefore, we named our isolate *P. cf. schumacherensis*. Despite their morphological similarities with the *P. cf. schumacherensis* isolate the two other *Bracteacoccus*-like isolates, SAG 2403 and DB9-3 (**Fig. 7b**), were in a distinct clade of the 18S

rRNA gene phylogeny, i.e. the *Bracteococcaceae* clade. Strain SAG 2403 had a close named relative, *Bracteococcus aerius* UTEX 1250 (**Fig. 3**), but shared less than 99.9% sequence similarity from p-distances with the latter and, thus, was named *Bracteococcus* cf. *aerius*. In contrast, isolate DB9-3 had no named closest relative; it was most closely to an unidentified *Bracteococcus* strain (KF-2011f), but rather distant from *B.* cf. *aerius* (**Fig. 3**). Both *Mychonastes/Nannochloris*-like isolates, RK48 and DB6-29 (**Figs. 7c-d**), were within a single well supported clade, representing the genus *Mychonastes* within the Chlorophyceae (**Fig. 3**). Isolate RK48 had *Mychonastes homosphaera* CAUP H6501 (which represents the authentic strain of the species) as its closest named relative, but with a sequence similarity of less than 99.9% (**Table S3**) and, therefore, was named *Mychonastes* cf. *homosphaera*. *Mychonastes*-like isolate DB6-29 was distant from RK48 and had an unidentified *Mychonastes* strain, Itas 9/21 14-8w (acc. no. AY543066) as closest relative (with 99.9% sequence similarity) and, therefore, was named *Mychonastes* sp. (**Fig. 3**). Several species of *Mychonastes* have recently been described as common members of freshwater phytoplankton (Krienitz *et al.* 2011), but our isolates were more distant to those and, therefore, were not included in our phylogenetic analyses. The *Monoraphidium*-like creek biofilm isolate, strain SAG 2393, was most closely related (with sequence similarity < 99.9%) to two reference strains named *M. dybowskii* within the *Selenastraceae* clade and, therefore, was named *M.* cf. *dybowskii*.




Figure 1. Maximum-likelihood (ML) phylogeny of 18S rRNA gene sequences of green algal species from the class Trebouxiophyceae isolated from biofilms of two freshwater creeks (highlighted sequence names) and other members of the Trebouxiophyceae as references. Additional members of Oocystaceae (Trebouxiophyceae) as well as other classes of the Chlorophyta, the Chlorophyceae, Ulvophyceae and Chlorodendrophyceae (outgroup), are included for comparisons and to root the tree (for their accession numbers see Table S4). Additional outgroups including *Picocystis salinarum* and *Nephroselmidophyceae* were removed from the figure due to lack of space (all outgroups were the same as in Fig. 2, for accession numbers see Table S4). Numbers at internal branches indicate support from bootstrap tests using ML (left) and posterior probabilities from Bayesian analysis (BI; right). Thick lines mark internal branches that were resolved by both, the ML and the BI tree topologies. Grey boxes highlight the clades comprising the new isolates.

Diversity of Ulvophyceae

Five species from creek biofilms were members of the Ulvophyceae. Interestingly, out of them four were of pseudo-filamentous organization, i.e. formed only short filaments and also coccoid stages, whereas only one exhibited a purely coccoid vegetative stage. This was in contrast to our isolates of Trebouxiophyceae and Chlorophyceae which were all of coccoid or monadoid organization. The pseudo-filamentous isolates exhibited characteristic cup-shaped chloroplasts with one or two prominent pyrenoids (**Figs. 7h-i**). Three filamentous species were members of the order Ulotrichales, one belonged together with the coccoid isolate to the Ulvales (**Fig. 4**). Filamentous strain SAG 2396 (**Fig. 7e**) had a rather small genetic distance to *Hazenia mirabilis* UTEX LB 846 (acc. no. AF387156) and, therefore, was assigned to this species (**Fig. 4**). *Pseudendoclonium basiliense* UTEX 2593 and two species of *Gloeotilopsis* formed up a clade with the former two strains which, however, received only insignificant support values (**Fig. 4**). Another pseudo-filamentous species we recovered was *Pseudendocloniopsis botryoides*; it was with nine isolates (with identical partial sequences and morphology) the second most frequently retrieved green algal species (**Table S1**). As represented by isolate SAG 2394 the close relationship with two available strains of *Pseudendocloniopsis botryoides* was well supported and, therefore, SAG 2394 was assigned to this species (**Fig. 4**). SAG 2394 may also form coccoid stages (**Fig. 7f**), a feature shared with *Planophila laetevirens* SAG 2008 which also was its next closest relative in the 18S rRNA gene phylogeny (**Fig. 4**). Morphology of *P. botryoides* was studied at isolate DB6-19 in more detail (**Fig. 7g**; **Table S2**). The filamentous isolate SAG 2404 (**Fig. 7h**) was a very closely relative to *Pseudendoclonium akinetum* UTEX 1912 and also two species of *Trichosarcina* were their next closest relatives, but these relationships did not receive significant support values (**Fig. 4**). Finally, for another pseudo-filamentous species, *Dilabifilium printzii*, a total of seven isolates was established (**Table S1**); it was the third most frequently recovered green algal species in our study. Unfortunately, we failed to obtain an almost full 18S rRNA gene sequence for these isolates. The partial sequence of isolate WB41 shared 99.9% similarity with the sequence (acc. no. KF144198) of *D. printzii* strain SAG 467-1 and, therefore, we assigned our isolate to this species. Morphological features of isolate WB41 are presented in **Fig. 7i** and **Table S2**. *D. printzii* SAG 467-1 had an independent position within the clade representing the Ulvales in our 18S rRNA gene phylogeny (**Fig. 4**). The single coccoid Ulvophyceae isolate SAG 2397 clearly fell within a subclade of the Ulvales comprising the coccoid genera *Chlorocystis*, *Desmochloris* and *Halochlorococcum* (**Fig. 4**). Strain SAG 2397 had short genetic distances with two strains of *Desmochloris halophila* and, therefore, we




Figure 5. Morphology of green algal isolates of the class Trebouxiophyceae. *Stichococcus bacillaris* isolate WB74 (a), *Stichococcus* sp.4 SAG 2406 (b), *Chlorella* sp. RK52 (c), *Chlorella* sp. D11-2 (d), *Muriella terrestris* D6-DB2 (e), *Marvania* sp. WB67 (f), *Neocystis* cf. *mucosa* SAG 2405 (g), *Elliptochloris subsphaerica* WB5-D1e (h), *Coccomyxa* cf. *pringsheimii* WB32 (i). Scale bars = 10 μ m.




Figure 6. Morphology of green algal isolates of the class Chlorophyceae. Chlamydomonadales (a) – (f); Sphaeropleales (g) – (i). *Chlorococcum sphacosum* SAG 2398 (a), *Chlorococcum minutum*-relative SAG 2399 (b), *Chlorococcum ellipsoideum*-relative1 GRK6-DB5 (c), *Chlorococcum ellipsoideum*-relative2 SAG 2400 (d), *Chlorococcum ellipsoideum*-relative2 GRK6-DB6, zoospores (e), *Chlamydomonas* sp. isolate RK68 (f), *Desmodesmus* cf. *armatus* isolate RK43 (g), *Acutodesmus obliquus* isolate D22-6-2B (h), *Scenedesmaceae* sp. isolate RK49 (i). Scale bars = 10 μm .




Figure 7. Morphology of green algal isolates of the classes Chlorophyceae and Ulvophyceae isolates. Chlorophyceae, Sphaeropleales (a) – (d); Ulvophyceae (e) – (i). *Pseudomuriella* cf. *schumacherensis* RK3 (a), *Bracteacoccus* sp. DB9-3 (b), *Mychonastes* cf. *homosphaera* RK48 (c), *Mychonastes* sp. isolate DB6-29 (d), *Hazenia mirabilis* strain SAG 2396 (e), *Pseudendoconiopsis botryoides* SAG 2394, coccoid stage (f), *Pseudendoconiopsis botryoides* DB6-19, filamentous stage (g), *Pseudendoclonium akinetum* SAG 2404 (h), *Dilabifilum printzii* isolate WB41 (i). Scale bars = 10 μ m.

Discussion

Distribution of the isolates at the sampling sites

In total, we detected nearly the same numbers of green algal species in the Deinschwanger (DB; 21) and Westerhöfer (WB; 18) creeks, regardless of the unequal number of sites sampled per creek (8 at the DB, 5 at the WB; **Table 1**). With respect to calcification the various sampling sites of the DB did not cluster according to high or low calcification when using their species compositions (**Fig. 8**). We conclude that we observed a rather accidental distribution of the biofilm microalgae which is not or only very little influenced by calcification. This is contrast to findings of Brinkmann *et al.* (2015) (this issue) who reported the diversity of cyanobacteria and diatoms of the same creek biofilms as studied here was clearly influenced by SI_{calcite} and pCO_2 which are reciprocally linked. The highest diversity per site (11 species) was detected at site DB6 which was highly calcified (**Table 1**). However, from the other calcified sampling sites (DB1-DB5) only 1-3 species per site could be retrieved in culture. At the spring site DBS with calcification almost absent also just four species were recovered. Two of them, *Coccomyxa* cf. *pringsheimii* and *Dilabifilum printzii* were retrieved only at the DB spring site, but *D. printzii* was recovered from the highly-calcified sites of the WB as well (**Table 1**). In addition, we compared the various DB sites for their “phylogenetic diversity”, i.e. whether the species recovered from a certain DB site belong to a single or several phylogenetically distinct clades. With respect to the number of phylogenetic clades retrieved per site, DB6 again was the most diverse site and the spring site DBS was the second most diverse site (**Table 2**). The Chlamydomonadales clade (Chlorophyceae; **Fig. 2**) seemed to have a preference towards the highly calcified DB sites, whereas the Chlorellales (Trebouxiophyceae; **Fig. 1**) and the




Figure 8. Ordination diagram. Nonmetric Multidimensional Scaling (NMDS) plot showing the grouping of high-calcified sampling sites (black symbols) based on presence/absence of the green algal lineages.

Ulotrichales (Ulvophyceae; **Fig. 4**) clades were found at the highly calcified sites as well as the spring site of the DB (**Table 2**). With respect to the three different green algal classes detected in both karst-water creeks, the WB was dominated by members of Trebouxiophyceae (12 out of the total of 18 species detected there belonged to Trebouxiophyceae),




Figure 9. Bar chart showing the relative representation of the three classes of Chlorophyta at the both investigated creeks. Black, Chlorophyceae; light grey, Trebouxiophyceae; dark grey, Ulvophyceae.

whereas the DB predominately harbored members of Chlorophyceae (12 out of the total of 21 species detected there belonged to Chlorophyceae). For the few members of Ulvophyceae no preference to any of the both creeks was found. Interestingly, out of the 34 detected green algal species only six were retrieved from both creeks, i.e. *Coccomyxa* cf. *pringsheimii*, *Muriella terrestris* and *Stichococcus* sp.1 of the Trebouxiophyceae, *Chlorococcum ellipsoideum*-relative2 of the Chlorophyceae, and *Dilabifilum printzii* and *Pseudendocloniopsis botryoides* of the Ulvophyceae (**Fig. 9**). It follows that the species richness of green algae at both creeks was rather different (**Fig. 9**).

Species identification and origins

The molecular phylogenetic diversity of inland freshwater green algae from running waters has been studied for the first time here. In particular, for estimating the diversity of microscopic green algae in epilithic biofilms of running freshwaters, studies are rare and all were exclusively based on morphology so far (for review see Lindstrøm *et al.* (2004) and Veselá (2006)). The majority of existing evidence about the molecular diversity of freshwater green microalgae refers to communities rather different from algal biofilms, e.g. phytoplankton (Fawley *et al.* 2004; Krienitz and Bock 2012). Other molecular data for the diversity of green algae from of running freshwaters have mainly been available from habitats with extreme environmental variables or artificial impact (e.g., Dorigo *et al.* 2002; Aguilera *et al.* 2007; Palacios *et al.* 2008; Baker *et al.* 2009; Aguilera *et al.* 2010). For example, molecular data are available for the periphyton of the extremely cold and remote Antarctica (De Wever *et al.* 2009; Vyverman *et al.* 2010). Compared to freshwater creeks, terrestrial habitats have more extensively been studied by a combined morphological/molecular approach to date (Lewis and Lewis 2005; Flechtner *et al.* 2013).

Sequence comparisons of 18S rRNA genes enable the unambiguous comparisons of new isolates to those previously isolated from other localities and habitats world-wide (De Wever *et al.* 2009; Němcová *et al.* 2011; Hodač *et al.* 2012). Using sequence comparisons most of our

isolates were closely related to those of various other inland freshwaters (**Table S3**). Within the Trebouxiophyceae, these were members of the Chlorellales clade, i.e. *Chlorella* (Bock *et al.* 2010; Bock *et al.* 2011), *Muriella* (Fawley *et al.* 2004) and *Marvania* (Fawley *et al.* 2004; Yamamoto *et al.* 2007). Within the Chlorophyceae, at least the *Scenedesmaceae*, *Selenastraceae* and *Mychonastes*, which we also retrieved in our study, are well known from freshwaters (Krienitz *et al.* 2011; Krienitz and Bock 2012). However, all these genera and families of green algae are known predominantly from major water bodies like lakes or ponds where they inhabit free-floating communities (phytoplankton) and, therefore, were not expected to occur in biofilm assemblages as well. While Trebouxiophyceae and many Chlorophyceae are predominately found in terrestrial and freshwater habitats, Ulvophyceae is a green algal class with a main divergence in marine habitats; it represents the only lineage of Chlorophyta that includes macroscopic sea-weeds (Leliaert *et al.* 2012). In contrast, our isolates were nested within the orders Ulotrichales and Ulvales which are known to occur also in freshwaters and terrestrial habitats as well. Interestingly, the characteristically branched filamentous thalli of the majority of our ulvophycean isolates strongly resembles the morphology of *Gongrosira* which has frequently been reported from tufa formations (Pentecost 1988; Pentecost and Spiro 1990; Freytet and Plet 1991; Johnson and John 1992). Therefore, the name *Gongrosira* may represent a collective morphotype whose members are distributed at least on several distant lineages within the Ulotrichales (Ulvophyceae). Moreover, the only so far sequenced strain of *Gongrosira*, *G. papuasica* UTEX 1916, has been identified as a member of Chlorophyceae (López-Bautista and Chapman 2003).

Using 18S rRNA gene sequence comparisons only for six out of the total of 34 species retrieved in our study had 100% and 99.9% identities with already available sequences from reference strains (**Table 1**, **Table S3**). Therefore, we suspect to have retrieved the same species as previous studies in these cases. However, for none of these species authentic strains were available and, therefore, our identification (species names) may be only as correct as the closest named neighbors (reference strains) of our isolates have been correctly named to species. Except for two reference strains where their origin is not known and *Elliptochloris subsphaerica*, all reference strains for these species have been isolated from various freshwater habitats (**Table S3**) and from a broad range of geographical regions, e.g. Europe, United States and Antarctica. Consequently, our findings support the world-wide distribution of these species. However, it is important to note that most of these freshwater species have been reported from open water bodies as phytoplankton so far, but are recovered by our study from submerged biofilms attached to stones in running waters for the first time. In other cases closest relatives to our isolates were

from terrestrial habitats. Our finding of *E. subsphaerica* is the first record of this species from a freshwater habitat; previously it has been reported only from terrestrial habitats (Ettl and Gärtner 1995). Similarly, our still unidentified isolates of *Stichococcus*, WB38 (*S. cf. deasonii*), SAG 2408 (*S. sp.3*) and SAG 2406 (*S. sp.4*), have their closest relatives from terrestrial habitats. Species of *Coccomyxa* have mostly been reported from terrestrial habitats (e.g. tree bark or lichen symbiosis) so far, but were recovered from creek biofilm in our study. The latter three examples indicate that at least parts of the creek biofilm algal communities were probably driven into their habitat through (rain-) flushes from the neighboring soils. For most other biofilm species which so far were reported from phytoplankton only it is still not known whether they can be found in terrestrial habitats as well. *Muriella terrestris* may be an example for a green microalgal species that is known from phytoplankton as well as soil; it was frequently found in the creek biofilms as well.

For 24 species we recovered no identification at the species level was possible, i.e. they were left unnamed despite they could be assigned to a certain genus in most cases. These represent species for which no 18S rRNA gene sequence have become available yet or which have been retrieved for the first time. In this respect we see the importance of molecular-phylogenetic investigations of inland freshwater algal biodiversity, i.e. to extend our knowledge about the distribution of species of green microalgae across various freshwater and terrestrial habitats.

Acknowledgements

We are indebted to Dr. Anke Behnke and Ms. Viviane Kipp who initiated this study through their culturing efforts and developed green algal PCR primer LR1650. This work was supported by the German Science Foundation (DFG) by a grant extended to T.F. (Fr 905/13-2). Parts of this work were also supported by the German Federal Ministry of Education and Research, BMBF (AlgaTerra project, grant 01 LC 0026) within the BIOLOG program.

Chapter 3

Phylogenetic analysis of polar *Chlorella* and *Stichococcus* suggests biogeography of airborne microalgae

Ladislav Hodač¹, Christine Hallmann¹, Karolin Spitzer¹, Josef Elster^{2,3}, Fabian Faßhauer¹, Nicole Brinkmann⁴, Daniela Lepka¹, Vaibhav Diwan¹ and Thomas Friedl¹

¹Experimental Phycology and Culture Collection of Algae (SAG), University of Göttingen, Germany

²Centre for Polar Ecology, University of South Bohemia, České Budějovice, Czech Republic

³Institute of Botany, Phycology Centrum, Academy of Sciences of the Czech Republic, Třeboň, Czech Republic

⁴Department of Forest Botany, University of Göttingen, Germany

(Submitted to *FEMS Microbiology Ecology*; FEMSEC-15-11-0606)

Abstract

Chlorella and *Stichococcus* are morphologically simple airborne microalgae, omnipresent in terrestrial and aquatic habitats. The minute cell size and resistance against environmental stress facilitate their long-distance dispersal. However, the actual distribution of *Chlorella*- and *Stichococcus*-like species was so far inferred from ambiguous morphology-based evidence. Here we contribute a phylogenetic analysis of an expanded SSU and ITS2 rDNA dataset including new accessions from terrestrial habitats of polar regions, the temperate zone and the tropics in order to determine whether related species of airborne microalgae show similar patterns in geographic distribution. We found that psychrotolerant strains of *Chlorella* and *Stichococcus* are without exception conspecific (or closely related) with strains originating from the temperate zone. Seven phylogenetic species of polar Chlorellaceae have been uncovered so far with five of them being our new psychrotolerant strains. *Chlorella vulgaris* and *Muriella terrestris* show the widest distribution, including polar and hot desert environments. The morphologically more diversified *Stichococcus*-like lineages were so far detected not only in polar regions and hot deserts, but also in tropical rainforests. However, distinct *Stichococcus* clades exhibit either temperate-polar or temperate-tropical distributions. Our data suggest that terrestrial airborne microalgae might exhibit biogeography and are not distributed everywhere.

Keywords: biogeography, *Chlorella*, *Stichococcus*, polar strains

Introduction

Terrestrial species of *Chlorella* (Beijerinck 1893) and *Stichococcus* (Nägeli 1849) are true survivalists among the green microalgae (Chlorophyta). They inhabit biofilms covering natural and artificial subaerial substrates and dwell in soils (Carson and Brown 1976; 1978; Ettl and Gärtner 1995; Sharma *et al.* 2007; Rindi *et al.* 2009). *Chlorella* and *Stichococcus* have been reported by microscopical observations from nearly all soil types, including desert soil crusts in Namibia (Büdel *et al.* 2009), humid tropical soils of India (Ray and Thomas 2012), Central America (Archibald 1972), Oceania (Arvik and Willson 1974; MacEntee *et al.* 1977; Carson and Brown 1978) and polar desert soils in Antarctica (Cavacini 2001; Fermani *et al.* 2007) and the Arctic (Elster *et al.* 1999; Kaštovská *et al.* 2005; Kaštovská *et al.* 2007; Patova and Dorokhova 2008). Langhans *et al.* (2009) recognized species of *Chlorella* and *Stichococcus* as key players for monitoring the succession of biological soil crust formation. Particular attention has been paid to the metabolic facilities of psychrotolerant terrestrial strains of *Chlorella vulgaris* and *Stichococcus bacillaris* (Kvíděrová and Lukavský 2005; Shukla *et al.* 2011; Chen *et al.* 2012; Hong *et al.* 2015); both species include strains with the potential for use in biotechnology (Lang *et al.* 2011; Olivieri *et al.* 2011; Cadoret *et al.* 2012; Barreiro *et al.* 2013; Olivieri *et al.* 2013; Goiris *et al.* 2014; Mudimu *et al.* 2014; Safi *et al.* 2014; Sivakumar *et al.* 2014; Slocombe *et al.* 2015). The resistance against environmental stresses connected with metabolic versatility is a common feature of polar microalgae, which have to cope with low temperatures and shortages of nutrients and liquid water (Elster 1999; Elster and Benson 2004; Kvíděrová and Lukavský 2005; Sharma *et al.* 2007). An extreme multi-stress resistance was detected in a psychrotolerant cryptoendolithic strain of *Stichococcus* (isolated from Antarctica), which survived an experimental exposure to an Earth orbital space environment (Scalzi *et al.* 2012).

Systematically, '*Chlorella*' is the name for minute coccoid microalgae phylogenetically nested within the Chlorellaceae clade, class Trebouxiophyceae (Huss *et al.* 2002; Krienitz *et al.* 2003; Luo *et al.* 2010; Pröschold *et al.* 2010; Leliaert *et al.* 2012; Krienitz *et al.* 2015). Whereas planktonic Chlorellaceae evolved into distinct forms, e.g., *Micractinium* (Pröschold *et al.* 2010; Krienitz and Bock 2012) or *Actinastrum* (Luo *et al.* 2010; Krienitz and Bock 2012), terrestrial members exhibit morphological convergence, which is characteristic for “true” *Chlorella* (*Chlorella* clade, Bock *et al.* 2011) and for multiple *Nannochloris*-like clades (Henley *et al.* 2004). Morphologically simple *Chlorella*- and *Nannochloris*-like species were repeatedly uncovered in the Antarctic (Gilichinsky *et al.* 2007; Hu *et al.* 2008; De Wever *et al.* 2009).

'*Stichococcus*' can be as well encountered in the harsh terrestrial and freshwater environments of the Antarctic and Arctic (De Wever *et al.* 2009; Vishnivetskaya 2009; Khan *et al.* 2011).

'*Stichococcus*' is the common name for multiple rod-shaped species phylogenetically nested within the *Prasiola* clade (Handa *et al.* 2003; Neustupa *et al.* 2007; Novis *et al.* 2008; Eliáš and Neustupa 2009). The *Prasiola* clade represents a trebouxiophyte lineage with the highest morphological diversity within the class (Leliaert *et al.* 2012), which is widely distributed in freshwaters, marine and terrestrial ecosystems (Karsten *et al.* 2005; Rindi *et al.* 2007). Despite their morphological simplicity, *Stichococcus* species veil a considerable phylogenetic diversity, which is still in need of taxonomic revision (Neustupa *et al.* 2007; Sluiman and Guihal 2008; Karbovska and Kostikov 2012a). The rod-shaped *Stichococcus* species are scattered across the whole *Prasiola* clade, but are not intermixed with morphologically distinct lineages such as *Pseudomarvania* (Eliáš and Neustupa 2009) or the thallos Prasiolales (Handa *et al.* 2003; Rindi *et al.* 2004; Rindi *et al.* 2007). SSU-based indications, that the *Prasiola* clade and the Chlorellaceae might represent sister lineages (Handa *et al.* 2003; Fučíková *et al.* 2014), appear to be questionable, while the *Prasiola* clade is undoubtedly nested within the core Trebouxiophyceae (Leliaert *et al.* 2012; Lemieux *et al.* 2014; Turmel *et al.* 2015), plastid genome phylogenies place Chlorellaceae outside the core Trebouxiophyceae (Lemieux *et al.* 2014; Leliaert and Lopez-Bautista 2015; Turmel *et al.* 2015).

Chlorella, *Stichococcus* and a handful of other green algal morphospecies are counted among the so called airborne algae (Sharma *et al.* 2007; Sharma and Rai 2010), which are supposed to be ubiquitous in terrestrial ecosystems (Rindi *et al.* 2009; Rindi *et al.* 2011). However, the existence of limitedly dispersed cryptic species which are morphologically hard to determine, became the rule since the introduction of molecular techniques into phycology (Boenigk *et al.* 2005; Rindi *et al.* 2008; Dal Grande *et al.* 2014; Řídká *et al.* 2014; Ryšánek *et al.* 2014; Škaloud *et al.* 2014a; Škaloud *et al.* 2015). Nevertheless, some species might be true cosmopolitans, as recently affirmed within some lineages of terrestrial green microalgae, e.g., *Coccomyxa* (Darienkov *et al.* 2015), *Klebsormidium* (Ryšánek *et al.* 2014) and *Diplosphaera* (Fontaine *et al.* 2012). Remote polar regions like Antarctica and the Arctic provide an opportunity for investigating isolation-by-distance and speciation of microorganisms (Martiny *et al.* 2006; Hahn *et al.* 2015). Pioneering molecular studies on algal diversity in Antarctica found genetic divergence between polar and non-polar species; some detected *Chlorella* and *Stichococcus* species were supposed to be Antarctic endemics (Lawley *et al.* 2004; De Wever *et al.* 2009). Some five years later, the amount of sequenced algal strains and environmental clones has dramatically increased, encouraging us to reconsider whether the Antarctic endemism might be

simply due to a sampling effect. We avoid the difficulty of the algal species concept (Leliaert *et al.* 2012; Leliaert *et al.* 2014); instead, we infer distribution patterns based on monophyletic clades of closely related sequences. By focusing on the polar strains of *Chlorella* and *Stichococcus*, we aim to show that related species of airborne microalgae have similar patterns of geographic distribution.

Materials and Methods

Data sampling and microscopy of the strains

We analyzed new SSU and ITS2 rDNA sequences obtained from the strains, isolates and clones listed in **Table S1** (*Chlorella*-like sequences, accession numbers XY000000-XY000000) and **Table S2** (*Stichococcus*-like sequences, accession numbers XY000000-XY000000). The geographic origin of new sequences from polar/temperate/tropical regions is shown in **Fig. 1**. Isolation of the algal strains and sequencing were described in Hodač *et al.* (2015) while PCR-cloning of the algal sequences from environmental samples was described in Hallmann *et al.* (2013a). Observations of the cultures were conducted using an Olympus BX60 microscope (Tokyo, Japan) with Nomarski DIC optics and an attached ColorView III camera (Soft Imaging System, Münster, Germany). Cell size measurements (diameter of spherical cells; length/width of elongated cells) were conducted in ImageJ (Schneider *et al.* 2012) and were based on a set of 50-100 cells per culture.




Figure 1. Geographic origin of the new *Chlorella* and *Stichococcus* 18S and/or ITS2 sequences. 1-5: *Chlorella*-like accessions (circles); 1-Canada (Ellesmere Island), 2-Svalbard, 3-Antarctica (Killingbeck Island), 4-Antarctica (Adelaide Island), 5-Antarctica (King George Island), 6-Antarctica (Anchorage Island), 7-German Biodiversity Exploratories. 8-20: *Stichococcus*-like accessions (squares); 8-German Biodiversity Exploratories, 9-Croatia, 10-Canada (New Scotia), 11-USA (Massachusetts), 12-USA (Alabama, Dauphin Island), 13-USA (Alaska), 14-Ecuador (Podocarpus National Park), 15-Antarctica (Victoria Land), 16-New Zealand (Wawaira Scenic Res.), 17-Indonesia (Bali, Lake Bratan), 18-Malaysia (Kampong Kuala Jenera), 19-Japan (Taishakukyo Gorge), 20-Antarctica (Marie Bird Land). Blue symbols represent polar accessions. Detailed information on sampling sites is listed in **Tables S1, S2**.

18S rDNA phylogenetic analyses

The closest relative sequences to the new accessions were searched for in GenBank using the BLAST algorithm (Altschul *et al.* 1997). New and the GenBank accessions were checked for chimeras using Bellerophon (Huber *et al.* 2004). Two separate 18S rDNA alignments were created, one for *Stichococcus*-like accessions (*Prasiola* clade) and one for *Chlorella*-like (Chlorellaceae) accessions. Both 18S rDNA sequence alignments were computed using MAFFT v.6 (Kato and Toh 2008) available online. The aligned sequences were checked for possible misaligned positions in BioEdit 7.0.9.0 (Hall 1999). The 18S rDNA alignment, which included the new *Stichococcus* accessions and the closest relatives from GenBank (*Prasiola* clade), contained 100 sequences/1709 positions (276 variable, 149 parsimony informative). Based on the AIC criterion in jModelTest 0.1.1 (Posada 2008), the GTR+ Γ +I nucleotide substitution model was selected as the best fitting the *Prasiola* clade. A maximum likelihood phylogeny was computed in RAxML 7.0.4 (Stamatakis *et al.* 2008) under the proposed model and statistical support values were derived from rapid bootstrapping (1000 replicates) in the same program. An alternative maximum likelihood tree search procedure, i.e., quartet puzzling, was applied for the same 18S alignment using Tree-Puzzle 5.2 (Strimmer and Von Haeseler 1996) available at the Moby-Pasteur webserver (Néron *et al.* 2009). For this, the GTR+ Γ substitution model was applied. To assess the putative age of dichotomies within the *Prasiola* clade, relative node ages were estimated using BEAST v1.8.2 (Drummond and Rambaut 2007; Drummond *et al.* 2012) with 5 000 000 steps (after 500 000 burn-in) under the relaxed molecular clock option. Absolute ages were estimated following De Wever *et al.* (2009) by setting the minimum and maximum age of the Chlorophyta–Streptophyta split at 700 and 1500 Ma. A tree was visualized using FigTree (Rambaut 2007). The 18S rDNA alignment of Chlorellaceae comprised 82 sequences/1668 positions (350 variable, 226 parsimony informative). The J2:G:5 substitution model was proposed by Treefinder (Jobb *et al.* 2004), which was subsequently used for maximum-likelihood tree reconstruction. Confidence values were inferred from 1000 bootstrap replicates in the same program. The resulting tree was visualized using MEGA5 (Tamura *et al.* 2011). For additional statistical support, Bayesian posterior probabilities were computed in MrBayes 3.2.1 x64 (Ronquist *et al.* 2012). We carried out two MCMC runs for three million generations each with one cold and three heated chains under the GTR+ Γ +I evolutionary model (parameters were estimated from the data); trees were sampled every 100 generations. To summarize the genetic similarities within the alignments, the sequences were clustered into operational taxonomic units (OTUs) in the program MOTHUR v.1.13.0 (Schloss *et al.* 2009) implementing the 0.00 and 0.01 thresholds under exclusion of the gap positions. For sequence comparisons, p-distances were

computed in MEGA6 (Tamura *et al.* 2013).

ITS2 rDNA phylogeny and secondary structure analysis

First, 63 new ITS2 rDNA sequences were submitted to the BLAST search (Altschul *et al.* 1997) in order to obtain their closest relatives for phylogenetic comparison. Subsequently, an online ITS2 database (Schultz *et al.* 2006; Selig *et al.* 2008; Koetschan *et al.* 2010; Koetschan *et al.* 2012) was used for ITS2 rRNA annotation (Keller *et al.* 2009). Minimum energy secondary structure models were computed from the annotated sequences using RNAstructure 5.3 (Reuter and Mathews 2010). For very closely related sequences, a few were folded in RNAstructure and the rest modelled using homology modelling in the ITS2 database (Wolf *et al.* 2005). Ambiguous models were compared to those computed in the mfold webserver (Zuker 2003). The ITS2 rRNA secondary structures were visualized by Varna 3.8 (Darty *et al.* 2009). Subsequently, two separate alignments of *Chlorella vulgaris* and *Stichococcus*-like sequences and structures were built using 4SALE 1.7. (Seibel *et al.* 2006; Seibel *et al.* 2008). In the same program, compensatory base changes (CBC) were computed between pairs of sequences. The Chlorellaceae ITS2 alignment contained 39 sequences/321 positions (58 variable, 39 parsimony informative). A statistical parsimony analysis was conducted with the very closely related ITS2 sequences of *Chlorella vulgaris* and *C. pituita* using TCS v.1.21 (Clement *et al.* 2000) to generate ribotype networks. The connection limit was set to 95%. The *Prasiola* clade ITS2 alignment contained 56 sequences/584 positions (252 variable, 189 parsimony informative). Based on sequence-structure alignment, a neighbor-joining tree was computed in ProfDist (Müller *et al.* 2004; Friedrich *et al.* 2005; Rahmann *et al.* 2006; Wolf *et al.* 2008) using the GTR+ Γ model and bootstrapping (1000 replicates). A splitting structure among closely related ITS2-ribotypes was reconstructed by neighbor-net analysis in SplitsTree4 4.10 (Huson and Bryant 2006), applying uncorrected p-distances and with ambiguities handled as averages. Bootstrap support values for internal splits were calculated with 1000 replicates. The assignation of sequences into ribotypes was computed in DnaSP v.5 (Librado and Rozas 2009).

Results

Polar Chlorellaceae and allies

The analyzed polar *Chlorella*-like strains exhibit two different gross morphologies: 1) *Chlorella*-like spherical to slightly elliptical cells ($\varnothing = 2.8\text{--}8.4\ \mu\text{m}$) bearing one cup-shaped chloroplast with one (**Fig. 2a**) or two (**Fig. 2b**) pyrenoids; 2) *Nannochloris*- and *Muriella*-like spherical cells (\varnothing

= 2.2-7.3) with a simple parietal chloroplast without a pyrenoid (**Figs. 2d-f**). According to the 18S-based phylogenetic analysis, the polar strains are nested in four different operational taxonomic units (OTUs; corresponding to genera) of the Chlorellaceae (**Fig. 3**). The OTUs 1-4 include further GenBank accessions, which are closely related to our polar strains at the $\geq 99.5\%$ sequence similarity level.




Figure 2. Microphotographs of polar strains of the Chlorellaceae. (a) *Chlorella* cf. *vulgaris* L5 ($\varnothing = 2.8-5.7 \mu\text{m}$); (b) *Chlorella* sp.1 N9 ($\varnothing = 2.9-5.4 \mu\text{m}$); (c) *Chlorella* sp.2 L3 ($\varnothing = 3.4-8.4 \mu\text{m}$); (d) *Muriella* sp. L20 ($\varnothing = 2.2-6.3 \mu\text{m}$); (e) *Marvania* relative L15 ($\varnothing = 2.3-5.8 \mu\text{m}$); (f) Unidentified Chlorellaceae L24 ($\varnothing = 2.6-7.3 \mu\text{m}$). Scale bars = $10 \mu\text{m}$.

***Chlorella* clade**

The Antarctic strains L1 and L4 (King George Island) and the Arctic strains L5 (**Fig. 2a**) and L6 (Ellesmere Island) show $\geq 99.5\%$ sequence similarity to the authentic strain *C. vulgaris* SAG 211-11b. The 18S phylogeny suggests divergence between the polar strains and the remaining mostly temperate sequences within OTU3 (**Fig. 3**). We analyzed the observed differences between the polar and temperate strains of *C. vulgaris* using more variable ITS2 (**Fig. 4a**), assigning them into 13 ribotypes (A, B, E-O), exhibiting a total variability of 54 polymorphic nucleotide positions (**Fig. S1a**). The ribotypes A-K differ from *C. vulgaris* SAG 211-11b by a few nucleotide polymorphisms (**Fig. S1a**). In contrast, ribotypes L-O are more divergent from *C. vulgaris* SAG 211-11b; they differ from both SAG 211-11b and ribotypes A-K by sharing some identical nucleotide sites with *C. pituita* ACOI 311 (**Fig. S1a**). The ITS2 secondary structure

model (**Fig. S1b**) shows that a considerable amount of nucleotide positions are conserved across all *C. vulgaris* ribotypes (including the polar ribotypes) and *C. pituita* (**Fig. S1b**). Moreover, none of the ribotypes (not even the polar ones) shows any compensatory base changes (CBCs) against *C. vulgaris* SAG 211-11b, but all show at least three CBCs against *C. pituita* ACOI 311 (**Fig. S2a**). Notably, the Antarctic strain L4 (ribotype M) differs in only a single nucleotide from the German isolate LH10HG2067 (ribotype L). Otherwise, the same ribotypes were detected from both aquatic and terrestrial habitats (i.e., ribotypes B, H, I; **Fig. 4b**). Ribotype B is the most common ITS2 variety of *C. vulgaris* in GenBank, including strains isolated from freshwaters in Europe, USA and China.

The Arctic strain N9 (**Fig. 2b**) and the Antarctic strain L3 (**Fig. 3c**) morphologically resemble members of the *Chlorella* clade. Considering the 18S phylogeny, both strains are nested within an unnamed clade (OTU1), which comprises closely related strains from freshwaters of the temperate northern hemisphere and Antarctica. The 18S inference points out the relatedness to the strain *Hindakia fallax* CCAP 222/29. However, the ITS2 phylogeny does not support the close relationship (**Fig. 4a**; **Fig. S2b**), since both strains exhibit considerable genetic divergence from both *Hindakia* species. The ITS2 inference further suggests that N9 and L3 represent two different species, provisionally denominated as *Chlorella* sp.1 and *C. sp.2*, respectively. We detected a remarkably high ITS2 similarity between the *Chlorella* sp.1 strain N9 from the Arctic and the strain KNUA034 from Antarctica, differing by three nucleotide positions (and a one-sided/hemi compensatory base change/hCBC in the helix IV, **Fig. 4c**). The *Chlorella* sp.2 strain L3 has no known close relatives.

***Nannochloris*-like clades**

The *Nannochloris*-like Antarctic strains L20 (**Fig. 2d**) and N5 are nested within OTU3, comprising *Muriella terrestris* ASIB V38 and other terrestrial and aquatic strains of almost identical 18S sequences (**Fig. 3**). The high 18S similarity of the strains L20 and N5 contrasts with their considerable genetic dissimilarity in the ITS2 marker (**Fig. 4**; **Fig. S2c**). The Antarctic *Muriella* sp.1 strain L20 exhibits high ITS2 similarity with the Antarctic clone Ant 8/104 and the isolate LH08SG3009 from a German soil. Strain L20 differs from the isolate LH08SG3009 by one hCBC in the helix III of the ITS2 secondary structure (**Fig. 4d**).

The Antarctic strains L13 and L15 (**Fig. 2e**) clustered within the OTU4 comprising further 18S accessions from Antarctica and the temperate regions (**Fig. 3**). The phylogenetically closest named species are *Marvania geminata* SAG 12.88 and *Nannochloris coccoides* CCAP 251/1b. However, any close relationship to OTU4 (provisionally denominated as *Marvania* relative1) is

supported neither in the 18S nor the ITS2 phylogeny (**Fig. 4a**, **Fig. S2d**).

Strain L24 (King George Island) represents a phylogenetically isolated lineage within the Chlorellaceae, so far without known close relatives. The strain L24 (**Fig. 2f**) exhibits morphology resembling *Nannochloris*-like lineages rather than the true *Chlorella* described above.




Figure 3. 18S ML-phylogeny of the polar *Chlorella*-like strains and relatives. All polar accessions are highlighted in blue and authentic strains are marked by a '*<*' sign. The numbers next to branches indicate statistical support values (maximum-likelihood bootstraps (ML)/Bayesian posterior probabilities (BI)); the clades of particular interest were additionally tested via maximum parsimony (MP) and bio-neighbor-joining (NJ) and the statistic support values are given in the following order: ML/BI/MP/NJ. Black bars to the right side give sequence assignments into OTUs at $\geq 99.5\%$ similarity level. Drawings show morphology characteristic for each clade.




Figure 4. Analysis of ITS2 sequences obtained from polar *Chlorella*-like strains and relatives. (a) An ITS2-based neighbor-joining tree computed from sequences and secondary structures. (b) Ribotype network of *Chlorella vulgaris* accessions assigned to ribotypes A–O. Each circle represents a distinct ITS2 ribotype and its size reflects the number of identical sequences. (c)–(d) Consensual ITS2 secondary structures of closely related accessions: (c) *Chlorella* sp.1, (d) *Muriella* sp. Nucleotide positions within the ITS2 secondary structures are colored as follows: grey dots=conserved positions, orange dots=nucleotide substitutions and black dots=nucleotide deletions.

Stichococcus-like lineages

The newly obtained strains of *Stichococcus*-like species all exhibited characteristic rod-shaped morphology (**Fig. 5a–l**) and varied in both cell length (1.3–10.0 μm) and width (1.0–4.4 μm). Deviations from the rod-shape morphotype—*Diplosphaera*-like cell packages—were observed in the isolate LH08SW1099 (**Fig. 5j**). The *Stichococcus* strains do not substantially differ in chloroplast shape and the detectability of pyrenoids using light microscope is ambiguous. The 18S analysis of all *Stichococcus*-like accessions (**Table S2**) revealed five monophyletic *Stichococcus*-like OTUs of highly similar sequences ($\geq 99.5\%$), i.e., OTU1, OTU3–6 (**Fig. 6; Fig S3**) and a further four *Stichococcus*-like OTUs of lower resolution $\geq 99.0\%$, i.e., OTU2, OTU7–9 (**Fig. 6; Fig S3**). The OTUs could be better resolved using more polymorphic ITS2 sequences (**Fig. 7; Fig. S4a**) and analyzing their secondary structures (**Fig. 7; Fig. S4b; Fig. S5**); this approach revealed a total of 12 unambiguously distinguishable monophyletic lineages (**Tab. S3**).




Figure 5. Microphotographs of the *Stichococcus*-like strains isolated from Germany and Ecuador. (a) *Stichococcus* clade2 LH08AW8104 (l = 4.0-10.0 µm; w = 1.8-3.2 µm); (b) *Stichococcus* clade7 LH10HG2063 (l = 4.5-8.7 µm; w = 1.8-2.8 µm); (c) *Diplosphaera* sp. LH08HW8075 (l = 3.2-6.2 µm, w = 1.9-4.4 µm); (d) *Diplosphaera* sp. KS120SM6L (l = 1.3-4.9 µm, w = 1.9-3.4 µm); (e) *Pseudostichococcus* sp. LH08SW8044 (l = 3.4-9.8 µm; w = 1.6-3.3 µm); (f) *Stichococcus* clade1 SAG 2481 (l = 3.1-8.7 µm; w = 1.6-2.9 µm); (g) *Stichococcus* clade1 KS075SM6T (l = 2.9-6.3 µm; w = 1.5-3.0 µm); (h) *Stichococcus* clade3 KS106CL6T (l = 2.5-5.0 µm; w = 1.5-2.4 µm); (i) *Stichococcus* clade5 LH08SW1099 (l = 3.0-6.3 µm; w = 1.0-4.2 µm); (j) *Stichococcus* clade5 KS305SM6L (l = 3.3-6.8 µm; w = 1.4-3.2 µm); (k) *Stichococcus jenerensis* KS126SM6L (l = 1.9-5.0 µm; w = 1.6-3.2 µm); (l) *Stichococcus* clade4 SAG 2406 (l = 4.5-8.2 µm; w = 1.7-2.7 µm). Scale bars = 10 µm.

***Stichococcus*-like clades including polar species**

The polar *Stichococcus*-like accessions (obtained from GenBank; **Table S2**) represent at least three different phylogenetic lineages, here provisionally designed as “*OTU2 Stichococcus deasonii* & allies”, “*OTU8 Diplosphaera*” and “*OTU9 Pseudostichococcus*” (**Fig. 8**; **Fig. S3**).

OTU2 consists of the authentic strain *Stichococcus deasonii* SAG 2139 (Alabama, USA) and further accessions originating from terrestrial habitats (**Table S2**). *OTU2* is poorly resolved in the 18S phylogenies (**Fig. 6**; **Fig. S3**). The ITS2 analyses suggest that, apart from *Stichococcus deasonii*, *OTU2* consists of another three different species, here provisionally named as *Stichococcus* clade2 (**Fig. 5a**), *S.* clade6, and *S.* clade7 (**Fig. 5b**). *S.* clade2 includes accessions from Antarctica (e.g., HM490287), differing from its European relatives (e.g., SAG 2482) by 6-9 nucleotide positions in ITS2 (**Fig. S5**). Another two Antarctic strains cluster within *OTU2*: *S. bacillaris* NJ-10 and *S. bacillaris* s3, but the 18S data do not provide sufficient resolution to assign them into any known species. *Stichococcus deasonii* is the only lineage within *OTU2* which comprises accessions from the tropics (e.g., clone KSK870SM6T; **Fig. S5**).

OTU8 Diplosphaera is a phylogenetically heterogeneous clade consisting of multiple lineages of *Stichococcus*-like (*S. chodati* UTEX 1177), *Diplosphaera*-like (*D. mucosa* SAG 48.86) and *Chlorella*-like (*C. sphaerica* SAG 11.88) species, recognizable in the 18S and/or ITS2 phylogenies, yet without sufficient resolution or statistical support. The Antarctic strain *Diplosphaera mucosa* SAG 48.86 is the only known polar member within *OTU8*. Further accessions within the clade originate from Central Europe (isolate LH08HW8075; **Fig. 5c**; **Fig. 6**) the tropics (isolate KS120SM6L; **Fig. 5d**; **Fig. 6**) and North-American deserts (e.g., *Stichococcus chlorelloides* BCP-CNP2-VF11B). *OTU8* is the only known *Stichococcus*-like clade that includes species from hot deserts (**Fig. 8**).

OTU9 Pseudostichococcus represents a phylogenetically isolated clade of *Stichococcus*-like (*S. mirabilis* CCAP 379/3, *Pseudostichococcus monallantoides* SAG 380-1) and *Desmococcus*-like (*D. spinocystis* SAG 2067) species (**Fig. 7**; **Fig. 8**; **Fig. S3**; **Fig. S5**). The clade consists of two strains isolated from Antarctica: Trebouxiophyceae sp. EO7-4 and *Stichococcus minutus* NJ-17. All other accessions within *OTU9* originate from terrestrial (e.g., *Pseudostichococcus* sp. LH08SW8044; **Fig. 5e**) and aquatic habitats of the temperate zone while no tropical relatives are known so far.

***Stichococcus*-like clades including tropical species not related to polar allies**

We identified four clades consisting of tropical and temperate species and designed them provisionally as: “*OTU1 Stichococcus* clade1”, “*OTU3 Stichococcus* clade3”, “*OTU5*

Stichococcus clade5” and “*OTU7 Stichococcus jenerensis*” (**Table S2**).

OTUs 1, 3 and 5 are statistically highly supported in both the 18S (**Fig. 6**) and ITS2 (**Fig. 7**; **Fig. S5**) phylogenies and represent three undescribed *Stichococcus*-like species. Few nucleotide differences in the ITS2 within the clades and no compensatory base changes further support close relationships among the tropical and European accessions. OTU1 *Stichococcus* clade1 consists of strains isolated from Germany (SAG 2481; **Fig. 5f**) and Ecuador (KS075SM6T; **Fig. 5g**). Another German strain KS075SM6T differs from KS075SM6T by only 10 nucleotide positions within ITS2 and exhibits no compensatory base changes (**Fig. S5**). OTU3 *Stichococcus* clade3 includes the European strain SAG 2408 which differs by only three ITS2 nucleotide positions from the tropical isolate KS106CL6T (**Fig. 5h**). OTU3 further consists of 18S accessions sampled in the Swiss Alps and Yellowstone NP (**Fig. 6**; **Table S2**). OTU5 *Stichococcus* clade5 is a terrestrial lineage consisting of the European isolate LH08SW1099 (exhibiting both *Stichococcus*- and *Diplosphaera*-like morphology; **Fig. 5i**) and the Ecuadorian isolate KS305SM6L (**Fig. 5j**). The tropical isolate differs by four ITS2 nucleotide substitutions from the isolate KP09AW1004 originating from Germany. OTU5 might be distributed as well in Hawaii, since the accession KM462543 shows a high 18S sequence similarity (99.46%) to the members of *Stichococcus* clade5.

OTU7 Stichococcus jenerensis contains the authentic strain *S. jenerensis* SAG 2138, which is the only *Stichococcus*-like species described from the tropics. The ITS2 data suggest a pantropical distribution of the clade: the Ecuadorean isolate KS126SM6L (**Fig. 5k**) differs by 13 ITS2 nucleotides from the Southeast-Asian strain SAG 2138, and exhibiting no CBCs (**Fig. S5**). *OTU7* further consists of temperate terrestrial and even marine accessions (**Fig. 6**; **Table S2**).

***Stichococcus*-like clades from the temperate zone**

Two highly supported clades of terrestrial and aquatic *Stichococcus*-like species are known so far only from the temperate zone: “*OTU4 Stichococcus* clade4” and “*OTU6 Stichococcus bacillaris*”. OTU4 contains the *Stichococcus*-like strain SAG 2406 (**Fig. 5l**) and further accessions (sampled in Germany; **Table S2**) with almost identical 18S sequences (**Fig. 6**). However, the ITS2 data point out a cryptic species diversity, since we detected one CBC between strain SAG 2406 and the isolate LH08SG1073 (**Fig. S5**). The strain CCALA 906 from Svalbard (KF355941) is the closest polar relative of OTU4, yet with low 18S similarity (98.7%). OTU6 includes multiple *Stichococcus* strains (e.g., *S. bacillaris* SAG 379-1b; *S. chloranthus* SAG 379-2) and strains morphologically assigned to *Gloeotila* (*Gloeotila scopulina* SAG 335-8; *Gloeotila* cf. *protogenita* SAG 56.91). OTU6 accessions originate from freshwaters (**Table S2**), soil (clone

HEW1B K3375; **Fig. 6**) and even the Southeast Pacific (KF899844). The ITS2 data suggest a close relationship between the OTU6 *Stichococcus bacillaris* and the *Desmococcus* clade (**Fig. 7; Fig. S5**), which consists of cell packages-forming/filamentous terrestrial species (distributed in the polar and temperate regions; **Fig. 8**). OTU6 and *Desmococcus* are sister lineages in both the 18S and ITS2 phylogenies and differ by two CBCs in the ITS2 secondary structures.




Figure 6. 18S ML (quartet puzzling) phylogeny of *Stichococcus*-like species and relatives. The numbers next to branches (italic letters) indicate bootstrap support values inferred from 1000 replicates. The numbers in white boxes close to clades studied in detail correspond to statistical values in the following order: ML/BI/MP/NJ. Schematic drawings within clades show cell size of representative *Stichococcus*-like strains as compared to *Stichococcus bacillaris* morphospecies after Nägeli (1849) and Ettl & Gärtner (1995). Assignations into operational taxonomic units (OTUs) are based on sequence similarities $\geq 99.5\%$ (= OTU) or $\geq 99.0\%$ (= OTU). The clades are colored according to geographic origin of the sequences: blue=polar/temperate, green=tropical/temperate, grey=only temperate or temperate/polar/tropical.




Figure 7. ITS2 phylogenetic network of *Stichococcus*-like species and relatives. Neighbor-net analysis of *Stichococcus* clades. The numbers near to splits correspond to bootstrap support values. Schematic ITS2 secondary structures are shown for representative strains. Assignations into operational taxonomic units (OTUs) are based on 18S inference (Fig. 6; Table S2). The comprehensive dataset is shown in Fig. S5. The clades are colored according to geographic origin of the sequences: blue=polar/temperate, green=tropical/temperate, grey=only temperate or temperate/polar/tropical.




Figure 8. Calibrated 18S phylogeny of *Stichococcus*-like species and relatives using relaxed molecular clock. Bayesian phylogenetic tree (BEAST) computed from 18S dataset identical to **Fig. 6**. Colored dots to the right of the tree summarize information on the geographic distribution of *Stichococcus*-like species and relatives (based on molecular data listed in **Table S2**). Schematic drawings illustrate the multiple occurrence of rod-shaped (*Stichococcus*-like) morphotypes throughout the *Prasiola* clade. Assignations into operational taxonomic units are based on sequence similarities $\geq 99.5\%$ (= OTU) and $\geq 99.0\%$ (= OTU), **Fig. 6**.

Discussion

We defined operational taxonomic units (= clades consisting of identical or closely related sequences) as those with $\geq 99.5\%$ SSU-sequence similarity. The most psychrotolerant strains of *Chlorella* and *Stichococcus* are conspecific (or closely related) with strains from the temperate zone. Including our new sequences and the sequences published in GenBank (De Wever *et al.* 2009), we recognized seven phylogenetic species of polar Chlorellaceae. Five of these species are the psychrotolerant Chlorellaceae isolated by Shukla *et al.* (2011). Such a low phylogenetic diversity of *Chlorella* is also known from arid regions in which six species have been reported so far (Lewis and Lewis 2005; Flechtner *et al.* 2013; Fučíková *et al.* 2014). Our results suggest that particularly two species of the Chlorellaceae, *Chlorella vulgaris* (Bock *et al.* 2011) and *Muriella terrestris* (Hanagata 1998), might be distributed in both polar and arid desert environments. Terrestrial *Stichococcus*-like microalgae might be even more widely dispersed than members of the Chlorellaceae, because in addition to the polar and temperate regions, they were detected in the tropics as well. However, six *Stichococcus* clades with a temperate-polar distribution do not include any tropical relatives. And vice versa, *Stichococcus* clades with a temperate-tropical distribution do not contain any psychrotolerant species.

***Chlorella*-like psychrotolerant strains have allies in temperate zones and hot deserts**

The strains morphologically resembling *Muriella* (Petersen 1935) are phylogenetically nested in *Muriella terrestris* (Hanagata 1998). We detected a considerably low genetic distance—a difference of three nucleotides within ITS2—among the *M. terrestris* strains sampled in Antarctica (L20; Shukla *et al.* 2011; Kochkina *et al.* 2014) and Germany (LH08SG3009). Their putative conspecificity is supported by the absence of compensatory base changes (CBCs; Müller *et al.* 2007b; Wolf *et al.* 2013). Further, SSU-based evidence suggests a widespread distribution of *M. terrestris* in streams of high CO₂-pressure (Hodač *et al.* 2015), freshwaters (Fawley *et al.* 2004) and even in hot deserts (Flechtner *et al.* 2013; Fučíková *et al.* 2014). Another *Nannochloris*-like clade includes the Antarctic isolates L13/L14/L15/L23/L32 (Shukla *et al.* 2011), which are weakly related to *Marvania geminata* (Henley *et al.* 2004; Eliáš and Neustupa 2009). These strains cluster together with another Antarctic *Nannochloris* sp. Ant-1 (Gilichinsky *et al.* 2007), with *Chlorella* sp. 193-GA188 from the Siberian permafrost (Vishnivetskaya 2009) and with multiple isolates from German soils (e.g., LH08SG2053) and from freshwater (*Nannochloris* sp. JL 4-6; Fawley *et al.* 2004). The relatives of the tropical species *Hindakia fallax* (Bock *et al.* 2010) exhibited low genetic distance (three nucleotides in ITS2) between the isolate N9 from

Svalbard (Shukla *et al.* 2011) and the Antarctic isolate KNUA034 (Hong *et al.* 2015). The Arctic isolate N9 is phylogenetically nested in a clade containing other accessions provisionally denominated as *Chlorella* spp., a putative sister lineage to the tropical *Hindakia fallax*. *Hindakia* spp. was already recorded multiple times from Antarctica (Hu *et al.* 2008; De Wever *et al.* 2009; Hong *et al.* 2015) and includes freshwater strains, which were screened for application in biotechnology, e.g., CCAP 211/79 (Germond *et al.* 2013; Osundeko *et al.* 2013; Driver *et al.* 2015). High ITS2 similarity between isolates from Nova Scotia (KP204587; Park *et al.* 2015) and the South China Sea (GU942201), which are both relatives of the Antarctic isolate L3 (Shukla *et al.* 2011), additionally supports the long-distance dispersal of *Chlorella* spp.

Chlorella vulgaris, the epitome of spherical green microalgae (Krienitz *et al.* 2015), was isolated from Antarctica and represented by the intensively investigated psychrotolerant strain NJ-7 (Li *et al.* 2009; Lu *et al.* 2009; Lu *et al.* 2010). The SSU sequences of our Antarctic (L1/L4) and Arctic (L5/L6) isolates (Shukla *et al.* 2011) clustered between *C. vulgaris* SAG 211-11b and its closest named relative *C. pituita* ACOI 311 (Bock *et al.* 2011). The Antarctic strains are more different from the temperate *C. vulgaris* SAG 211-11b than the Arctic strains; the most similar strain to the Antarctic strains L1/L4 was isolated from a hot desert (JX446471; Flechtner *et al.* 2013) and a German soil (LH10HG2081; this study). ITS2 analysis of multiple *C. vulgaris* strains revealed considerable intraspecific diversity; the particular ITS2 ribotypes do not cluster according to their original habitat (freshwater vs. terrestrial). Remarkably, the ITS2 of the Antarctic strain L4 differs from all *C. vulgaris* strains, but is almost identical to the isolate LH10HG2081. Taking the ITS2 difference between *C. vulgaris* SAG 211-11b and *C. pituita* ACOI 311 (32 nucleotides and 3-4 CBCs) into account, then the psychrotolerant (L4) and *C. vulgaris* SAG 211-11b might be conspecifics (differing in 14 nucleotides; 0 CBC). In comparison, another trebouxiophycean genus *Coccomyxa* exhibits intraspecific variability of 26 nucleotide positions in ITS2 (Darienکو *et al.* 2015). Evidence of identical ITS2 ribotypes from Europe (SAG 211-11b) and North America (CCAP 254/5) confirms the long-distance dispersal of *C. vulgaris*. *Chlorella sorokiniana* (Bock *et al.* 2011) is another remarkable *Chlorella*, so far known from both cold and hot environments, e.g., the Antarctic (De Wever *et al.* 2009) and North-American deserts (Flechtner *et al.* 2013). In contrast to the terrestrial *Chlorella*-like species mentioned above, *C. sorokiniana* might be dispersed even in the humid tropics and was recorded (based on DNA data) from Central America (de-Bashan *et al.* 2008), South America (de-Bashan *et al.* 2008; Bashan *et al.* 2015) and Southeast Asia (identified by S. Marimuthu; <http://studentsrepo.um.edu.my/3562/>). The species is thermotolerant (de-Bashan *et al.* 2008; Zheng *et al.* 2013). Other thermotolerant Chlorellaceae recorded from hot deserts have close

relatives from phytoplankton, e.g., *Meyerella* (Fučíková *et al.* 2014), *Micractinium* (Flechtner *et al.* 2013); the distribution of the recently described *Chlorella thermophila* (Ma *et al.* 2015) remains unknown. Apart from the psychrotolerant strains described above, another two polar strains, i.e., L24 (Shukla *et al.* 2011) and VPL1-3 (FJ946890; De Wever *et al.* 2009) are incertae sedis in the Chlorellaceae.

***Stichococcus*-like clades including psychrotolerant species and species from hot deserts**

The psychrotolerant strains of *Stichococcus* which have been sequenced so far (Hughes 2006; Chen *et al.* 2012) cluster within three different lineages of the *Stichococcus* clade sensu Sluiman and Guihal (2008). We provisionally denominated these clades as *Pseudostichococcus* (SAG 2067; SAG 380-1; CCAP 379/3), *Stichococcus* clade2 (SAG 2482) and *S. clade7* (SAG 2060; SAG 2119). The *Pseudostichococcus* clade (including the strain *P. monallantoides* SAG 380-1; Moewus 1951) is an early diverging lineage consisting of species morphologically resembling *Stichococcus* (Nägeli 1849) and *Desmococcus* (Brandt and Stockmayer 1925; Gärtner and Ingolić 2003). Two Antarctic strains are nested within the *Pseudostichococcus* clade: Trebouxioephyceae EO7-4 (De Wever *et al.* 2009) and *Stichococcus minutus* NJ-17 (Chen *et al.* 2012). The freshwater strain EO7-4 is genetically similar to German isolates from soil (LH08SW8044; this study) and a karst-water stream (WB69; Hodač *et al.* 2015). The allies of the subaerial isolate NJ-17 were detected as environmental clones from German soils and isolated from a lichen (Thüs *et al.* 2011). Further Antarctic strains—phylogenetically distant from the *Pseudostichococcus* clade—were isolated from permafrost (*Stichococcus bacillaris* s3; Hughes 2006) and wet rocks (*S. bacillaris* NJ-10; Chen *et al.* 2012) and belong to a poorly resolved clade provisionally denominated as *Stichococcus* clade7. As a result of the SSU-based phylogeny, the species within the *Stichococcus* clade7 might be allies of *S. deasonii* (Neustupa *et al.* 2007). The ITS2-based inference, however, did not unambiguously support the monophyletic origin of both clades, suggesting a considerable evolutionary distance between *S. clade7* (including the above mentioned polar species) and the *S. deasonii* lineage (including tropical species). Notably, the *Stichococcus* clade7 further consists of species from soils and subaerial habitats (Karsten *et al.* 2005), but, in contrast to most *Stichococcus*-like clades, no aquatic species are known so far. The only psychrotolerant *Stichococcus*-like strain obtained from the Arctic permafrost is *Stichococcus* sp. 594-GA18 (Vishnivetskaya 2009) isolated from a 4.65 m depth in Siberia. This strain clusters within the *Stichococcus* clade2, an unnamed monophyletic lineage closely related to the above mentioned *S. clade7*. *Stichococcus* clade7 contains German isolates from a karst-water stream (D4-2A; Hodač *et al.* 2015) and soils (e.g., SAG 2482). Remarkably, *S. clade7* is the only known

Stichococcus-like clade which might have a bipolar distribution; it includes environmental clones (HM490287, HM490288; Khan *et al.* 2011) from the extreme McMurdo Dry Valleys, Antarctica, which are highly similar to the above mentioned soil strain SAG 2482 (6-9 nucleotide differences in ITS2, no CBCs).

The phylogeny of Chlorellaceae in hot deserts might be higher than that of *Stichococcus*, which is limited to a single known clade. This clade including desert species we denominated *Diplosphaera*, based on the species *D. mucosa* (Broady 1983) and *D. sphaerica* (Karbovska and Kostikov 2012b). *Diplosphaera mucosa* SAG 48.86 is the only polar strain within the *Diplosphaera* clade, however, the clade contains another putatively psychrotolerant species, represented by the environmental clone QE28 detected in the Tibetan tundra (FJ790654; Wong and Lacap 2010). The members of the *Diplosphaera* clade inhabit not only extremely cold and hot deserts (Lewis and Lewis 2005; Flechtner *et al.* 2013), but also acidic freshwater (Aguilera *et al.* 2007), oceanic sediment (AB183601) and occur as common lichen photobionts (Thüs *et al.* 2011; Fontaine *et al.* 2012; Fontaine *et al.* 2013). *Diplosphaera* ITS2 sequences, which were retrieved from leaf surfaces in an Ecuadorian tropical rainforest (e.g., KS120SM6L), were almost identical (i.e., differing in only three nucleotides; no CBC) to the isolate LH08HW8075 (this study) from a German forest soil. The SSU data suggest close relatedness of both temperate-tropical strains, the Antarctic *Diplosphaera mucosa* SAG 48.86 and lichen photobiont *Diplosphaera* sp. J4028B (Thüs *et al.* 2011). In *Stichococcus* clade7, we also observed genetic relatedness of a lichen photobiont (*Diplosphaera* sp. W1118; Thüs *et al.* 2011), tropical clone KSK870SM6T and polar isolates (e.g., *Stichococcus bacillaris* NJ-10; Chen *et al.* 2012), however, this evidence is based on lower genetic similarities as compared to the *Diplosphaera* clade.

Biogeography of terrestrial *Chlorella* and *Stichococcus*

Patterns in species distribution have been accepted for freshwater microalgae (De Wever *et al.* 2009; Naselli-Flores and Padisák 2015). Some freshwater species of “true” *Chlorella*, which were described from the tropics (e.g., *C. pulchelloides*, *C. rotunda*, *C. singularis*, *C. volutis*; Bock *et al.* 2011), have not yet been uncovered in polar regions. Furthermore, the *Chlorella*-like species we detected in Antarctica and the Arctic (*C. vulgaris*, *C. spp.*, *Muriella terrestris*, *Marvania*-relative) belong to clades with temperate-polar distributions; they are genetically divergent from their tropical relatives (e.g., *Hindakia fallax* or *Chlorella sorokiniana*). However, concerning terrestrial green microalgae, Earth’s biodiversity hotspots such as tropical rainforests still remain largely unexplored. Morphology-based observations suggest that true *Chlorella* (and other

Chlorellaceae) might be less common in tropical rainforests than *Stichococcus* (Neustupa and Škaloud 2008; 2010). Accordingly, the most *Chlorella*-like species, which were described from tropical rainforests using molecular data, belong either to the trebouxiophycean *Watanabea* clade (Zhang *et al.* 2008; Neustupa *et al.* 2009; Suutari *et al.* 2010; Song *et al.* 2015) or to the class Chlorophyceae (e.g., Jenufa; Němcová *et al.* 2011; Hodač *et al.* 2015). Preliminary cloning studies (unpublished data of the coauthors of this study) on the diversity of terrestrial green microalgae in Ecuadorian rainforests revealed only *Chlorella* sequences related to *C. sorokiniana*, a putatively pantropic and thermotolerant species. Remarkably, common terrestrial species such as *Chlorella vulgaris* or *Muriella terrestris* were not detected, although the approach did uncover common species of *Stichococcus*, which were almost identical to isolates from European soils and freshwaters. The *Stichococcus* species with temperate-tropical distribution were provisionally denominated as *Stichococcus* clade1, *S.* clade3 and *S.* clade5, and do not contain any psychrotolerant strains. The same is true for the first *Stichococcus* described from the tropics, *S. jenerensis* SAG 2138 (Southeast Asia; Neustupa *et al.* 2007). The discovery of *S. jenerensis* as well in Ecuador (KS126SM6L) confirmed the pantropic-temperate distribution of the *S. jenerensis* clade.

The deepest divergence within the *Stichococcus* clade (Sluiman and Guihal 2008), or *Stichococcus/Prasiola* clade (Neustupa *et al.* 2007) occurred in the Mesozoic, during the transition from the Lower Jurassic to Middle Cretaceous (De Wever *et al.* 2009). According to the same study, the deepest splits within the Chlorellaceae would be dated much earlier, i.e., approximately in the Lower Devonian to Lower Jurassic. The earliest divergence of the Antarctic versus non-Antarctic species within the *Chlorella* clade would thus have happened during the Cenozoic (Upper Paleocene to Upper Oligocene). Our accordingly calibrated phylogeny suggests a remarkable coincidence of (1) species divergence in the *Chlorella* clade and (2) species divergence in the *Stichococcus* clade. During the same period in the Triassic, the following splits resulting in new lineages might have appeared: split 1) *Stichococcus jenerensis* (rod-shapes; temperate-pantropic distribution) and *Diplosphaera* (rod-shapes/cell packages; temperate-tropical-arid-polar distribution); split 2) *Stichococcus bacillaris* (rod-shapes; temperate distribution) and *Desmococcus* (cell packages/filaments; temperate-polar distribution); split 3) *Stichococcus* clade3 (rod-shapes; temperate-tropical distribution) and *S.* clade4+clade5 (rod-shapes/cell packages; temperate-tropical-polar(?) distribution). In summary, *Stichococcus*- and *Diplosphaera/Desmococcus*-like morphologies evolved multiple times within the *Stichococcus* clade, possibly in successive order. The morphological plasticity of the *Stichococcus* lineages could explain its success in colonizing almost all terrestrial substrates (soil, epiphytic, epixylic,

epilithic, endolithic, lichens). In contrast to *Stichococcus*, the lineages of terrestrial Chlorellaceae are morphologically uniform, without any signs of multicellularity. The strikingly poor phylogenetic diversity of the Chlorellaceae found in humid tropical rainforests might be, apart from a simple sampling effect, due to low competitiveness. Instead, Chlorellaceae successfully colonized low-competitive microbial communities of extreme cold and hot terrestrial environments. Nonetheless, most regions on Earth have not been sampled yet. Although polar research undoubtedly contributes valuable biodiversity data, the picture of algal biogeography should be completed by much more samplings in rapidly disappearing biodiversity hotspots such as tropical rainforests.

Funding

The work was supported by the Priority Program 1374 "Infrastructure-Biodiversity-Exploratories" of the German Research Foundation (DFG) [grant number DFG-FR 905/16-1, FR 905/17-1] extended to T.F.; by the German Federal Ministry of Education and Research [grant number BMBF UKR 08/038] extended to T.F.; by the German Academic Exchange Service (DAAD; doctoral scholarship extended to L.H.) and by Ministry of Education, Youth and Sports of the Czech Republic [grant number: LM2010009, RVO67985939] extended to J.E.

Acknowledgements

We thank the managers of the three Exploratories, Kirsten Reichel-Jung, Swen Renner, Katrin Hartwich, Sonja Gockel, Kerstin Wiesner, and Martin Gorke for their work in maintaining the plot and project infrastructure; Christiane Fischer and Simone Pfeiffer for giving support through the central office, Michael Owonibi for managing the central data base, and Markus Fischer, Eduard Linsenmair, Dominik Hessenmöller, Jens Nieschulze, Daniel Prati, Ingo Schöning, François Buscot, Ernst-Detlef Schulze, Wolfgang W. Weisser and the late Elisabeth Kalko for their role in setting up the Biodiversity Exploratories project. Field work permits were issued by the responsible state environmental offices of Baden-Württemberg, Thüringen, and Brandenburg (according to §72 BbgNatSchG). We thank Alena Lukešová and Klára Řeháková for providing us with the cultures of polar *Chlorella*. We further thank Kathrin I. Mohr and Kristin C. Pahlmann for their sequencing efforts and Maike Lorenz for depositing new algal strains in the Culture Collection of Algae at the University of Göttingen (SAG).

Chapter 4

Molecular evidence for the wide distribution of two lineages of terrestrial green algae (Chlorophyta) from tropics to temperate zone

Ladislav Hodač¹, Christine Hallmann¹, Helen Rosenkranz², Fabian Faßhauer¹ and Thomas Friedl¹

¹University of Göttingen, Experimental Phycology and Culture Collection of Algae (SAG), Göttingen, Germany

²School of Biological Sciences, University of Bristol, Bristol, United Kingdom

(Published in *ISRN Ecology*; doi: 10.5402/2012/795924)

Abstract

Phylogenetic analyses of 18S rDNA sequences from environmental clones and culture strains revealed a widespread distribution of two subaerial green algal lineages, *Jenufa* and *Xylochloris*, recently described from rainforests in Southeast Asia. A new lineage of *Jenufa* (Chlorophyceae), most closely related to or even conspecific with *J. minuta*, was formed by sequences of European origin. Two more lineages of *Jenufa* were formed by three additional sequences from Ecuador and Panama. The other lineage was a close relative of *Xylochloris irregularis* (Trebouxiophyceae), probably representing a new species of the genus and distinct from the only so far described species, *X. irregularis*. It comprised two distinct clades each containing almost identical sequences from Germany and Ecuador. Analyses of the new sequences for both genera allowed to presume a preference of *J. minuta* to subaerial growth on rock or artificial hard substrates combined with a remarkable adaptation to extended periods of darkness, whereas *Xylochloris* may preferably occur on tree bark or in the soil.

Keywords: green algae; *Jenufa*; *Xylochloris*; molecular phylogeny; 18S rDNA; biogeography; substrate specificity

Abbreviations: BioNJ, Bio-neighbor-joining; CS, chlorophyte superclades including Chlamydomonadales and Sphaeropleales; MB, Bayesian inference; ML, Maximum likelihood; MP, Maximum parsimony; NMDS, Non-metric multidimensional scaling

Introduction

The neutral dispersal model (Fenchel and Finlay 2004) suggests that microorganisms do not exhibit biogeography, i.e. they are so small that no distributions barriers exist for them. This has been subject of debates by several authors, i.e. geographic distribution was found in ciliates (Foissner *et al.* 2008) and some microalgae (Řezáčová and Neustupa 2007). A recent study based on 18S rDNA sequence comparisons provided evidence for endemism in an Antarctic habitat, i.e. several new independent lineages of green algae were found in the benthos of an Antarctic lake (Lawley *et al.* 2004; De Wever *et al.* 2009). However, caution needs to be applied because there may still be an insufficient number of sequences available for many groups within the green algae. A lineage may appear “endemic” only as long as no other close relatives to it are known and this may easily change with the availability of new sequences from close relatives. In contrast to benthic freshwater algae, subaerial algae may be widely dispersed due to their remarkable adaptation to fast changing and adverse environmental conditions. Their resting cysts may easily survive transportation over long distances in the air being resistant to drought, high as well as low light intensities, and high UV radiation, e.g. due to the presence of thickened cell walls or light protection pigments (Řezanka *et al.* 2004; Karsten *et al.* 2005; Häubner *et al.* 2006; Gustavs *et al.* 2010).

Recently, subaerial microalgae from Southeast Asian tropical rainforest habitats gained increased interest by a series of studies (Eliáš *et al.* 2008; Neustupa *et al.* 2009; Eliáš *et al.* 2010; Němcová *et al.* 2011; Neustupa *et al.* 2011) which lead to the establishment of two new genera of green algae, *Jenufa* (Němcová *et al.* 2011) and *Xylochloris* (Neustupa *et al.* 2011). Both genera appeared as somehow distant to any known genera of green algae, despite a taxon sampling as large as possible has been applied in the 18S rDNA sequence analyses of both studies. The environmental sequencing approach extends the data pool available for diversity comparisons and may enable us to assess distribution patterns of protists (Šlapeta *et al.* 2005; Edgcomb *et al.* 2011). In the present study we took the advantage of an extended taxon sampling available through new 18S rDNA environmental sequences we obtained from our own ongoing studies which focus on changes in the algal diversity of certain terrestrial habitats along gradients of abiotic parameters in Europe (Germany, Ukraine) and South America (Ecuador). In addition, a variety of closely related environmental sequences have become available from GenBank. Here we demonstrate the presence of new lineages for both recently described genera, *Jenufa* (Chlorophyceae) and *Xylochloris* (Trebouxiophyceae), their wide distribution over long geographic distances and that both differ in their preferences towards certain substrates.

Materials and methods

Origin of analyzed sequences

The analyzed sequences and culture strains were provided by several our own on-going studies which focus on the algal diversities of various terrestrial habitats, see **Table 1**. They were derived from soil, tree bark or stone surfaces samples at four temperate habitats and one tropical habitat (**Table 1**). The majority of sequences were from environmental 18S rDNA clone libraries (334-1761 base pairs long), three from cultured strains which were accessioned by the SAG culture collection as strains SAG 2379, SAG 2382, and SAG 2383. Soil and tree bark samples originating from Germany were collected within the frame of the German Biodiversity Exploratories project (<http://www.biodiversity-exploratories.de>), for project description see Fischer *et al.* (2010) (**Fig. 1a**). Samples of biofilms dominated by green algae were investigated from the surface of sandstone samples within the framework of another study, focusing on wall sections of the castle “Burg Gleichen”, near Gotha (Thuringia, Germany) (Hallmann *et al.* 2011a). The biofilm samples were either from a sun-exposed wall area (**Fig. 1b**) or from the inner face of a scale from a dark basement vault (**Fig. 1c**). Samples from an even less light-exposed locality were from the concrete walls inside a World War II bunker monument on the North Sea offshore island Helgoland, Germany (**Fig. 1d-e**). Here the algal biofilms were exposed to extended periods of darkness because light is available from fluorescent tubes only during guided tours for tourists, i.e., 1.5 hours per day and around 8 months per years. The soil sample from Ukraine was from the arid steppe zone (Chernomorskiy Biosphere Reserve). The soil and tree bark samples from South America originated from the tropical mountain rain forest in Southern Ecuador along an elevation transect at 1000 m, 2000 m and 3000 m above sea level (a.s.l.) corresponding to three different regions (**Table 1; Fig. 1f-g**).

Microscopy and rDNA sequence determination

Microscopic observations of cultures were conducted using an Olympus BX60 microscope (Tokyo, Japan) with Nomarski DIC optics and an attached ColorView III camera (Soft Imaging System, Münster, Germany). Micrographs were processed using Cell[^]D image software (Soft Imaging System, Münster, Germany). The detailed experimental procedures on how the 18S rDNA clone libraries and cultures were established will be described elsewhere. Briefly, DNA was extracted from environmental samples using MoBio PowerSoil DNA isolation Kit (MoBio Laboratories Inc. Carlsbad, CA, USA) and from mechanically disrupted cultured algal cells using Invisorb® Spin Plant Mini Kit (Stratec, Berlin, Germany). Almost full length 18S rDNA

sequences were amplified from the cultures and European environmental samples with primers preferentially binding to green algal rDNAs, for the samples from Ecuador a region from about 350 base pairs upstream of the 3' end of 18S rDNA downstream to the 26S rDNA which included the ITS2 rDNA for DNA barcoding purposes was amplified. Either almost full 18S rDNA sequences were obtained or partial sequences comprising either the hypervariable regions V2, V3 and V4 (European environmental samples) or hypervariable regions V8 and V9 regions (Ecuador environmental samples) of 18S rDNA (Neefs and De Wachter 1990; Lee and Gutell 2012). The newly determined sequences are available from GenBank under the accession numbers JQ988923-JQ988943.

Table 1. List of new sequences obtained for this study.

| Taxon | Clone identifier (*cultured strains) | Accession number | Length (base pairs) | Region of origin | GPS-coordinates | Habitat | Substrate |
|--------------------|---|---------------------|------------------------|---------------------------|-------------------|--------------------------|---|
| <i>Xylochloris</i> | AEQ1B-K1547 | JQ988937 | 748 | Germany, Swabian Alb | 48°23'N, 9°20'E | Grassland | Soil |
| | AEW2R-K255 | JQ988938 | 1734 | Germany, Swabian Alb | 48°37'N, 9°35'E | Spruce forest | Tree bark, <i>Picea abies</i> |
| | AEW2R-K265 | JQ988934 | 842 | Germany, Swabian Alb | 48°37'N, 9°35'E | Spruce forest | Tree bark, <i>Picea abies</i> |
| | BOAEW3R1-K03 | JQ988939 | 716 | Germany, Swabian Alb | 48°41'N, 9°35'E | Spruce forest | Tree bark, <i>Picea abies</i> |
| | SAG 2382* | JQ988942 | 1719 | Germany, Swabian Alb | 48°39'N, 9°26'E | Beech forest | Soil |
| | HEG9B-K2617 | JQ988940 | 1702 | Germany, Hainich | 51°13'N, 10°22'E | Grassland | Soil |
| | HEW10SB-K5621 | JQ988935 | 854 | Germany, Hainich | 51°05'N, 10°27'E | Beech forest | Soil |
| | HEW4SB-K5577 | JQ988936 | 854 | Germany, Hainich | 51°24'N, 10°32'E | Beech forest | Soil |
| | HEW4SB-K5606 | JQ988941 | 733 | Germany, Hainich | 51°24'N, 10°32'E | Beech forest | Soil |
| | U33B-K2708 | JQ988943 | 628 | Ukraine, Ascania Nova | 46°27'N, 33°54'E | Steppe | Soil |
| | C32U6-13 | JQ988925 | 334 | Ecuador, Cajanuma | 04°06'S, 079°10'W | Rainforest 3000 m a.s.l. | Soil |
| | S30L6-32 | JQ988926 | 334 | Ecuador, San Francisco | 03°58'S, 079°04'W | Rainforest 2000 m a.s.l. | Tree bark |
| <i>Jenufa</i> | 3GA1K3658 | JQ988931 | 851 | Germany, Drei Gleichen | 50°52'N, 10°50'E | Castle Gleichen | Sandstone, surface |
| | GS1K32 | JQ988932 | 1761 | Germany, Drei Gleichen | 50°52'N, 10°50'E | Castle Gleichen | Sandstone, scale (endolithic) Concrete |
| | KB2B11-06 | JQ988928 | 1596 | Germany, Helgoland | 54°11'N, 7°53'E | Bunker | Concrete |
| | KB4B5-06 | JQ988929 | 1570 | Germany, Helgoland | 54°11'N, 7°53'E | Bunker | Concrete |
| | SAG 2379* | JQ988927 | 1563 | Germany, Helgoland | 54°11'N, 7°53'E | Bunker | Concrete |
| | SAG 2383* | JQ988933 | 1700 | Germany, Swabian Alb | 48°38'N, 9°38'E | Beech forest | Soil |
| | FLUL-B69M5-34 | JQ988923 | 335 | Ecuador, Bombuscaro | 04°07'S, 078°58'W | Rainforest 1000 m a.s.l. | Tree bark |
| | S46L1-7 | JQ988924 | 338 | Ecuador, San Francisco | 03°58'S, 079°04'W | Rainforest 2000 m a.s.l. | Soil |




Figure 1. Sampling sites. (a) Young spruce forest in the Swabian Alb, (b-c) wall sections of castle Gleichen (Germany), (b) sun-exposed wall area, (c) inner face of a scale from a dark basement vault (the white arrow shows the sampled green biofilm), (d-e) biofilm samples from concrete walls in a World War II bunker on the North Sea offshore island Helgoland, (e) sampled green biofilm in detail, (f-g) tropical sub-mountain rainforest in Ecuador (the study site in Bombuscaro, 1000 m a.s.l.), and (g) green biofilms on tree bark in detail.

Phylogenetic analyses

The newly determined sequences were used in database queries using BLAST (Altschul *et al.* 1997) at NCBI (<http://www.ncbi.nlm.nih.gov/>) to retrieve their closest neighboring sequences available from public sequence data bases (state in March 2012). In addition, the newly determined and their retrieved closest relative sequences were compared to a broad selection of green algal sequences which was available in the internal 18S rDNA sequence database of our laboratory and is maintained in the ARB program (version 05.05.26, 2004, www.arb-home.de) (Ludwig *et al.* 2004). This database was updated with all currently available 18S rDNA sequences from green algae. The comparisons identified sequences of both genera, *Jenufa* and *Xylochloris*, as well as their next relatives. Potential chimeras were identified by Bellerophon (Huber *et al.* 2004) and excluded from the analyses. The almost full 18S rDNA sequences of *Jenufa*, *Xylochloris* and their close relatives were aligned in two separated data sets, i.e. together with representatives of all known lineages from the two green algal classes, Chlorophyceae and Trebouxiophyceae. First the sequences were aligned with MAFFT ver. 6 (Katoh and Toh 2008) online (<http://mafft.cbrc.jp/alignment/server/index.html>) in order to identify and exclude intron positions. Then the sequences from each class were aligned with MUSCLE online (<http://www.ebi.ac.uk/Tools/msa/muscle>). The alignments were carefully checked for possible misaligned positions by eye in BioEdit (Hall 1999). The final alignment of Chlorophyceae comprised 97 sequences and 1790 positions (742 variable/535 parsimony informative sites), the alignment of Trebouxiophyceae included 99 sequences and 1807 positions (710 variable/497 parsimony informative sites). Based on the AIC criterion in jModelTest 0.1.1 (Posada 2008), the GTR+ Γ +I nucleotide substitution model was selected as to fit best both alignments. ML phylogenies were obtained from RAxML 7.0.4 (Stamatakis *et al.* 2008). Confidence values for the obtained groups (edge support) were inferred from rapid bootstrapping algorithm (1000 replicates) as implemented in RAxML. Bayesian posterior probabilities for internal nodes were obtained using MrBayes 3.2 (Ronquist *et al.* 2012). Two MCMC runs for four million generations each with one cold and three heated chains using the GTR+ Γ +I+COV evolutionary model (parameters were estimated from the data) were performed with trees sampled every 100 generations. For comparisons of European with Ecuadorian environmental clones for which different regions of the 18S rDNA were determined (see above), only the hypervariable regions V8/V9 of the 18S rDNA could be used because they represent an overlapping sequence region available in both data sets (European data set comprised the full sequences whereas the Ecuadorian one the partial ones). Then only sequences of closest relatives were used for the

phylogenetic analyses, i.e. the investigated datasets comprised less than 10 sequences and were only about 300 base pairs long. Bio-Neighbor-Joining distance phylogenies based on Jukes-Cantor genetic distances and 1000 bootstrap replicates were computed in the program SeaView ver. 4 (Gouy *et al.* 2010). The parsimony analysis was conducted using DNAPARS as implemented in SeaView with 10 times randomization of the sequence order and 1000 bootstrap replicates. Evolutionary distances among selected sequences were computed as p-distances in MEGA5 (Tamura *et al.* 2011) with ambiguous alignment positions removed for each sequence pair.

Multivariate statistical analysis

In order to compare the clones originating from the different substrates within Europe (tree bark, soil, stone surfaces) Non-metric Multidimensional Scaling (NMDS) analysis based on genetic distances was computed. The analysis was conducted on the matrix of Jukes-Cantor genetic distances (within alignments comprising hypervariable regions V2, V3 and V4 of the 18S rDNA) as implemented in the program package PAST ver. 2.12 (Hammer *et al.* 2001). The reliability of the resulting ordination was examined by checking the stress values from Shepard-plot. The alignment comprising the new sequences of *Jenufa minuta* relatives included 8 sequences and 658 positions (**Fig. 2d**), whereas the other comprising the new sequences of *Xylochloris*-relatives included 10 sequences and 595 positions (**Fig. 3d**).

Results and Discussion

Jenufa lineage

The 18S rDNA phylogeny revealed the position of several of the obtained new environmental sequences and the three cultured isolates into the immediate vicinity of the genera *Jenufa* (Chlorophyceae, **Fig. 2a, 2b, 2d; Fig. 1 in Appendix**) and *Xylochloris* (Trebouxiophyceae, **Fig. 3a, 3b, 3d; Fig. 2 in Appendix**). We recovered sequences assigned to *Jenufa* from environmental clone libraries from Germany (soil, hard substrates), Ukraine (soil) and Ecuador (soil, tree bark) (**Table 1**). The full European environmental sequences and the culture strains formed a well-supported monophyletic cluster (p-distances 0.001-0.004) closely related to *Jenufa minuta*. The p-distances among European sequences and *J. minuta* (0.016 – 0.019) were about two times lower than between *J. minuta* and *J. perforata* (0.038). The partial environmental sequences from Ecuador constitute a lineage by its own positioned between *J. minuta* and *J. perforata*, but without significant bootstrap support (**Fig. 2d**). As already discussed previously in Němcová *et al.* (2011),

Jenufa may represent a distinct probably new lineage of Chlorophyceae of still unresolved relationship to the orders Sphaeropleales and Chlamydomonadales. Within the 18S rDNA phylogeny presented here, *Jenufa* is placed together with *Golenkinia* on the very basis of Chlamydomonadales, although without statistical support (ML/MB: 69/0.83 in **Fig. 2a**; **Fig. 1 in supplement**). In contrast, in Bayesian tree topology, *Jenufa*- and *Golenkinia* clades formed a lineage independent of the Chlamydomonadales and Sphaeropleales. The light-microscopic observations on the two available culture strains SAG 2383 and SAG 2379 were in agreement with their assignment to *Jenufa* inferred from the 18S rDNA phylogenetic analyses. Both strains exhibited morphological characters as previously described for *J. minuta* and *J. perforata* (**Fig. 4a**). Both species were found indistinguishable from each other without the use of a confocal microscope and the analysis of cell wall structures (Němcová *et al.* 2011).




Figure 2. (a) Schematic 18S-based phylogenetic tree of the Chlorophyceae showing the position of sequences assigned to *Jenufa minuta*, (b) *Jenufa*/*Golenkinia* clades in detail shown as a section of the full 18S rDNA ML-phylogenetic tree (**Fig. S1 in Appendix**), (c) world map with dots representing regions where *Jenufa* sequences were detected, and (d) BioNJ-tree based on partial 18S sequences (hypervariable regions V8 and V9). Colors in figures (c) and (d) indicate three geographical regions where *Jenufa* occurs (orange—southeast Asia, green—south and Latin America, blue—Europe). Names in bold represent specimens analyzed as both partial- and full-length 18S rDNA sequences.

Xylochloris lineage

Other new environmental sequences and the sequence of strain SAG 2382 were closely neighboring to *Xylochloris irregularis* (Trebouxiophyceae) in the 18S rDNA phylogenetic analyses, but without significant bootstrap support (**Fig. 3b**). The p-distances among the full European sequences and *X. irregularis* (0.031 – 0.035) was slightly smaller than those between

the two previously described species of *Jenufa* (Chlorophyceae) which may imply that the European sequences and SAG 2382 represent a species independent of *X. irregularis*. The *Xylochloris* clade comprised environmental clones from soils and tree bark from Germany, Singapore and Ecuador (**Fig. 3d**). The addition of new sequences supported a sister-group relationship of *Xylochloris* with a clade comprising of environmental clones from cold-fumarole soils of the Andes (Costello *et al.* 2009) and two culture strains from terrestrial habitats in Germany assigned to *Dictyochloropsis* (**Fig. 3b, Fig. 2 in Appendix**). The latter clade was well supported in bootstrap tests, whereas its sister-group relationship with *Xylochloris* was only supported in MB analyses. Strain SAG 2382 shared several morphological characters with *X. irregularis* (Neustupa *et al.* 2011), i.e. the presence of a pyrenoid, the lobed chloroplast in vegetative cells, similar cell size, and the irregular cell shape. However, the strain also exhibited distinguishing characters of its own, namely the presence cylindrical as well as spherical autospores at the same time (**Fig. 4b-d**).




Figure 3. (a) Schematic 18S-based phylogenetic tree of the Trebouxiophyceae showing the placement of sequences assigned to *Xylochloris*, (b) the clade including *Xylochloris irregularis* and its closest relatives from Europe shown as a section of the full 18S rDNA ML-phylogenetic tree of the Trebouxiophyceae (**Fig. S2**), (c) world map with dots representing regions where *Xylochloris* sequences were detected, and (d) BioNJ-tree based on partial 18S sequences (hypervariable regions V8 and V9). Colors in Figures (c) and (d) indicate three geographical regions where *Xylochloris* occurs (orange—southeast Asia, green—South and Latin America, blue—Europe). Names in bold represent specimens analyzed as both partial- and full-length 18S rDNA sequences.




Figure 4. Microphotographs of the strains SAG 2383 (a) and SAG 2382 (b–d), (a) vegetative cells and autospores of the European *Jenufa* strain SAG 2383 isolated from a soil sample from Swabian Alb (Germany), (b) vegetative cell of *Xylochloris* strain SAG 2382 exhibiting the presence of one pyrenoid, (c) irregular cylindrical to oval autospores typical for the strain, and (d) vegetative cells with lobed margins of the chloroplast. Scale bar represents 20 µm.

Substrate preferences of *Jenufa* and *Xylochloris*

Jenufa (Chlorophyceae) and *Xylochloris* (Trebouxiophyceae) represent distinct lineages of terrestrial green algae without any known members from freshwater habitats. Despite *Jenufa* and *Xylochloris* were detected in both habitats, soils and tree barks, in the tropics, the sequence analysis of their closest relatives from Europe suggests a pattern reflecting different substrate preferences for both genera. In our study *Jenufa* was recovered from hard substrates, except for one isolate (SAG 2383) established from soil. The hard substrates were epilithic biofilms (surface of bunker walls on Helgoland, outer surface of walls of castle Gleichen, walls of Altamira cave in Spain) as well as endolithic biofilms (rocks in Piora Valley/1,965 m a.s.l./Swiss Alps; Horath and Bachofen 2009), or and the upper side of wall scales on castle Gleichen) Although the NMDS analysis (**Fig. 5**) of European partial sequences did not point out any clear groupings due to substrate type (endolithic *versus* epilithic/soil), the second ordination axis indicated variation among sequences due to different light intensities at the sampling sites. This may be seen among *Jenufa* sequences from the Helgoland bunker monument. Two sequences, (KB2B11-06 and SAG 2379, originating from two sites of low irradiance within the bunker were genetically closer than both to another *Jenufa* sequence, KB4B5-06, from a site of three times higher irradiance (HR unpublished results). In addition, the genetic differences between two *Jenufa* sequences, 3GA1K3658 from epilithic and GS1K32 from endolithic habitats of the Gleichen castle sandstone walls, may be explained by adaptation to different light intensities as well, as pointed out by the NMDS-plot (**Fig. 5**). The newly determined *Xylochloris* sequences were all recovered from tree bark as well as from soils. The NMDS-analysis pointed out a separation of two groups of very closely related sequences from tree bark against from those from soils (**Fig. 6**). These two groups may represent even two separate taxa at the level of species or below, as already discussed

above. Members of both groups were recovered also from Ecuador, hence both may be widespread as well.




Figure 5. Nonmetric multidimensional Scaling (NMDS) diagram. The ordination analysis based on the Jukes-Cantor genetic distances among six environmental clones and two isolates representing sequences of *Jenufa minuta* detected in Europe on various hard substrates. For the analysis, 18S partial sequences comprising hypervariable regions V2, V3, and V4 were used. The red-colored points represent sequences from the walls of castle Gleichen/Germany, the blue ones are from the bunker on Helgoland, an offshore island in the North Sea. For the corresponding accession numbers of the sequences see Table 1.




Figure 6. Nonmetric multidimensional scaling (NMDS) diagram. The ordination based on Jukes-Cantor genetic distances among nine environmental clones and one isolate representing a yet undescribed European species of the genus *Xylochloris*. For the analysis 18S partial sequences comprising hypervariable regions V2, V3, and V4 were used. The colors encode for three different habitats: green—tree bark, blue—forest soil, brown—grassland soil. For the corresponding accession numbers of the sequences see Table 1.

Identification of environmental rDNA clones using new cultured isolates

A high 18S rDNA sequence similarity between strains of closely related terrestrial green algae from tropics and temperate habitats has already been presented recently (Němcová *et al.* 2011). Two cultured isolates of *J. minuta* from the tropics were found closely related to hitherto unidentified environmental clones from the Swiss Alps. This example shows that unidentified environmental clones may become identified as soon as cultures of close relatives become available, at least at the generic level if a morphology-based species or generic concept is applied. In our study we revealed two new isolates, SAG 2379 and SAG 2383, whose sequences were almost identical (p-distances 0.001-0.004) to the environmental clones from the Swiss Alps. Another case where environmental clones became identified due to the availability of culture strains was found in the “*Dictyochloropsis*” clade, sister with *Xylochloris* in the Trebouxiophyceae (**Fig. 3d**). Due to the very low genetic distances (p-distances 0.001 – 0.005) present within the clade, the strains of *Dictyochloropsis* from tree barks in Germany and the environmental clones from the high Andes Mountains (Chile) may be regarded as member of a single species which again exhibits a wide geographic distribution. The genetic distances between the Andes clones/*Dictyochloropsis* pairs were even shorter (0.003-0.004) than between both Andes environmental clones (0.005). In conclusion, for a better understanding of the biogeography of terrestrial microalgae it seems to be of crucial importance not only to sequence environmental clones from somehow exotic or extreme habitats, but also to establish and sequence cultures from seemingly well investigated regions like Central Europe. For the studies on biogeography of terrestrial algae we are still faced with the lack of sufficient molecular data to compare new findings to, because the most diversity evidences from either Central Europe or many other geographical regions relies on morphological observations only (Hoffmann *et al.* 2007; Neustupa and Škaloud 2008; Büdel *et al.* 2009; Khaybullina *et al.* 2010). We expect the situation to change in the future through extended application of the next generation sequencing (e.g. 454-pyrosequencing) concentrated on hypervariable regions of the 18S rDNA (Amaral-Zettler *et al.* 2009).

Acknowledgements

This study would not have been possible without the stipend of the DAAD extended to L.H. This work was supported by two grants from the German Research Foundation (DFG) extended to T.F. (contract no. FR 905/16-1 and FR 905/17-1) and a grant from the German Federal Ministry of Education and Research (BMBF) extended to T.F. (contract no. UKR 08/038). Parts of the

work were supported by the USA National Science Foundation (DEB-0529737) to Louise A. Lewis, University of Connecticut, in collaboration with T.F. T.F. and H.R. thank Kathrin I. Mohr for initiating and supervising the study of algal biofilms of the Helgoland bunker, T.F. thanks Igor Y. Kostikov for provision of the opportunity to study samples from the steppe zone of Ukraine within the project BMBF UKR 08/038. C.H. acknowledges the financial support provided by the Deutsche Bundesstiftung Umwelt (DBU) extended to M. Hoppert, Göttingen. We acknowledge provision of environmental sequences determined by a student summer course in 2011 (project FoLL of Göttingen University). We also thank two anonymous reviewers for their critical comments on the manuscript.

General Summary

Green microalgae (Chlorophyta) dominate soils in the northern temperate climate zone, however, their biodiversity is still poorly understood. Available diversity inventories rely on light microscopy and morphospecies, which are hardly distinguishable without molecular markers. Whereas particular attention was paid to molecular diversity of terrestrial microalgae in extreme regions, temperate climate zones remain almost unexplored. We aim to uncover the phylogenetic diversity of green microalgae isolated from Central European soils and from periodically desiccating freshwater creek biofilms. Such terrestrial and semi-aquatic habitats are inhabited by green microalgae, which are presumably able of long-distance dispersal. Their cosmopolitan distribution is presumed, however, supporting molecular evidence is almost missing. By assembling newly obtained green algal sequences together with accessions from remote geographic regions, we further aim to address a question of biogeography of terrestrial microalgae.

Soil samples were taken from grassland and forest plots within the German Biodiversity Exploratories. Freshwater biofilms dominated by green microalgae were sampled in two karstwater creeks in Germany. In total, 280 new monoclonal cultures of green microalgae were examined by molecular phylogenetic methods and by light microscopy. By using ribosomal 18S and ITS2 sequences, we recognized about 100 monophyletic species of green microalgae. The newly obtained sequences were blasted against public databases in order to infer taxonomy, distribution and ecology of the detected species. Further monoclonal cultures and environmental clones originated from additional samplings in Germany, Ecuador, the Arctic and Antarctic.

Most Chlorophyta isolated from German soils were highly similar ($\geq 99.5\%$ threshold) to cultured relatives already known from Europe, predominantly from soils and further terrestrial substrates such as tree barks and rocks. Considering a lower similarity threshold ($\geq 99\%$), about 90% of our cultures matched environmental clones inferred from various terrestrial and aquatic habitats. Similarly, Chlorophyta detected in creek biofilms partly represent terrestrial species, but some are known also from planktonic communities. In soils, we detected novel species mostly within established lineages. Only one soil isolate, provisionally named *Navichloris fusiformis*, was recognized as a member of a novel monophyletic lineage, comprising accessions from South and North American deserts. In multiple cases, molecular data supported close relatedness between the European soil isolates and species from tropics and polar regions. We show, that well supported monophyletic clades of *Stichococcus* exhibit either temperate-polar *or* temperate-

tropical distributions. In contrast, some monophyletic clades of *Chlorella*-like microalgae were so far evidenced only from polar regions and hot deserts. The long-distance dispersal was finally confirmed for particular species of *Stichococcus*, *Chlorella* and *Klebsormidium*.

Forest and grassland soils in Central Europe host species of green microalgae otherwise known from a broad spectrum of terrestrial as well as aquatic habitats. New species and even novel lineages can still be uncovered by using standard culturing techniques. Despite the striking similarity between some European and exotic species, unambiguous molecular evidence of the intraspecific long-distance dispersal is still scant. At the present state of knowledge, our data suggest the existence of biogeography of airborne microalgae.

References

- Achtman M, Wagner M. 2008. Microbial diversity and the genetic nature of microbial species. *Nature Reviews Microbiology* 6(6):431-440.
- Adl SM, Simpson AG, Farmer MA, Andersen RA, Anderson OR, Barta JR, Bowser SS, Brugerolle G, Fensome RA, Fredericq S, James TY, Karpov S, Kugrens P, Krug J, Lane CE, Lewis LA, Lodge J, Lynn DH, Mann DG, McCourt RM, Mendoza L, Moestrup O, Mozley-Standridge SE, Nerad TA, Shearer CA, Smirnov AV, Spiegel FW, Taylor MF. 2005. The new higher level classification of eukaryotes with emphasis on the taxonomy of protists. *Journal of Eukaryotic Microbiology* 52(5):399-451.
- Aguilera A, Zettler E, Gomez F, Amaral-Zettler L, Rodriguez N, Amils R. 2007. Distribution and seasonal variability in the benthic eukaryotic community of Rio Tinto (SW, Spain), an acidic, high metal extreme environment. *Systematic and Applied Microbiology* 30(7):531-546.
- Aguilera Á, Souza-Egipsy V, González-Toril E, Rendueles O, Amils R. 2010. Eukaryotic microbial diversity of phototrophic microbial mats in two Icelandic geothermal hot springs. *International Microbiology* 13(1):21-32.
- Al-Fredan M, Fathi A. 2007. Preliminary survey of edaphic algae in Al-Hasa region, Saudi Arabia. *Pakistan Journal of Biological Sciences* 10(18):3210-3214.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25(17):3389-3402.
- Amaral-Zettler LA, McCliment EA, Ducklow HW, Huse SM. 2009. A method for studying protistan diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA genes. *PLoS ONE* 4(7):e6372.
- Andreyeva V. 1973. Novye vidy *Chlorella* Beijer. *Botanicheskij Zhurnal* 58:1735-1741.
- Andreyeva VM. 1998. Pochvennye i aerofilnye zelenye vodorosli: Chlorophyta (Tetrasporales, Chlorococcales, Chlorosarcinales). Nauka. 315 pp.
- Androsova EJ. 1964. O sostave vodoroslej pochv g. Novosibirska i ego okrestnostej. Aktualnye problemy sovremennoj algologii. Tscherkassy. p. 23-25.
- Archibald PA, Bold HC. 1970. Phycological Studies XI. The Genus *Chlorococcum* Meneghini. University of Texas Publication No. 7015. 115 pp.
- Archibald PA. 1972. A preliminary survey of the edaphic algae of Costa Rica and San Andreas Isle. *Soil Science* 113:63-64.
- Archibald PA, Bold HC. 1975. Notes on the edaphic algae of the Galapagos. *Soil Science* 120:400-402.
- Arp G, Bissett A, Brinkmann N, Cousin S, Reimer A, Shiraishi F, Stackebrandt E, Zippel B. 2010. Tufa-forming biofilms of German karstwater streams: microorganisms, exopolymers, hydrochemistry and calcification. Geological Society, London, Special Publications. p. 83-118.
- Arvik JH, Willson DL. 1974. Soil algae of Eniwetok Atoll, The Marshall Islands. *Pacific Science* 28(2):189-190.
- Asfandijarova LZ. 2008. Analiz pochvennoj algoflory posevov mnogoletnikh trav Preduralja respubliki Bashkortostan. Fundamentalnye i prikladnye problemy botaniki v nachale XXI veka. Algologija. p. 10-12.

- Ashley J, Rushforth SR, Johansen JR. 1985. Soil algae of cryptogamic crusts from the Uintah Basin, Utah, USA. *Western North American Naturalist* 45(3):432-442.
- Bachura JM, Khranchenkova OM. 2008. Osobennosti vosstanovleniya algogruppirovok pochv na kostrishchakh. *Fundamentalnye i prikladnye problemy botaniki v nachale XXI veka. Algologiya*. p. 13-16.
- Bailly J, Fraissinet-Tachet L, Verner M-C, Debaud J-C, Lemaire M, Wésolowski-Louvel M, Marmeisse R. 2007. Soil eukaryotic functional diversity, a metatranscriptomic approach. *The ISME Journal* 1(7):632-642.
- Baker BJ, Tyson GW, Goosherst L, Banfield JF. 2009. Insights into the diversity of eukaryotes in acid mine drainage biofilm communities. *Applied and Environmental Microbiology* 75(7):2192-2199.
- Balezina L. 1974. Effect of concentrated mineral fertilizers on the growth of soil algae. *Soviet Soil Science*: 550-552.
- Barreiro DL, Zamalloa C, Boon N, Vyverman W, Ronsse F, Brilman W, Prins W. 2013. Influence of strain-specific parameters on hydrothermal liquefaction of microalgae. *Bioresource Technology* 146:463-471.
- Barsanti L, Frassanito AM, Passarelli V, Evangelista V, Etebari M, Paccagnini E, Lupetti P, Lenzi P, Verni F, Gualtieri P. 2013. *Tetraflagellochloris mauritanica* gen. et sp. nov. (Chlorophyceae), a new flagellated alga from the Mauritanian Desert: morphology, ultrastructure, and phylogenetic framing. *Journal of Phycology* 49(1):178-193.
- Bashan Y, Lopez BR, Huss VA, Amavizca E, de-Bashan LE. 2015. *Chlorella sorokiniana* (formerly *C. vulgaris*) UTEX 2714, a non-thermotolerant microalga useful for biotechnological applications and as a reference strain. *Journal of Applied Phycology*:doi:10.1007/s10811-015-0571-z.
- Bates ST, Clemente JC, Flores GE, Walters WA, Parfrey LW, Knight R, Fierer N. 2013. Global biogeography of highly diverse protistan communities in soil. *The ISME Journal* 7(3):652-659.
- Bazin P, Jouenne F, Friedl T, Deton-Cabanillas A-F, Le Roy B, Véron B. 2014. Phytoplankton diversity and community composition along the estuarine gradient of a temperate macrotidal ecosystem: combined morphological and molecular approaches. *PLoS ONE* 9(4):e94110.
- Becker B, Marin B. 2009. Streptophyte algae and the origin of embryophytes. *Annals of Botany* 103(7):999-1004.
- Becker B. 2013. Snow ball earth and the split of Streptophyta and Chlorophyta. *Trends in Plant Science* 18(4):180-183.
- Behnke A, Bunge J, Barger K, Breiner H-W, Alla V, Stoeck T. 2006. Microeukaryote community patterns along an O₂/H₂S gradient in a supersulfidic anoxic fjord (Framvaren, Norway). *Applied and Environmental Microbiology* 72(5):3626-3636.
- Beijerinck M. 1893. Bericht über meine Kulturen niederer Algen auf Nährgelatine. *Zentralblatt Für Bakteriologie* 13:781-786.
- Benson D, Karsch-Mizrachi I, Clark K, Lipman D, Ostell J, Sayers E. 2012. GenBank. *Nucleic Acids Research* 40:D48-D53.
- Bérard A, Rimet F, Capowiez Y, Le Boulanger C. 2004. Procedures for determining the pesticide sensitivity of indigenous soil algae - a possible bioindicator of soil contamination? *Archives of Environmental Contamination and Toxicology* 46:24-31.

- Bérard A, Dorigo U, Humbert JF, Martin-Laurent F. 2005. Microalgae community structure analysis based on 18S rDNA amplification from DNA extracted directly from soil as a potential soil bioindicator. *Agronomie* 25:1-7.
- Bhatnagar A, Makandar MB, Garg MK, Bhatnagar M. 2008. Community structure and diversity of cyanobacteria and green algae in the soils of Thar Desert (India). *Journal of Arid Environments* 72(2):73-83.
- Bilan M, Usov A. 2001. Polysaccharides of calcareous algae and their effect on the calcification process. *Russian Journal of Bioorganic Chemistry* 27(1):2-16.
- Bischoff HW, Bold HC. 1963. *Phycological Studies IV. Some soil algae from enchanted rock and related algal species.* The University of Texas Press No. 6318. 95 pp.
- Bock C, Pröschold T, Krienitz L. 2010. Two new *Dictyosphaerium*-morphotype lineages of the Chlorellaceae (Trebouxiophyceae): *Heynigia* gen. nov. and *Hindakia* gen. nov. *European Journal of Phycology* 45(3):267-277.
- Bock C, Krienitz L, Pröschold T. 2011. Taxonomic reassessment of the genus *Chlorella* (Trebouxiophyceae) using molecular signatures (barcodes), including description of seven new species. *Fottea* 11(2):293-312.
- Boenigk J, Pfandl K, Stadler P, Chatzinotas A. 2005. High diversity of the 'Spumella-like' flagellates: an investigation based on the SSU rRNA gene sequences of isolates from habitats located in six different geographic regions. *Environmental Microbiology* 7(5):685-697.
- Borzi A. 1895. *Studi algologici: saggio di ricerche sulla biologia delle alghe.* Fasc. 1-2. G. Capara. 476 pp.
- Brandt F, Stockmayer S. 1925. Analyse der aerophilen Grünalgenanflüge, insbesondere der proto-pleurococcoiden Formen. *Archiv für Protistenkunde* 52:265-355.
- Bråte J, Klaveness D, Rygh T, Jakobsen KS, Shalchian-Tabrizi K. 2010. Telonemia-specific environmental 18S rDNA PCR reveals unknown diversity and multiple marine-freshwater colonizations. *BMC Microbiology* 10(1):168.
- Brinkmann N, Hodač L, Mohr KI, Hodačová A, Jahn R, Ramm J, Hallmann C, Arp G, Friedl T. 2015. Diversity of Cyanobacteria and Diatom communities along pCO₂ and calcite saturation gradients in calcifying biofilms covering karst-water stromatolites in Germany. *Geomicrobiology Journal* 32(3-4):255-274.
- Bristol BM. 1920. On the alga-flora of some desiccated English soils: an important factor in soil biology. *Annals of Botany* 34(1):35-80.
- Bristol BM. 1927. On the algae of some normal English soils. *The Journal of Agricultural Science* 17(4):563-588.
- Broady P. 1983. New records of Chlorophycean micro-algae cultured from antarctic terrestrial habitats. *Nova Hedwigia* 36(2-4):445-84.
- Broady PA. 1978. The terrestrial algae of Glerardalur Akureyri, Iceland. *Acta Botanica Islandica* 5:3-60.
- Broady PA. 1979a. A preliminary survey of the terrestrial algae of the Antarctic Peninsula and South Georgia. *British Antarctic Survey Bulletin* 48:47-70.
- Broady PA. 1979b. Qualitative and quantitative observations on green and yellow-green algae in some English soils. *British Phycological Journal* 14:151-160.
- Broady PA. 1984. Taxonomic and ecological investigations of algae on steam-warmed soil on Mt. Erebus, Ross Island, Antarctica. *Phycologia* 23(3):257-271.
- Broady PA. 1989. Survey of algae and other terrestrial biota at Edward VII Penninsula, Mary Bird Land. *Antarctic Science* 1:215-224.

- Broady PA. 1996. Diversity, distribution and dispersal of Antarctic terrestrial algae. *Biodiversity and Conservation* 5:1307-1335.
- Brown MV, Philip GK, Bunge JA, Smith MC, Bissett A, Lauro FM, Fuhrman JA, Donachie SP. 2009. Microbial community structure in the North Pacific ocean. *The ISME Journal* 3(12):1374-1386.
- Buchheim MA, Keller A, Koetschan C, Förster F, Merget B, Wolf M. 2011. Internal transcribed spacer 2 (nu ITS2 rRNA) sequence-structure phylogenetics: towards an automated reconstruction of the green algal tree of life. *PLoS ONE* 6(2):e16931.
- Büdel B. 2003. Biological soil crusts of South America. In: Belnap J, Lange OL, editors. *Biological soil crusts: Structure, function, and management*. Springer. p. 51-55.
- Büdel B, Darienko T, Deutschewitz K, Dojani S, Friedl T, Mohr KI, Salisch M, Reisser W, Weber B. 2009. Southern African biological soil crusts are ubiquitous and highly diverse in drylands, being restricted by rainfall frequency. *Microbial Ecology* 57(2):229-247.
- Buzer J, Dohmeier R, Du Toit D. 1985. The survival of algae in dry soils exposed to high temperatures for extended time periods. *Phycologia* 24(2):249-251.
- Cadoret J-P, Garnier M, Saint-Jean B. 2012. Microalgae, functional genomics and biotechnology. *Advances in Botanical Research* 64:285-341.
- Caisová L, Marin B, Melkonian M. 2013. A consensus secondary structure of ITS2 in the Chlorophyta Identified by phylogenetic reconstruction. *Protist* 164:482-496.
- Cantino PD, de Queiroz K. International Code of Phylogenetic Nomenclature, Version 4b. <https://www.ohio.edu/phylocode/index.html>.
- Caron DA, Countway PD, Savai P, Gast RJ, Schnetzer A, Moorthi SD, Dennett MR, Moran DM, Jones AC. 2009. Defining DNA-based operational taxonomic units for microbial-eukaryote ecology. *Applied and Environmental Microbiology* 75(18):5797-5808.
- Carson JL, Brown RM. 1976. The correlation of soil algae, airborne algae, and fern spores with meteorological conditions on the Island of Hawaii. *Pacific Science* 30(2):197-205.
- Carson JL, Brown RM. 1978. Studies on Hawaiian freshwater and soil algae II. Algal colonization and succession on a dated volcanic substrate. *Journal of Phycology* 14(2):171-178.
- Cary SC, Fierer N. 2014. The importance of sample archiving in microbial ecology. *Nature Reviews Microbiology* 12(12):789-790.
- Cavacini P. 2001. Soil algae from northern Victoria Land (Antarctica). *Polar Bioscience* 14:45-60.
- Chariton AA, Court LN, Hartley DM, Colloff MJ, Hardy CM. 2010. Ecological assessment of estuarine sediments by pyrosequencing eukaryotic ribosomal DNA. *Frontiers in Ecology and the Environment* 8(5):233-238.
- Chen Z, He C, Hu H. 2012. Temperature responses of growth, photosynthesis, fatty acid and nitrate reductase in Antarctic and temperate *Stichococcus*. *Extremophiles* 16(1):127-133.
- Cheung MK, Au CH, Chu KH, Kwan HS, Wong CK. 2010. Composition and genetic diversity of picoeukaryotes in subtropical coastal waters as revealed by 454 pyrosequencing. *The ISME Journal* 4(8):1053-1059.
- Chornevych TM, Maltseva IA, Nikorych VA. 2008. Soil algae of the coniferous forests of Ukraine. *Ekologia ta noosferologija* 18(3-4):77-82.
- Chunxiang H, Yongding L. 2003. Primary succession of algal community structure in desert soil. *Acta Botanica Sinica* 45(8):917-924.
- Clement M, Posada D, Crandall KA. 2000. TCS: a computer program to estimate gene genealogies. *Molecular Ecology* 9(10):1657-1659.

- Coleman AW. 2009. Is there a molecular key to the level of "biological species" in eukaryotes? A DNA guide. *Molecular Phylogenetics and Evolution* 50(1):197-203.
- Costello EK, Halloy SRP, Reed SC, Sowell P, Schmidt SK. 2009. Fumarole-supported islands of biodiversity within a hyperarid, high-elevation landscape on Socompa volcano, Puna de Atacama, Andes. *Applied and Environmental Microbiology* 75(3):735-747.
- Cox E, Deason T. 1969. *Heterochlamydomonas*, a new alga from Tennessee. *Journal of the Tennessee Academy of Sciences* 44:105-107.
- Dahlin S, Witter E, Mårtensson A, Turner A, Bååth E. 1997. Where's the limit? Changes in the microbiological properties of agricultural soils at low levels of metal contamination. *Soil Biology and Biochemistry* 29(9–10):1405-1415.
- Dal Grande F, Beck A, Cornejo C, Singh G, Cheenacharoen S, Nelsen MP, Scheidegger C. 2014. Molecular phylogeny and symbiotic selectivity of the green algal genus *Dictyochloropsis* s.l. (Trebouxiophyceae): a polyphyletic and widespread group forming photobiont-mediated guilds in the lichen family Lobariaceae. *New Phytologist* 202(2):455-470.
- Daniel R. 2005. The metagenomics of soil. *Nature Reviews Microbiology* 3(6):470-478.
- Darienko T, Friedl T, Pröschold T. 2009. *Desmochloris mollenhaueri* – a new terrestrial ulvophycean alga from south-west African soils. *Algological Studies* 129(1):25-40.
- Darienko T, Gustavs L, Mudimu O, Menendez CR, Schumann R, Karsten U, Friedl T, Pröschold T. 2010. *Chloroidium*, a common terrestrial coccoid green alga previously assigned to *Chlorella* (Trebouxiophyceae, Chlorophyta). *European Journal of Phycology* 45(1):79-95.
- Darienko T, Gustavs L, Eggert A, Wolf W, Pröschold T. 2015. Evaluating the species boundaries of Green microalgae (*Coccomyxa*, Trebouxiophyceae, Chlorophyta) using integrative taxonomy and DNA barcoding with further implications for the species identification in environmental samples. *PLoS ONE* 10(6):e0127838.
- Darienko T, Pröschold T. 2015. Genetic variability and taxonomic revision of the genus *Auxenochlorella* (Shihira et Krauss) Kalina et Punčochářová (Trebouxiophyceae, Chlorophyta). *Journal of Phycology* 51(2):394-400.
- Darienko TM. 2012. Terrestrial algae of Zmiinyj island (Black sea): general characteristic and peculiarities of species composition. *Ukrajinskij Botanichnij Zhurnal* 69(1):111-124.
- Darty K, Denise A, Ponty Y. 2009. VARNA: Interactive drawing and editing of the RNA secondary structure. *Bioinformatics* 25(15):1974-1975.
- Day JG, Pröschold T, Friedl T, Lorenz M, Silva PC. 2010. Conservation of microalgal type material: approaches needed for 21st century science. *Taxon* 59(1):3-6.
- de-Bashan LE, Trejo A, Huss VA, Hernandez J-P, Bashan Y. 2008. *Chlorella sorokiniana* UTEX 2805, a heat and intense, sunlight-tolerant microalga with potential for removing ammonium from wastewater. *Bioresource Technology* 99(11):4980-4989.
- De Wever A, Leliaert F, Verleyen E, Vanormelingen P, Van der Gucht K, Hodgson DA, Sabbe K, Vyverman W. 2009. Hidden levels of phylodiversity in Antarctic green algae: further evidence for the existence of glacial refugia. *Proceedings of the Royal Society B: Biological Sciences* 276(1673):3591-3599.
- Deason T, Bold H. 1960. Phycological studies 1. Exploratory studies of Texas soil algae. The University of Texas Press No. 6022. 71 pp.
- Deason TR, Floyd GL. 1987. Comparative ultrastructure of three species of *Chlorosarcina* (Chlorosarcinaceae, Chlorophyta). *Journal of Phycology* 23(1):187-195.

- del Campo J, Sieracki ME, Molestina R, Keeling P, Massana R, Ruiz-Trillo I. 2014. The others: our biased perspective of eukaryotic genomes. *Trends in Ecology and Evolution* 29(5):252-259.
- Dickie IA. 2010. Insidious effects of sequencing errors on perceived diversity in molecular surveys. *New Phytologist* 188(4):916-918.
- Doherty M, Costas BA, McManus GB, Katz LA. 2007. Culture-independent assessment of planktonic ciliate diversity in coastal northwest Atlantic waters. *Aquatic Microbial Ecology* 48(2):141-154.
- Dorigo U, Bérard A, Humbert J. 2002. Comparison of eukaryotic phytobenthic community composition in a polluted river by partial 18S rRNA gene cloning and sequencing. *Microbial Ecology* 44(4):372-380.
- Dorokhova MF, Kosheleva NE, Terskaya EV. 2015. Algae and Cyanobacteria in soils of Moscow. *American Journal of Plant Sciences* 6(15):2461-2471.
- Dotsenko KA. 2008. Study of soil algae as bioindicators in areas of petroleum production. *Nauchnyj Zhurnal of the Kuban State Agrarian University* 41(7):1-6.
- Driver T, Bajhaiya AK, Allwood JW, Goodacre R, Pittman JK, Dean AP. 2015. Metabolic responses of eukaryotic microalgae to environmental stress limit the ability of FT-IR spectroscopy for species identification. *Algal Research* 11:148-155.
- Drummond AJ, Rambaut A. 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* 7(214):doi:10.1186/1471-2148-7-214.
- Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* 29(8):1969-1973.
- Edgcomb V, Orsi W, Bunge J, Jeon S, Christen R, Leslin C, Holder M, Taylor GT, Suarez P, Varela R. 2011. Protistan microbial observatory in the Cariaco Basin, Caribbean. I. Pyrosequencing vs Sanger insights into species richness. *The ISME Journal* 5(8):1344-1356.
- Ehrenberg CG. 1833. Dritter Beitrag zur Erkenntniss grosser Organisation in der Richtung des kleinsten Raumes. Berlin: Königliche Akademie der Wissenschaften. p. 145-336
- Eliáš M, Neustupa J, Škaloud P. 2008. *Elliptochloris bilobata* var. *corticola* var. nov. (Trebouxiophyceae, Chlorophyta), a novel subaerial coccal green alga. *Biologia* 63(6):791-798.
- Eliáš M, Neustupa J. 2009. *Pseudomarvania*, gen. nov. (Chlorophyta, Trebouxiophyceae), a new genus for “budding” subaerial green algae *Marvania aerophytica* Neustupa et Šejnohová and *Stichococcus ampulliformis* Handa. *Fottea* 9(2):169-177.
- Eliáš M, Němcová Y, Škaloud P, Neustupa J, Kaufnerová V, Šejnohová L. 2010. *Hylodesmus singaporensis* gen. et sp. nov., a new autosporic subaerial green alga (Scenedesmaceae, Chlorophyta) from Singapore. *International Journal of Systematic and Evolutionary Microbiology* 60:1224-1235.
- Eliáš M, Neustupa J, Pažoutová M, Škaloud P. 2013. A case of taxonomic inflation in coccoid algae: *Ellipsoidion parvum* and *Neocystis vischeri* are conspecific with *Neocystis* (= *Nephrodiella*) *brevis* (Chlorophyta, Trebouxiophyceae). *Phytotaxa* 76(2):15-27.
- Elster J. 1999. Algal versatility in various extreme environments. Enigmatic microorganisms and life in extreme environments. Springer. p. 215-227.
- Elster J, Lukešová A, Svoboda J, Kopecký J, Kanda H. 1999. Diversity and abundance of soil algae in the polar desert Sverdrup Pass, central Ellesmere Island. *Polar Record* 35:231-254.

- Elster J, Benson E. 2004. Life in the polar terrestrial environment with a focus on algae and cyanobacteria. In: Fuller BJ, Lane N, Benson EE, editors. Life in the frozen state. CRC Press. p. 111-150.
- Ettl H. 1963. Massenhaftes Auftreten von Chloromonas-Arten in Regenpfützen. *Nova Hedwigia Beiheft* 6:395-400.
- Ettl H, Gärtner G. 1995. Syllabus der Boden-, Luft-und Flechtenalgen. Springer-Verlag. 721. pp.
- Ettl H, Gärtner G. 2014. Syllabus der Boden-, Luft-und Flechtenalgen. Springer-Verlag. 773. pp.
- Fairchild ED, Willson DL. 1967. The algal flora of two Washington soils. *Ecology* 48(6):1053-1055.
- Faßhauer F, Beck A, Hodač L, Kaufholz F, Probst P, Sasonowa M, Weber A, Friedl T. 2011. Biodiversity of terrestrial algae in tropical mountain habitats in Podocarpus National Park, Ecuador. Posters, *European Journal of Phycology* 46(Supplement). p. 166.
- Faßhauer F. in prep. Biodiversity of terrestrial algae in tropical mountain habitats in Podocarpus National Park, Ecuador. PhD thesis, Georg August University of Göttingen.
- Fawley MW, Fawley KP, Buchheim MA. 2004. Molecular diversity among communities of freshwater microchlorophytes. *Microbial Ecology* 48:489-499.
- Feher D. 1948. Researches on the geographical distribution of soil microflora. II. The geographical distribution of soil algae. *Communications of the Botanical Institute of the Hungarian University of Technical and Economic Sciences, Sopron (Hungary)* 21:1-35.
- Fenchel T, Finlay BJ. 2004. The ubiquity of small species: patterns of local and global diversity. *BioScience* 54(8):777-784.
- Fermani P, Mataloni G, Van de Vijver B. 2007. Soil microalgal communities on an antarctic active volcano (Deception Island, South Shetlands). *Polar Biology* 30(11):1381-1393.
- Fierer N, Cary C. 2014. Archiving: Don't let microbial samples perish. *Nature* 512(7514):253-253.
- Filker S, Gimmler A, Dunthorn M, Mahé F, Stoeck T. 2015. Deep sequencing uncovers protistan plankton diversity in the Portuguese Ria Formosa solar saltern ponds. *Extremophiles* 19(2):283-295.
- Finlay BJ, Clarke KJ. 1999. Ubiquitous dispersal of microbial species. *Nature* 400(6747):828-828.
- Finlay BJ. 2002. Global dispersal of free-living microbial eukaryote species. *Science* 296(5570):1061-1063.
- Fischer M, Bossdorf O, Gockel S, Hänsel F, Hemp A, Hessenmöller D, Korte G, Nieschulze J, Pfeiffer S, Prati D, Renner S, Schöning I, Schumacher U, Wells K, Buscot F, Kalko EKV, Linsenmair KE, Schulze E-D, Weisser WW. 2010. Implementing large-scale and long-term functional biodiversity research: The Biodiversity Exploratories. *Basic and Applied Ecology* 11(6):473-485.
- Flechtner VR, Johansen JR, Clark WH. 1998. Algal composition of microbiotic crusts from the central desert of Baja California, Mexico. *The Great Basin Naturalist* 58(4):295-311.
- Flechtner VR, Ng RA, Johansen JR, Antonio S. 2005. Algal diversity in North American desert soils. *Proc. SPIE* 5906, Astrobiology and Planetary Missions. p. 1-9.
- Flechtner VR, Johansen JR, Belnap J. 2008. The biological soil crusts of the San Nicolas island. *Western North American Naturalist* 68(4):405-436.
- Flechtner VR, Pietrasiak N, Lewis LA. 2013. Newly revealed diversity of green microalgae from wilderness areas of Joshua tree national park (JTNP). *Monographs of the Western North American Naturalist* 6:43-63.

- Foissner W, Chao A, Katz L. 2008. Diversity and geographic distribution of ciliates (Protista: Ciliophora). *Biodiversity and Conservation* 17(2):345-363.
- Fontaine KM, Beck A, Stocker-Wörgötter E, Piercey-Normore MD. 2012. Photobiont relationships and phylogenetic history of *Dermatocarpon luridum* var. *luridum* and related *Dermatocarpon* species. *Plants* 1(2):39-60.
- Fontaine KM, Stocker-Woergötter E, Booth T, Piercey-Normore MD. 2013. Genetic diversity of the lichen-forming alga, *Diplosphaera chodatii*, in North America and Europe. *The Lichenologist* 45(6):799-813.
- Ford T, Pedley H. 1996. A review of tufa and travertine deposits of the world. *Earth-Science Reviews* 41(3):117-175.
- Forney LJ, Zhou X, Brown CJ. 2004. Molecular microbial ecology: land of the one-eyed king. *Current Opinion in Microbiology* 7(3):210-220.
- Freeman KR, Pescador MY, Reed SC, Costello EK, Robeson MS, Schmidt SK. 2009. Soil CO₂ flux and photoautotrophic community composition in high-elevation, 'barren' soil. *Environmental Microbiology* 11(3):674-686.
- Frey B, Bühler L, Schmutz S, Zumsteg A, Furrer G. 2013. Molecular characterization of phototrophic microorganisms in the forefield of a receding glacier in the Swiss Alps. *Environmental Research Letters* 8(1):doi:10.1088/1748-9326/8/1/015033.
- Freytet P, Plet A. 1991. The modern stromatolitic formations (calcareous-tufa) of the Tournus area (Burgundy, France). *Geobios* 24(2):123-139.
- Freytet P, Plet A. 1996. Modern freshwater microbial carbonates: the *Phormidium* stromatolites (tufa-travertine) of southeastern Burgundy (Paris Basin, France). *Facies* 34(1):219-237.
- Freytet P, Verrecchia EP. 1998. Freshwater organisms that build stromatolites: a synopsis of biocrystallization by prokaryotic and eukaryotic algae. *Sedimentology* 45(3):535-563.
- Friedl T. 1996. Evolution of the polyphyletic genus *Pleurastrum* (Chlorophyta): inferences from nuclear-encoded ribosomal DNA sequences and motile cell ultrastructure. *Phycologia* 35(5):456-469.
- Friedl T, Rokitta C. 1997. Species relationships in the lichen alga *Trebouxia* (Chlorophyta, Trebouxiophyceae): molecular phylogenetic analyses of nuclear-encoded large subunit rRNA gene sequences. *Symbiosis* 23:125-148.
- Friedl T, O'Kelly CJ. 2002. Phylogenetic relationships of green algae assigned to the genus *Planophila* (Chlorophyta): evidence from 18S rDNA sequence data and ultrastructure. *European Journal of Phycology* 37(3):373-384.
- Friedl T, Rybalka N. 2012. Systematics of the Green Algae: A Brief Introduction to the current status. In: Lüttge U, editor. *Progress in Botany*. p. 259-280.
- Friedmann EI, Lipkin Y, Ocampo-Paus R. 1967. Desert algae of the Negev (Israel). *Phycologia* 6(4):185-200.
- Friedrich J, Dandekar T, Wolf M, Müller T. 2005. ProfDist: a tool for the construction of large phylogenetic trees based on profile distances. *Bioinformatics* 21(9):2108-2109.
- Frouz J, Keplin B, Pižl V, Tajovský K, Starý J, Lukešová A, Nováková A, Balík Vr, Háněl L, Materna J. 2001. Soil biota and upper soil layer development in two contrasting post-mining chronosequences. *Ecological Engineering* 17(2):275-284.
- Fučíková K, Rada JC, Lewis LA. 2011a. The tangled taxonomic history of *Dictyococcus*, *Bracteacoccus* and *Pseudomuriella* (Chlorophyceae, Chlorophyta) and their distinction based on a phylogenetic perspective. *Phycologia* 50(4):422-429.

- Fučíková K, Rada JC, Lukešová A, Lewis LA. 2011b. Cryptic diversity within the genus *Pseudomuriella* Hanagata (Chlorophyta, Chlorophyceae, Sphaeropleales) assessed using four barcode markers. *Nova Hedwigia* 93(1-2):29-46.
- Fučíková K, Flechtner VR, Lewis LA. 2012. Revision of the genus *Bracteacoccus* Tereg (Chlorophyceae, Chlorophyta) based on a phylogenetic approach. *Nova Hedwegia* 96(1-2):15-59.
- Fučíková K, Lewis LA. 2012. Intersection of *Chlorella*, *Muriella* and *Bracteacoccus*: Resurrecting the genus *Chromochloris* Kol et Chodat (Chlorophyceae, Chlorophyta). *Fottea* 12(1):83-93.
- Fučíková K, Lewis PO, Lewis LA. 2013. Putting incertae sedis taxa in their place: a proposal for ten new families and three new genera in Sphaeropleales (Chlorophyceae). *Journal of Phycology* 50(1):14-25.
- Fučíková K, Lewis PO, Lewis LA. 2014. Widespread desert affiliation of trebouxiophycean algae (Trebouxiophyceae, Chlorophyta) including discovery of three new desert genera. *Phycological Research* 62(4):294-305.
- Garcia-Pichel F, Al-Horani FA, Farmer JD, Ludwig R, Wade BD. 2004. Balance between microbial calcification and metazoan bioerosion in modern stromatolitic oncolites. *Geobiology* 2:49-57.
- Gärtner G. 1996. ASIB - the Culture Collection of Algae at the Botanical Institute of the University at Innsbruck (Austria) Catalogue of Strains. p. 45-69.
- Gärtner G, Ingolić E. 2003. Further studies on *Desmococcus* Brand emend. Vischer (Chlorophyta, Trebouxiophyceae) and a new species *Desmococcus spinocystis* sp. nov. from soil. *Biologia* 58(4):517-523.
- Gaysina L, Němcová Y, Škaloud P, Ševčíková T, Eliáš M. 2013. *Chloropyrula uraliensis* gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a new green coccoid alga with a unique ultrastructure, isolated from soil in South Urals. *Journal of Systematics and Evolution* 51(4):476-484.
- Genitsaris S, Moustaka-Gouni M, Kormas KA. 2011. Airborne microeukaryote colonists in experimental water containers: diversity, succession, life histories and established food webs. *Aquatic Microbial Ecology* 62(2):139-152.
- Germond A, Hata H, Fujikawa Y, Nakajima T. 2013. The phylogenetic position and phenotypic changes of a *Chlorella*-like alga during 5-year microcosm culture. *European Journal of Phycology* 48(4):485-496.
- Gilichinsky D, Wilson G, Friedmann E, McKay C, Sletten R, Rivkina E, Vishnivetskaya T, Erokhina L, Ivanushkina N, Kochkina G. 2007. Microbial populations in Antarctic permafrost: biodiversity, state, age, and implication for astrobiology. *Astrobiology* 7(2):275-311.
- Goiris K, Muylaert K, Voorspoels S, Noten B, De Paepe D, Baart E, Gino J, De Cooman L. 2014. Detection of flavonoids in microalgae from different evolutionary lineages. *Journal of Phycology* 50(3):483-492.
- Gollerbach MM, Shtina EA. 1969. Soil algae. Nauka. 228 pp.
- Gouy M, Guindon S, Gascuel O. 2010. SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution* 27(2):221-224.
- Gratetpanche J-D, Santoferrara LF, McManus GB, Katz LA. 2014. Diversity of diversity: conceptual and methodological differences in biodiversity estimates of eukaryotic microbes as compared to bacteria. *Trends in Microbiology* 22(8):432-437.

- Guiry MD. 2012. How many species of algae are there? *Journal of Phycology* 48(5):1057-1063.
- Guiry MD, Guiry GM. 2015. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <http://www.algaebase.org>.
- Gustavs L, Eggert A, Michalik D, Karsten U. 2010. Physiological and biochemical responses of green microalgae from different habitats to osmotic and matric stress. *Protoplasma* 243(1-4):3-14.
- Hahn MW, Koll U, Jezberová J, Camacho A. 2015. Global phylogeography of pelagic *Polynucleobacter* bacteria: Restricted geographic distribution of subgroups, isolation by distance and influence of climate. *Environmental Microbiology* 17(3):829-840.
- Hall JD, Fucikova K, Lo C, Lewis LA. 2010. An assessment of proposed DNA barcodes in freshwater green algae. *Cryptogamie, Algologie* 31(4):529-555.
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*. Oxford University Press. p. 95-98.
- Hallmann C, Hodač L, Pahlmann KC, Faßhauer F, Friedl T. 2009. Changes in the biodiversity of algae in soils along gradients of land use intensities. *Posters, Phycologia* 48 (Supplement). p. 32.
- Hallmann C, Fritzlar D, Stannek L, Hoppert M. 2011a. Ascomycete fungi on dimension stone of the “Burg Gleichen”, Thuringia. *Environmental Earth Sciences* 63(7-8):1713-1722.
- Hallmann C, Rüdrieh J, Enseleit M, Friedl T, Hoppert M. 2011b. Microbial diversity on a marble monument: a case study. *Environmental Earth Sciences* 63:1701–1711.
- Hallmann C, Stannek L, Fritzlar D, Hause-Reitner D, Friedl T, Hoppert M. 2013a. Molecular diversity of phototrophic biofilms on building stone. *FEMS Microbiology Ecology* 84(2):355-372.
- Hallmann C, Wedekind W, Hause-Reitner D, Hoppert M. 2013b. Cryptogam covers on sepulchral monuments and re-colonization of a marble surface after cleaning. *Environmental Earth Sciences* 69:1149–1160.
- Hallmann C, Hodač L, Faßhauer F, Hoppert M, Friedl T. in prep.-a. Green algal communities from soil and tree bark in middle European forests: a molecular comparison.
- Hallmann C, Hodač L, Friedl T. in prep.-b. Green algae in grassland and forest soils under different management types - a culture-independent approach.
- Hamby KR, Sims L, Issel L, Zimmer E. 1988. Direct ribosomal RNA sequencing: Optimization of extraction and sequencing methods for work with higher plants. *Plant Molecular Biology Reporter* 6(3):175-192.
- Hammer Ø, Harper DAT, Ryan PD. 2001. PAST: Paleontological Statistics software package for education and data analysis. *Palaeontologia Electronica* 4(1):1-9.
- Hanagata N. 1998. Phylogeny of the subfamily Scotielloccystoideae (Chlorophyceae, Chlorophyta) and related taxa inferred from 18S ribosomal RNA gene sequence data. *Journal of Phycology* 34(6):1049-1054.
- Handa S, Nakahara M, Tsubota H, Deguchi H, Nakano T. 2003. A new aerial alga, *Stichococcus ampulliformis* sp. nov. (Trebouxiophyceae, Chlorophyta) from Japan. *Phycological Research* 51(3):203-210.
- Hansgirg A. 1888. *Prodromus der Algenflora von Böhmen*. Commissions-Verlag von Fr. Řivnáč. Prague. 282 pp.
- Hastings KL, Smith LE, Lindsey ML, Blotsky LC, Downing GR, Zellars DQ, Downing JK, Corena-McLeod M. 2014. Effect of microalgae application on soil algal species diversity,

- cation exchange capacity and organic matter after herbicide treatments. *F1000Research* 3(281):doi:10.12688/f1000research.4016.1
- Häubner N, Schumann R, Karsten U. 2006. Aeroterrestrial microalgae growing in biofilms on facades - response to temperature and water stress. *Microbial Ecology* 51(3):285-293.
- Heath CR, Leadbeater BC, Callow ME. 1995. Effect of inhibitors on calcium carbonate deposition mediated by freshwater algae. *Journal of Applied Phycology* 7(4):367-380.
- Heeg JS, Wolf M. 2015. ITS2 and 18S rDNA sequence-structure phylogeny of *Chlorella* and allies (Chlorophyta, Trebouxiophyceae, Chlorellaceae). *Plant Gene* 4:20-28.
- Hegewald E, Hanagata N. 2000. Phylogenetic studies on Scenedesmaceae (Chlorophyta). *Archiv für Hydrobiologie. Supplementband, Algological Studies* 136:29-49.
- Hegewald E, Bock C, Krienitz L. 2013. A phylogenetic study on Scenedesmaceae with the description of a new species of *Pectinodesmus* and the new genera *Verrucodesmus* and *Chodatodesmus*. *Fottea* 13:149-164.
- Henley WJ, Hironaka JL, Guillou L, Buchheim MA, Buchheim JA, Fawley MW, Fawley KP. 2004. Phylogenetic analysis of the 'Nannochloris-like' algae and diagnoses of *Picochlorum oklahomensis* gen. et sp. nov. (Trebouxiophyceae, Chlorophyta). *Phycologia* 43(6):641-652.
- Hepperle D, Krienitz L. 1996. The extracellular calcification of zoospores of *Phacotus lenticularis* (Chlorophyta, Chlamydomonadales). *European Journal of Phycology* 31(1):11-21.
- Hill MO. 1979. TWINSpan - A FORTRAN programme for arranging multivariate data in an ordered two-way table by classification of individuals and attributes. *Section of Ecology and Systematics*. 90. pp.
- Hirsch PR, Mauchline TH, Clark IM. 2010. Culture-independent molecular techniques for soil microbial ecology. *Soil Biology and Biochemistry* 42(6):878-887.
- Hodač L, Hallmann C, Rosenkranz H, Faßhauer F, Friedl T. 2012. Molecular evidence for the wide distribution of two lineages of terrestrial Green algae (Chlorophyta) over tropics to temperate zone. *ISRN Ecology*:doi:10.5402/2012/795924.
- Hodač L, Brinkmann N, Mohr KI, Arp G, Hallmann C, Ramm J, Spitzer K, Friedl T. 2015. Diversity of microscopic green algae (Chlorophyta) in calcifying biofilms of two karstic streams in Germany. *Geomicrobiology Journal* 32(3-4):275-290.
- Hodač L, Hallmann C, Spitzer K, Elster J, Faßhauer F, Brinkmann N, Lepka D, Diwan V, Friedl T. subm. Phylogenetic analysis of polar *Chlorella* and *Stichococcus* suggests biogeography of airborne microalgae. *FEMS Microbiology Ecology*.
- Hoffmann L. 1989. Algae of terrestrial habitats. *The Botanical Review* 55(2):77-105.
- Hoffmann L, Ector L, Kostikov I. 2007. Algal flora from limed and unlimed forest soils in the Ardenne (Belgium). *Systematics and Geography of Plants*:15-90.
- Hong JW, Jo SW, Cho HW, Nam SW, Shin W, Park KM, Lee KI, Yoon HS. 2015. Phylogeny, morphology, and physiology of *Micractinium* strains isolated from shallow ephemeral freshwater in Antarctica. *Phycological Research* 63(3):212-218.
- Hope ACA. 1968. A simplified Monte Carlo significance test procedure. *Journal of the Royal Statistical Society: Series B* 30:581-598.
- Hoppert M, Reimer R, Kemmling A, Schröder A, Günzl B, Heinken T. 2004. Structure and reactivity of a biological soil crust from a xeric sandy soil in Central Europe. *Geomicrobiology Journal* 21(3):183-191.
- Horath T, Bachofen R. 2009. Molecular characterization of an endolithic microbial community in dolomite rock in the Central Alps (Switzerland). *Microbial Ecology* 58:290-306.

- Hu C, Liu Y, Song L, Zhang D. 2002. Effect of desert soil algae on the stabilization of fine sands. *Journal of Applied Phycology* 14:281-292.
- Hu C, Liu Y, Paulsen BS, Petersen D, Klaveness D. 2003a. Extracellular carbohydrate polymers from five desert soil algae with different cohesion in the stabilization of fine sand grain. *Carbohydrate Polymers* 54(1):33-42.
- Hu C, Zhang D, Huang Z, Liu Y. 2003b. The vertical microdistribution of cyanobacteria and green algae within desert crusts and the development of the algal crusts. *Plant and Soil* 257(1):97-111.
- Hu H, Li H, Xu X. 2008. Alternative cold response modes in *Chlorella* (Chlorophyta, Trebouxiophyceae) from Antarctica. *Phycologia* 47(1):28-34.
- Huber-Pestalozzi G, Fott B, Komárek J, Förster K. 1983. *Das Phytoplankton des Süßwassers: Systematik und Biologie*. E. Schweizerbart. 606 pp.
- Huber T, Faulkner G, Hugenholtz P. 2004. Bellerophon: a program to detect chimeric sequences in multiple sequence alignments. *Bioinformatics* 20(14):2317-2319.
- Hughes KA. 2006. Solar UV-B radiation, associated with ozone depletion, inhibits the Antarctic terrestrial microalga, *Stichococcus bacillaris*. *Polar Biology* 29:327-336.
- Hunt ME, Floyd GL, Stout BB. 1979. Soil algae in field and forest environments. *Ecology* 60(2):362-375.
- Huson DH, Bryant D. 2006. Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution* 23(2):254-267.
- Huss VAR, Ciniglia C, Cennamo P, Cozzolino S, Pinto G, Pollio A. 2002. Phylogenetic relationships and taxonomic position of *Chlorella*-like isolates from low pH environments (pH < 3.0). *BMC Evolutionary Biology* 2:doi:10.1186/1471-2148-2-13.
- Iljushenko AE. 2008. Algotkompleksy pochvennykh vodoroslej bora-brusnichnika. *Fundamentalnye i prikladnye problemy botaniki v nachale XXI veka. Algologija*. p. 37-39.
- Islam N. 1960. Some subaerial green algae from East Pakistan. *Transactions of the American Microscopical Society* 10(4):471-479.
- Jaarsma CF, van Langevelde F, Botma H. 2006. Flattened fauna and mitigation: Traffic victims related to road, traffic, vehicle, and species characteristics. *Transportation Research Part D: Transport and Environment* 11(4):264-276.
- Jobb G, Von Haeseler A, Strimmer K. 2004. TREEFINDER: a powerful graphical analysis environment for molecular phylogenetics. *BMC Evolutionary Biology* 4(1):doi:10.1186/1471-2148-4-18.
- Johansen JR, Casamatta DA. 2005. Recognizing cyanobacterial diversity through adoption of a new species paradigm. *Algological Studies* 117(1):71-93.
- Johnson LR, John DM. 1992. Taxonomic observations on some uncommon and new *Gongrosira* species (Chaetophorales sensu stricto, division Chlorophyta). *European Journal of Phycology* 27(2):153-163.
- Jurdíková K, Kulichová J, Bestová H, Leliaert F, Škaloud P. 2014. Exploration of nuclear DNA markers for population structure assessment in the desmid *Micrasterias rotata* (Zygnematophyceae, Streptophyta). *Journal of Eukaryotic Microbiology* 61(5):509-519.
- Kabirov R, Gaisina L, Safiullina L. 2010. Universal criteria for assessing the ecological state of soil algocenoses. *Russian Journal of Ecology* 41(4):302-306.
- Kabirov RR. 1993. Algoindications with using soil algae (methodological aspects). *Algologia* 3(3):73-83.

- Kalinowska R, Trzcinska M, Pawlik-Skowronska B. 2008. Glony glebowe terenow pogornicznych skazonych metalami ciezkimi. *Wiadomości Botaniczne* 52(3-4):63-79.
- Kalinowska R, Pawlik-Skowrońska B. 2010. Response of two terrestrial green microalgae (Chlorophyta, Trebouxiophyceae) isolated from Cu-rich and unpolluted soils to copper stress. *Environmental Pollution* 158(8):2778-2785.
- Karbovska VM, Kostikov IY. 2012a. New morphological features of the members of genus *Stichococcus* Nägeli (Trebouxiophyceae, Chlorophyta). *Modern Phytomorphology* 1:111-114.
- Karbovska VM, Kostikov IY. 2012b. *Chlorella sphaerica* and its position in the genus *Diplosphaera* (Chlorophyta, Trebouxiophyceae). *Visnyk of Dnipropetrovsk University. Biology, Ecology* 20(1):34-42.
- Karsten U, Friedl T, Schumann R, Hoyer K, Lembcke S. 2005. Mycosporine-like amino acids and phylogenies in green algae: *Prasiola* and its relationships from the Trebouxiophyceae (Chlorophyta). *Journal of Phycology* 41(3):557-566.
- Kaštovská K, Elster J, Stibal M, Šantrůčková H. 2005. Microbial assemblages in soil microbial succession after glacial retreat in Svalbard (High Arctic). *Microbial Ecology* 50(3):396-407.
- Kaštovská K, Stibal M, Šabacká M, Černá B, Šantrůčková H, Elster J. 2007. Microbial community structure and ecology of subglacial sediments in two polythermal Svalbard glaciers characterized by epifluorescence microscopy and PLFA. *Polar Biology* 30(3):277-287.
- Katoh K, Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics* 9(4):286-298.
- Kaufnerová V, Eliáš M. 2013. The demise of the genus *Scotiellopsis* Vinatzer (Chlorophyta). *Nova Hedwigia* 97(3-4):415-428.
- Keller A, Schleicher T, Schultz J, Müller T, Dandekar T, Wolf M. 2009. 5.8S-28S rRNA interaction and HMM-based ITS2 annotation. *Gene* 430(1-2):50-57.
- Keller A. 2011. Secondary (and tertiary) structure of the ITS2 and its application for phylogenetic tree reconstructions and species identification. PhD thesis. Universität Würzburg.
- Khan N, Tuffin M, Stafford W, Cary C, Lacap DC, Pointing SB, Cowan D. 2011. Hypolithic microbial communities of quartz rocks from Miers Valley, McMurdo Dry Valleys, Antarctica. *Polar Biology* 34(11):1657-1668.
- Khaybullina LS, Gaysina L, Johansen JR, Krautová M. 2010. Examination of the terrestrial algae of the Great Smoky Mountains National Park, USA. *Fottea* 10(2):201-215.
- Kim GH, Klochkova TA, Kang SH. 2008. Notes on freshwater and terrestrial algae from Ny-Alesund, Svalbard (high Arctic sea area). *Journal of Environmental Biology* 29(4):485-491.
- King JM, Ward C. 1977. Distribution of edaphic algae as related to land usage. *Phycologia* 16(1):23-30.
- Kochkina G, Ozerskaya S, Ivanushkina N, Chigineva N, Vasilenko O, Spirina E, Gilichinskii D. 2014. Fungal diversity in the Antarctic active layer. *Microbiology* 83(1-2):94-101.
- Koetschan C, Forster F, Keller A, Schleicher T, Ruderisch B, Schwarz R, Müller T, Wolf M, Schultz J. 2010. The ITS2 Database III - sequences and structures for phylogeny. *Nucleic Acids Research* 38:275-279.
- Koetschan C, Hackl T, Müller T, Wolf M, Forster F, Schultz J. 2012. ITS2 database IV: interactive taxon sampling for internal transcribed spacer 2 based phylogenies. *Molecular Biology and Evolution* 63(3):585-588.

- Kol E. 1967. Algal growth experiments in the Baradla cave at Aggtelek. *International Journal of Speleology* 2:457-474.
- Komárek J, Anagnostidis K. 1998. Cyanoprokaryota – 1st Part: Chroococcales. Stuttgart, Germany: Gustav Fischer Verlag. 548 pp.
- Komárek J, Anagnostidis K. 2005. Cyanoprokaryota – 2nd Part: Oscillatoriales. Stuttgart, Germany: Gustav Fischer Verlag. 759 pp.
- Komárek J. 2013. Cyanoprokaryota – 3rd Part: Nostocales. Stuttgart, Germany: Gustav Fischer Verlag. 1131 pp.
- Komaromy ZP. 1976. Soil algal growth types as edaphic adaptation in Hungarian forest and grassland ecosystems. *Acta Botanica Hungarica* 22:373-379.
- Komaromy ZP. 1983. A comparative study on the algal synusia of Hungarian grasslands and deciduous forests. *Annales Historico-Naturales Musei Nationalis Hungarici* 75:47-53.
- Kostikov I, Carnol M, Dulière J, Hoffman L. 2001a. Effects of limiting on forest soil algal communities. *Algological Studies* 102:161-178.
- Kostikov I, Romanenko PO, Demchenko EM, Darienko TM, Mikhailyuk TI, Rybchnnskiy OV, Solonenko AM. 2001b. Soil algae of Ukraine: methods of study, system, checklist. In: Kondratiuk SJ, Massjiuk NP, editors. *Phytosociocenter*. 300 pp.
- Krienitz L, Hegewald E, Hepperle D, Wolf M. 2003. The systematics of coccoid green algae: 18S rRNA gene sequence data versus morphology. *Biologia* 58(4):437-446.
- Krienitz L, Hegewald EH, Hepperle D, Huss VA, Rohr T, Wolf M. 2004. Phylogenetic relationship of *Chlorella* and *Parachlorella* gen. nov. (Chlorophyta, Trebouxiophyceae). *Phycologia* 43(5):529-542.
- Krienitz L, Bock C, Dadheech P, Pröschold T. 2011. Taxonomic reassessment of the genus *Mychonastes* (Chlorophyceae, Chlorophyta) including the description of eight new species. *Phycologia* 50(1):89-106.
- Krienitz L, Bock C. 2012. Present state of the systematics of planktonic coccoid green algae of inland waters. *Hydrobiologia* 698(1):295-326.
- Krienitz L, Huss VAR, Bock C. 2015. *Chlorella*: 125 years of the green survivalist. *Trends in Plant Science* 20(2):67-69.
- Krivorotov SB, Bukareva OV. 2005. Soil algae as biological indicators of soil pollution of protected areas of the north-western Caucasus with heavy metals. *Uspechi Sovremennogo Estestvovania* 11:12-15.
- Kroken S, Taylor JW. 2000. Phylogenetic species, reproductive mode, and specificity of the green alga *Trebouxia* forming lichens with the fungal genus *Letharia*. *The Bryologist* 103(4):645-660.
- Kulichová J, Škaloud P, Neustupa J. 2014. Molecular diversity of green corticolous microalgae from two sub-Mediterranean European localities. *European Journal of Phycology* 49(3):345-355.
- Kutovaya O, Vasilenko E, Lebedeva M. 2012. Microbiological and micromorphological characteristics of extremely arid desert soils in the Ili Depression (Kazakhstan). *Eurasian Soil Science* 45(12):1147-1158.
- Kützing FT. 1843. *Phycologia generalis: oder Anatomie, Physiologie und Systemkunde der Tange*. Leipzig: Brockhaus. 458 pp.
- Kuzyakhmetov GG. 1998. Productivity of algocenoses in zonal arable soils of steppe and forest-steppe. *Eurasian Soil Science* 31(4):406-410.

- Kvíděrová J, Lukavský J. 2005. The comparison of ecological characteristics of *Stichococcus* (Chlorophyta) strains isolated from polar and temperate regions. *Algological Studies* 118(1):127-40.
- Lang I, Hodač L, Friedl T, Feussner I. 2011. Fatty acid profiles and their distribution patterns in microalgae: a comprehensive analysis of more than 2000 strains from the SAG culture collection. *BMC Plant Biology* 11(124):doi:10.1186/1471-2229-11-124.
- Langhans TM, Storm C, Schwabe A. 2009. Community assembly of biological soil crusts of different successional stages in a temperate sand ecosystem, as assessed by direct determination and enrichment techniques. *Microbial Ecology* 58(2):394-407.
- Lara E, Berney C, Harms H, Chatzinotas A. 2007. Cultivation-independent analysis reveals a shift in ciliate 18S rRNA gene diversity in a polycyclic aromatic hydrocarbon-polluted soil. *FEMS Microbiology Ecology* 62(3):365-373.
- Lauber CL, Hamady M, Knight R, Fierer N. 2009. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applied and Environmental Microbiology* 75(15):5111-5120.
- Lawley B, Ripley S, Bridge P, Convey P. 2004. Molecular analysis of geographic patterns of eukaryotic diversity in Antarctic soils. *Applied and Environmental Microbiology* 70(10):5963-5672.
- Lee JC, Gutell RR. 2012. A comparison of the crystal structures of eukaryotic and bacterial SSU ribosomal RNAs reveals common structural features in the hypervariable regions. *PLoS ONE* 7(5):e38203.
- Leff JW, Jones SE, Prober SM, Barberán A, Borer ET, Firn JL, Harpole WS, Hobbie SE, Hofmockel KS, Knops JM. 2015. Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *Proceedings of the National Academy of Sciences*:doi:10.1073/pnas.1508382112.
- Leliaert F, Verbruggen H, Wysor B, De Clerck O. 2009. DNA taxonomy in morphologically plastic taxa: algorithmic species delimitation in the *Boodlea* complex (Chlorophyta: Cladophorales). *Molecular Phylogenetics and Evolution* 53(1):122-133.
- Leliaert F, Smith DR, Herron MD, Verbruggen H, Delwiche CF, Clerck OD. 2012. Phylogeny and molecular evolution of the Green algae. *Critical Reviews in Plant Sciences* 31(1):1-46.
- Leliaert F, Verbruggen H, Vanormelingen P, Steen F, Lopez-Bautista JM, Zuccarello GC, De Clerck O. 2014. DNA-based species delimitation in algae. *European Journal of Phycology* 49(2):179–196.
- Leliaert F, Lopez-Bautista JM. 2015. The chloroplast genomes of *Bryopsis plumosa* and *Tydemania expeditiones* (Bryopsidales, Chlorophyta): compact genomes and genes of bacterial origin. *BMC Genomics* 16(1):doi:10.1186/s12864-015-1418-3.
- Lemieux C, Otis C, Turmel M. 2014. Chloroplast phylogenomic analysis resolves deep-level relationships within the green algal class Trebouxiophyceae. *BMC Evolutionary Biology* 14(1):doi:10.1186/s12862-014-0211-2.
- Lepka D. 2007. Phylogenie *Chlorella*-ähnlicher Grünalgen von rDNA-Sequenzanalysen. MSc thesis, Georg-August-Universität Göttingen. 94 pp.
- Lesaulnier C, Papamichail D, McCorkle S, Ollivier B, Skiena S, Taghavi S, Zak D, Van Der Lelie D. 2008. Elevated atmospheric CO₂ affects soil microbial diversity associated with trembling aspen. *Environmental Microbiology* 10(4):926-941.
- Lewis LA, Flechtner VR. 2002. Green algae (Chlorophyta) of desert microbiotic crusts: diversity of North American taxa. *Taxon* 41:443-451.

- Lewis LA, Flechtner VR. 2004. Cryptic species of *Scenedesmus* (Chlorophyta) from desert soil communities of Western North America. *Journal of Phycology* 40:1127-1137.
- Lewis LA, Lewis PO. 2005. Unearthing the molecular phylodiversity of desert soil green algae (Chlorophyta). *Systematic Biology* 54(6):936-947.
- Li H, Liu X, Wang Y, Hu H, Xu X. 2009. Enhanced expression of antifreeze protein genes drives the development of freeze tolerance in an Antarctica isolate of *Chlorella vulgaris*. *Progress in Natural Science* 19(9):1059-1062.
- Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25(11):1451-1452.
- Lie AA, Liu Z, Hu SK, Jones AC, Kim DY, Countway PD, Amaral-Zettler LA, Cary SC, Sherr EB, Sherr BF. 2014. Investigating microbial eukaryotic diversity from a global census: insights from a comparison of pyrotag and full-length sequences of 18S rRNA genes. *Applied and Environmental Microbiology* 80(14):4363-4373.
- Lin C-s, Lin Y-h, Wu J-t. 2012. Biodiversity of the epiphyllous algae in a *Chamaecyparis* forest of northern Taiwan. *Botanical Studies* 53:489-499.
- Lin C-S, Chou T-L, Wu J-T. 2013. Biodiversity of soil algae in the farmlands of mid-Taiwan. *Botanical Studies* 54(41):doi:10.1186/1999-3110-54-41.
- Lin C-S, Wu J-T. 2014. Environmental factors affecting the diversity and abundance of soil photomicrobes in arid lands of subtropical Taiwan. *Geomicrobiology Journal* 31(4):350-359.
- Lindstrøm E-A, Johansen SW, Saloranta T. 2004. Periphyton in running waters - long-term studies of natural variation. *The Atna River: Studies in an Alpine-Boreal Watershed*. Springer. p. 63-86.
- López-Bautista JM, Chapman RL. 2003. Phylogenetic affinities of the Trentepohliales inferred from small-subunit rDNA. *International Journal of Systematic and Evolutionary Microbiology* 53(6):2099-2106.
- López-Bautista JM, Rindi F, Casamatta D. 2007. The systematics of subaerial algae. In: Seckbach J, editor. *Algae and cyanobacteria in extreme environments*. Springer. p. 599-617.
- Lu Y, Chi X, Yang Q, Li Z, Liu S, Gan Q, Qin S. 2009. Molecular cloning and stress-dependent expression of a gene encoding $\Delta 12$ -fatty acid desaturase in the Antarctic microalga *Chlorella vulgaris* NJ-7. *Extremophiles* 13(6):875-84.
- Lu Y, Chi X, Li Z, Yang Q, Li F, Liu S, Gan Q, Qin S. 2010. Isolation and characterization of a stress-dependent plastidial $\Delta 12$ fatty acid desaturase from the Antarctic microalga *Chlorella vulgaris* NJ-7. *Lipids* 45(2):179-187.
- Ludwig W, Strunk O, Westram R. 2004. ARB: a software environment for sequence data. *Nucleic Acids Research* 32(4):1363-1371.
- Lukešová A, Komárek J. 1987. Succession of soil algae on dumps from strip coal-mining in the Most region (Czechoslovakia). *Folia Geobotanica et Phytotaxonomica* 22(4):355-362.
- Lukešová A. 1993. Soil algae in four secondary succesional stages on abandoned fields. *Algological Studies* 71:81-102.
- Lukešová A, Hoffmann L. 1996. Soil algae from acid rain impacted forest areas of the Krušné hory Mts. 1. Algal communities. *Vegetatio* 125(2):123-136.
- Lukešová A. 2001. Soil algae in brown coal and lignite post-mining areas in Central Europe (Czech Republic and Germany). *Restoration Ecology* 9(4):341-350.
- Lukešová A, Kociánová M, Váňa J, Štursová H, Elster J, Harčarik J, Halda J, Kocourková J, Jankovská V. 2010. Mud boils of the Giant Mts and Abisko Mts tundra – preliminary comparative study. *Opera Corcontica* 47:55-82.

- Lund JW. 1945. Observations on soil algae. *New Phytologist* 44(2):196-196.
- Lund JW. 1947. Observations on soil algae. II. Notes on groups other than diatoms. *New Phytologist* 46(1):35-60.
- Luo W, Pröschold T, Bock C, Krienitz L. 2010. Generic concept in *Chlorella*-related coccoid green algae (Chlorophyta, Trebouxiophyceae). *Plant Biology* 12:545-553.
- Lutz S, Anesio AM, Edwards A, Benning LG. 2015. Microbial diversity on Icelandic glaciers and ice caps. *Frontiers in Microbiology* 6:doi:10.3389/fmicb.2015.00307.
- Ma S, Han B, Huss VA, Hu X, Sun X, Zhang J. 2015. *Chlorella thermophila* (Trebouxiophyceae, Chlorophyta), a novel thermo-tolerant *Chlorella* species isolated from an occupied rooftop incubator. *Hydrobiologia* 760:81-89.
- MacEntee FJ, Bold HC, Archibald PA. 1977. Notes on some edaphic algae of the South Pacific and Malaysian areas, with special reference to *Pseudotetraedron polymorphum* gen. et spec. nov. *Soil Science* 124(3):161-166.
- Maistro S, Broady PA, Andreoli C, Negrisola E. 2009. Phylogeny and Taxonomy of Xanthophyceae (Stramenopiles, Chromalveolata). *Protist* 160(3):412-426.
- Mallet J. 2005. Hybridization as an invasion of the genome. *Trends in Ecology and Evolution* 20(5):229-237.
- Maltsev EI. 2013. Ecological features of algae communities in forest floor of floodplain oak woods in steppe area of Ukraine. *Gruntoznavstvo* 14(3):330-339.
- Maltsev EI, Negrulja DM. 2013. Algal flora of forest floor in plantings of park-monument landscape art "forest nursery". *Gruntoznavstvo* 14(3):98-105.
- Maltseva IA. 2005a. Soil algae of flooded forests of the steppe zone of Ukraine. Melitopol: Liuks. p. 44-53.
- Maltseva IA. 2005b. Soil algae from the forests of gorges of the steppe zone of Ukraine. *Ekologija ta Noosferologija* 16(3):150-156.
- Maltseva IA. 2007. Algae groups of pine-woods of sandy terracing of river valleys in steppe area of Ukraine. *Visnik Charkivskogo natsionalnogo agrarnogo universitetu, serija biologija*. p. 106-112.
- Maltseva IA, Baranova OO, Maltsev EI. 2009. Algogrouppings of biogeocenoses landscape-technogenic systems of the Kryvoy Rog region. *Visnik Zaporizkogo Nacionalnogo Universitetu*. p. 20-23.
- Mansour HA, Shaaban AS. 2010. Algae of soil surface layer of Wadi Al-Hitan Protective Area (World Heritage Site), El-Fayum Depression, Egypt. *Journal of American Science* 6(8):243-255.
- Marande W, López-García P, Moreira D. 2009. Eukaryotic diversity and phylogeny using small- and large-subunit ribosomal RNA genes from environmental samples. *Environmental Microbiology* 11(12):3179-3188.
- Marschner P, Kandeler E, Marschner B. 2003. Structure and function of the soil microbial community in a long-term fertilizer experiment. *Soil Biology and Biochemistry* 35(3):453-461.
- Martiny JBH, Bohannan BJ, Brown JH, Colwell RK, Fuhrman JA, Green JL, Horner-Devine MC, Kane M, Krumins JA, Kuske CR. 2006. Microbial biogeography: putting microorganisms on the map. *Nature Reviews Microbiology* 4(2):102-112.
- Mataloni G, Tell G, Wynn-Williams D. 2000. Structure and diversity of soil algal communities from Cierva Point (Antarctic Peninsula). *Polar Biology* 23(3):205-211.
- Matula J, Pietryka M, Richter D, Wojtun B. 2007. Cyanoprokaryota and algae of Arctic terrestrial ecosystems in the Hornsund area, Spitsbergen. *Polish Polar Research* 28(4):283-315.

- Maxwell CD. 1991. Floristic changes in soil algae and cyanobacteria in reclaimed metal-contaminated land at Sudbury, Canada. *Water, Air, and Soil Pollution* 60:381-393.
- McNeill J, Barrie F, Buck W, Demoulin V, Greuter W, Hawksworth D, Herendeen P, Knapp S, Marhold K, Prado J. 2012. International code of nomenclature for algae, fungi, and plants (Melbourne Code). Koeltz Scientific Books. ISBN 978-3-87429-425-6. 232 pp.
- Medinger R, Nolte V, Pandey RV, Jost S, Ottenwaelder B, Schloetterer C, Boenigk J. 2010. Diversity in a hidden world: potential and limitation of next-generation sequencing for surveys of molecular diversity of eukaryotic microorganisms. *Molecular Ecology* 19:32-40.
- Megharaj M, Venkateswarlu K, Rao AS. 1986. Growth response of four species of soil algae to monocrotophos and quinalphos. *Environmental Pollution Series A, Ecological and Biological* 42(1):15-22.
- Megharaj M, Kantachote D, Singleton I, Naidu R. 2000. Effects of long-term contamination of DDT on soil microflora with special reference to soil algae and algal transformation of DDT. *Environmental Pollution* 109(1):35-42.
- Metting B, Rayburn W. 1979. Algal communities and soil microenvironments in an eastern Washington silt loam. *Soil Science* 127(2):doi:10.1097/00010694-197902000-00003.
- Metting B. 1981. The systematics and ecology of soil algae. *The Botanical Review* 47(2):195-312.
- Meyen FJF. 1829. Beobachtungen über einige niedere Algenformen. Weber. p. 769-778.
- Mikhailyuk TI, Sluiman HJ, Massalski A, Mudimu O, Demchenko EM, Kondratyuk SY, Friedl T. 2008. New streptophyte green algae from terrestrial habitats and an assessment of the genus *Interfilum* (Klebsormidiophyceae, Streptophyta). *Journal of Phycology* 44(6):1586-1603.
- Mocali S, Benedetti A. 2010. Exploring research frontiers in microbiology: the challenge of metagenomics in soil microbiology. *Research in Microbiology* 161(6):497-505.
- Moewus L. 1951. Systematische Bestimmung einzelliger grüner Algen auf Grund von Kulturversuchen. *Botaniska Notiser* 4:287-318.
- Mondav R, Woodcroft BJ, Kim E-H, McCalley CK, Hodgkins SB, Crill PM, Chanton J, Hurst GB, VerBerkmoes NC, Saleska SR, Hugenholtz P, Rich VI, Tyson GW. 2014. Discovery of a novel methanogen prevalent in thawing permafrost. *Nature Communications* 5: doi:10.1038/ncomms4212.
- Moon-van der Staay S, De Wachter R, Vaultot D. 2001. Oceanic 18S rRNA gene sequences from picoplankton reveal unsuspected eukaryotic diversity. *Nature* 409:607-610.
- Moon-van der Staay SY, Tzeneva VA, van Der Staay GW, De Vos WM, Smidt H, Hackstein JH. 2006. Eukaryotic diversity in historical soil samples. *FEMS Microbiology Ecology* 57(3):420-428.
- Moreira D, Lopez-Garcia P. 2002. The molecular ecology of microbial eukaryotes unveils a hidden world. *Trends in Microbiology* 10(1):31-38.
- Mudimu O, Rybalka N, Bauersachs T, Born J, Friedl T, Schulz R. 2014. Biotechnological screening of microalgal and cyanobacterial strains for biogas production and antibacterial and antifungal effects. *Metabolites* 4(2):373-393.
- Müller J, Day JG, Harding K, Hepperle D, Lorenz M, Friedl T. 2007a. Assessing genetic stability of a range of terrestrial microalgae after cryopreservation using amplified fragment length polymorphism (AFLP). *American Journal of Botany* 94(5):799-808.

- Müller T, Rahmann S, Dandekar T, Wolf M. 2004. Accurate and robust phylogeny estimation based on profile distances: a study of the Chlorophyceae (Chlorophyta). *BMC Evolutionary Biology* 4(1):doi:10.1186/1471-2148-4-20.
- Müller T, Philippi N, Dandekar T, Schultz J, Wolf M. 2007b. Distinguishing species. *RNA* 13(9):1469-1472.
- Musabaeva IA. Osobennosti organizacii gruppirovok pochvennykh vodoroslej stepnogo fitocenoza. http://iesen.nspu.ru/images/sborn/nr2011/kaf_botaniki/ch2.html.
- Nacke H, Thurmer A, Wollherr A, Will C, Hodac L, Herold N, Schoning I, Schrupf M, Daniel R. 2011. Pyrosequencing-based assessment of bacterial community structure along different management types in German forest and grassland soils. *PLoS ONE* 6(2):doi:10.1371/journal.pone.0017000.
- Nägeli C. 1849. *Gattungen einzelliger Algen*. Schulthess. 139 pp.
- Nagy ML, Johansen JR, Clair LLS, Webb BL. 2005. Recovery patterns of microbiotic soils crusts 70 years after arsenic contamination. *Journal of Arid Environments* 63:304-323.
- Nakada T, Misawa K, Nozaki H. 2008. Molecular systematics of Volvocales (Chlorophyceae, Chlorophyta) based on exhaustive 18S rRNA phylogenetic analyses. *Molecular Phylogenetics and Evolution* 48(1):281-291.
- Naselli-Flores L, Padisák J. 2015. Blowing in the wind: how many roads can a phytoplankton walk down? A synthesis on phytoplankton biogeography and spatial processes. *Hydrobiologia*:doi:10.1007/s10750-015-2519-3.
- Neefs J-M, De Wachter R. 1990. A proposal for the secondary structure of a variable area of eukaryotic small ribosomal subunit RNA involving the existence of a pseudoknot. *Nucleic Acids Research* 18(19):5695-5704.
- Němcová Y, Eliáš M, Škaloud P, Hodač L, Neustupa J. 2011. *Jenufa* gen. nov.: a new genus of coccoid green algae (Chlorophyceae, incertae sedis) previously recorded by environmental sequencing. *Journal of Phycology* 47(4):928-938.
- Néron B, Ménager H, Maufrais C, Joly N, Maupetit J, Letort S, Carrere S, Tuffery P, Letondal C. 2009. Mobyle: a new full web bioinformatics framework. *Bioinformatics* 25(22):3005-3011.
- Neustupa J. 2001. Soil algae from marlstone-substratum based biotopes in the Nature park Džbán (Central Bohemia, Czech Republic) with special attention to the natural treeless localities. *Archiv für Hydrobiologie. Supplementband, Algological Studies* 137:109-120.
- Neustupa J, Škaloud P. 2004. Contribution to the knowledge of soil algae of two abandoned industrial sedimentation basins in Eastern Bohemia. In: Kovář P, editor. *Natural recovery of human-made deposits in landscape (Biotic interactions and ore/ash-slag artificial ecosystems)*. Prague: Academia. p. 194-199.
- Neustupa J, Eliáš M, Šejnohová L. 2007. A taxonomic study of two *Stichococcus* species (Trebouxiophyceae, Chlorophyta) with a starch-enveloped pyrenoid. *Nova Hedwigia* 84(1-2):51-63.
- Neustupa J, Škaloud P. 2008. Diversity of subaerial algae and cyanobacteria on tree bark in tropical mountain habitats. *Biologia* 63(6):806-812.
- Neustupa J, Němcová Y, Eliáš M, Škaloud P. 2009. *Kalinella bambusicola* gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a novel coccoid *Chlorella*-like subaerial alga from Southeast Asia. *Phycological Research* 57(3):159-169.
- Neustupa J, Škaloud P. 2010. Diversity of subaerial algae and cyanobacteria growing on bark and wood in the lowland tropical forests of Singapore. *Plant Ecology and Evolution* 143(1):51-62.

- Neustupa J, Eliáš M, Škaloud P, Němcová Y, Šejnohová L. 2011. *Xylochloris irregularis* gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a novel subaerial coccoid green alga. *Phycologia* 50(1):57-66.
- Neustupa J, Němcová Y, Veselá J, Steinová J, Škaloud P. 2013a. *Leptochlorella corticola* gen. et sp. nov. and *Kalinella apyrenoidosa* sp. nov.: two novel *Chlorella*-like green microalgae (Trebouxiophyceae, Chlorophyta) from subaerial habitats. *International Journal of Systematic and Evolutionary Microbiology* 63:377-387.
- Neustupa J, Němcová Y, Veselá J, Steinová J, Škaloud P. 2013b. *Parachloroidium* gen. nov. (Trebouxiophyceae, Chlorophyta), a novel genus of coccoid green algae from subaerial corticolous biofilms. *Phycologia* 52(5):411-421.
- Nikorych V, Chornevych T. 2012. Ecobiomorph structure of algocenoses organization of brownish-podzolic gleied soils of Ukrainian Ciscarpathia. *Soil Forming Factors and Processes from the Temperate Zone* 10:141-151.
- Novakovskaya IV, Patova EN. 2007. Changes in soil algal communities in spruce phytocenoses under the influence of aerotechnogenic pollution. *Eurasian Soil Science* 40(5):576-582.
- Novakovskaya IV, Patova EN. 2008. Green algae in spruce forests in the north-east of European Russia. *Biologia* 63(6):836-842.
- Novichkova-Ivanova LN. 1972. Soil algae of middle Asia deserts. In: Rodin LE, editor. *Eco-physiology foundation ecosystems productivity in arid zones*. Nauka. p. 180-182.
- Novis PM, Beer T, Vallance J. 2008. New records of microalgae from the New Zealand alpine zone, and their distribution and dispersal. *New Zealand Journal of Botany* 46(3):347-366.
- Novis PM, Aislabie J, Turner S, McLeod M. 2015. Chlorophyta, Xanthophyceae and Cyanobacteria in Wright Valley, Antarctica. *Antarctic Science* 27(5):439-454.
- Olivieri G, Marzocchella A, Andreozzi R, Pinto G, Pollio A. 2011. Biodiesel production from *Stichococcus* strains at laboratory scale. *Journal of Chemical Technology and Biotechnology* 86(6):776-783.
- Olivieri G, Gargano I, Andreozzi R, Marotta R, Marzocchella A, Pinto G, Pollio A. 2013. Effects of photobioreactors design and operating conditions on *Stichococcus bacillaris* biomass and biodiesel production. *Biochemical Engineering Journal* 74:8-14.
- Osundeko O, Davies H, Pittman JK. 2013. Oxidative stress-tolerant microalgae strains are highly efficient for biofuel feedstock production on wastewater. *Biomass Bioenergy* 56:284-294.
- Palacios C, Zettler E, Amils R, Amaral-Zettler L. 2008. Contrasting microbial community assembly hypotheses: a reconciling tale from the Rio Tinto. *PLoS ONE* 3(12):e3853.
- Park KC, Whitney C, Kozera C, O'Leary SJ, McGinn PJ. 2015. Seasonal isolation of microalgae from municipal wastewater for remediation and biofuel applications. *Journal of Applied Microbiology* 119:76-87.
- Patova EN, Dorokhova MF. 2008. Green algae in tundra soils affected by coal mine pollutions. *Biologia* 63(6):831-835.
- Patterson D. 1989. Stramenopiles: chromophytes from a protistan perspective. In: Green J, Leadbeater B, Diver W, editors. *The chromophyte algae: problems and perspectives*. Oxford Science Publications. p. 357-379.
- Paulsen BS, Vieira AAH, Klaveness D. 1992. Structure of extracellular polysaccharides produced by a soil *Cryptomonas* sp. (Cryptophyceae). *Journal of Phycology* 28(1):61-63.
- Peer T, Türk R, Gruber J, A T. 2010. Species composition and pedological characteristics of biological soil crusts in a high alpine ecosystem, Hohe Tauern, Austria. *Eco.mont* 2(1):23-30.

- Pentecost A. 1988. Observations on growth rates and calcium carbonate deposition in the green alga *Gongrosira*. *New Phytologist* 110(2):249-253.
- Pentecost A, Spiro B. 1990. Stable carbon and oxygen isotope composition of calcites associated with modern freshwater cyanobacteria and algae. *Geomicrobiology Journal* 8(1):17-26.
- Perůtková K. 2014. Diverzita terestrických řas urbánních habitatů ve střední Evropě. MSc thesis, Masaryk University (Brno). 70 pp.
- Petersen JB. 1935. Studies on the biology and taxonomy of soil algae. *Dansk Botanisk Arkiv* 8. 183 pp.
- Pia J. 1926. *Pflanzen als Gesteinsbildner*. Berlin: Verlag Gebrüder Bornträger. 355 pp.
- Pimm SL, Raven P. 2000. Biodiversity: extinction by numbers. *Nature* 403(6772):843-845.
- Posada D. 2008. jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* 25(7):1253-1256.
- Posrednikova AV, Pakhomov OY, Kulbachko YL. 2009. Soil algae and mesofauna communities in biotopes of forest rehabilitation in Zhovti Vody (Dnipropetrovsk region). *Visnyk of Dnipropetrovsk University, Biology, Ecology* 17(1):172-177.
- Preston-Mafham J, Boddy L, Randerson PF. 2002. Analysis of microbial community functional diversity using sole-carbon-source utilisation profiles – a critique. *FEMS Microbiology Ecology* 42(1):1-14.
- Prober SM, Leff JW, Bates ST, Borer ET, Firn J, Harpole WS, Lind EM, Seabloom EW, Adler PB, Bakker JD. 2015. Plant diversity predicts beta but not alpha diversity of soil microbes across grasslands worldwide. *Ecology Letters* 18(1):85-95.
- Pröschold T, Marin B, Gert U, Melkonian M. 2001. Molecular phylogeny and taxonomic revision of *Chlamydomonas* (Chlorophyta). I. Emendation of *Chlamydomonas* Ehrenberg and *Chloromonas* Gobi, and description of *Oogamochlamys* gen. nov. and *Lobochlamys* gen. nov. *Protist* 152:265-300.
- Pröschold T, Bock C, Luo W, Krienitz L. 2010. Polyphyletic distribution of bristle formation in Chlorellaceae: *Micractinium*, *Diacanthos*, *Didymogenes* and *Hegewaldia* gen. nov. (Trebouxiophyceae, Chlorophyta). *Phycological Research* 58(1):doi:10.1111/j.1440-1835.2009.00552.x.
- Prosser JI. 2002. Molecular and functional diversity in soil micro-organisms. *Plant and Soil* 244(1-2):9-17.
- Ragon M, Fontaine MC, Moreira D, López-García P. 2012. Different biogeographic patterns of prokaryotes and microbial eukaryotes in epilithic biofilms. *Molecular Ecology* 21(15):3852-3868.
- Rahmann S, Müller T, Dandekar T, Wolf M. 2006. Efficient and robust analysis of large phylogenetic datasets. In: Hsu H-H, editor. *Advanced Data Mining Technologies in Bioinformatics*. Hershey, PA, USA: Idea Group, Inc. p. 104-117.
- Rahmonov O, Piątek J. 2007. Sand colonization and initiation of soil development by Cyanobacteria and algae. *Ekológia* 26(1):52-63.
- Rambaut A. FigTree, a graphical viewer of phylogenetic trees. <http://tree.bio.ed.ac.uk/software/figtree>.
- Rastogi G, Sani RK. 2011. Molecular techniques to assess microbial community structure, function, and dynamics in the environment. *Microbes and Microbial Technology*. Springer. p. 29-57.
- Raven JA. 2012. Algal biogeography: metagenomics shows distribution of a picoplanktonic pelagophyte. *Current Biology* 22(17):R682-R683.

- Ray JG, Thomas BT. 2012. Fertility characteristics of oxic dystropepts under natural forest, rubber, and teak plantations in different seasons, Kerala, South India. *Communications in Soil Science and Plant Analysis* 43(17):2247-2261.
- Reisigl H. 1958. Über die Verteilung der Bodenalgae in der Gipfelstufe der Ötztaler Alpen. *Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck* 53:163-172.
- Reisigl H. 1964. Zur Systematik und Ökologie alpiner Bodenalgae. *Österreichische botanische Zeitschrift* 111 (4): 402-499.
- Reuter JS, Mathews DH. 2010. RNAstructure: software for RNA secondary structure prediction and analysis. *BMC Bioinformatics* 11:doi:10.1186/1471-2105-11-129.
- Reynaud PA, Lumpkin TA. 1988. Microalgae of the Lanzhou (China) cryptogamic crust. *Arid Land Research and Management* 2(3):145-155.
- Řezáčová M, Neustupa J. 2007. Distribution of the genus *Mallomonas* (Synurophyceae) - ubiquitous dispersal in microorganisms evaluated. *Protist* 158(1):29-37.
- Řezanka T, Temina M, Tolstikov A, Dembitsky V. 2004. Natural microbial UV radiation filters—mycosporine-like amino acids. *Folia Microbiologica* 49(4):339-352.
- Richards TA, Bass D. 2005. Molecular screening of free-living microbial eukaryotes: diversity and distribution using a meta-analysis. *Current Opinion in Microbiology* 8(3):240-252.
- Richards TA, Vepritskiy AA, Gouliamova DE, Nierzwicki-Bauer SA. 2005. The molecular diversity of freshwater picoeukaryotes from an oligotrophic lake reveals diverse, distinctive and globally dispersed lineages. *Environmental Microbiology* 7(9):1413-1425.
- Riding R. 1991. Cambrian calcareous cyanobacteria and algae. In: Riding R, editor. *Calcareous algae and stromatolites*. Berlin: Springer-Verlag. 571 pp.
- Řídká T, Peksa O, Rai H, Upreti DK, Škaloud P. 2014. Photobiont diversity in Indian *Cladonia* lichens, with special emphasis on the geographical patterns. In: Rai H, Upreti DK, editors. *Terricolous lichens in India*. Springer. p. 53-71.
- Rindi F, McIvor L, Guiry GM. 2004. The Prasiolales (Chlorophyta) of Atlantic Europe: An assessment based on morphological, molecular, and ecological data, including the characterization of *Rosenvingiella radicans* (Kützinger) comb. nov. *Journal of Phycology* 40(5):977-997.
- Rindi F, McIvor L, Sherwood A, Friedl T, Guiry M, Sheath R. 2007. Molecular phylogeny of the green algal order Prasiolales (Trebouxiophyceae, Chlorophyta). *Journal of Phycology* 43:811-822.
- Rindi F, Guiry MD, Lo JM. 2008. Distribution, morphology, and phylogeny of *Klebsormidium* (Klebsormidiales, Charophyceae) in urban environments in Europe. *Journal of Phycology* 44:1529-1540.
- Rindi F, Allali HA, Lam DW, López-Bautista JM. 2009. An overview of the biodiversity and biogeography of terrestrial green algae. In: Rescigno V, editor. *Biodiversity Hotspots*. Nova Science Publishers. 25 pp.
- Rindi F, Hodkinson T, Jones M, Waldren S, Parnell J. 2011. Terrestrial green algae: systematics, biogeography and expected responses to climate change. In: Hodkinson TR, editor. *Climate Change, Ecology and Systematics*. Cambridge University Press. p. 201-227.
- Rodó JDC-P, Molinari-Novoa EA. 2015. A nomenclatural note and cultural note on *Chlorella peruviana* G. Chacón and other species of the genus *Chlorella* Beij. (Chlorellaceae, Chlorellales). *The Biologist (Lima)* 13(1):71-74.
- Roger P-A, Reynaud P-A. 1982. Free-living blue-green algae in tropical soils. In: Dommergues Y, Diem H, editors. *Microbiology of tropical soils and plant productivity*. La Hague: Martinus Nijhoff Publisher. p. 147-168.

- Rondon MR, August PR, Bettermann AD, Brady SF, Grossman TH, Liles MR, Loiacono KA, Lynch BA, MacNeil IA, Minor C. 2000. Cloning the soil metagenome: a strategy for accessing the genetic and functional diversity of uncultured microorganisms. *Applied and Environmental Microbiology* 66(6):2541-2547.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Hohna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3):539-542.
- Rosa K, Lhotský O. 1971. Edaphische Algen und Protozoen im Isergebirge, Tschechoslowakei. *Oikos* 22:21-29.
- Rybalka N, Andersen RA, Kostikov I, Mohr KI, Massalski A, Olech M, Friedl T. 2009. Testing for endemism, genotypic diversity and species concepts in Antarctic terrestrial microalgae of the Tribonemataceae (Stramenopiles, Xanthophyceae). *Environmental Microbiology* 11(3):554-565.
- Rybalka N, Wolf M, Andersen RA, Friedl T. 2013. Congruence of chloroplast- and nuclear-encoded DNA sequence variations used to assess species boundaries in the soil microalga *Heterococcus* (Stramenopiles, Xanthophyceae). *BMC Evolutionary Biology* 13(39): doi:10.1186/1471-2148-13-39.
- Ryšánek D, Hrčková K, Škaloud P. 2014. Global ubiquity and local endemism of free-living terrestrial protists: phylogeographic assessment of the streptophyte alga *Klebsormidium*. *Environmental Microbiology* 17(3):689-698.
- Safi C, Zebib B, Merah O, Pontalier P-Y, Vaca-Garcia C. 2014. Morphology, composition, production, processing and applications of *Chlorella vulgaris*: A review. *Renewable & Sustainable Energy Reviews* 35:265-278.
- Şalaru V, Trofim A, Melniciuc C, Dontu N. 2008. Taxonomic and ecologic structure of communities of edaphic algae from the agrophytocenoses of the northern districts of Moldova. *Journal of Plant Development* 15:3-6.
- Santoferrara LF, Grattepanche JD, Katz LA, McManus GB. 2014. Pyrosequencing for assessing diversity of eukaryotic microbes: analysis of data on marine planktonic ciliates and comparison with traditional methods. *Environmental Microbiology* 16(9):2752-2763.
- Scalzi G, Selbmann L, Zuconi L, Rabbow E, Horneck G, Albertano P, Onofri S. 2012. LIFE Experiment: isolation of cryptoendolithic organisms from Antarctic colonized sandstone exposed to space and simulated Mars conditions on the International Space Station. *Origins of Life and Evolution of Biospheres* 42:253-262.
- Scherbina V, Maltseva I, Solonenko A. 2014. Peculiarities of postpyroge development of algae in steppe biocenoses at Askania Nova Biospheric National Park. *Contemporary Problems of Ecology* 7(2):187-191.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ. 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* 75(23):7537-7541.
- Schmidle W. 1901. Ueber drei Algengenera. *Berichte der Deutschen Botanischen Gesellschaft* 19:149-158.
- Schmidt SK, Lynch RC, King AJ, Karki D, Robeson MS, Nagy L, Williams MW, Mitter MS, Freeman KR, Schmidt SK, Lynch RC, King AJ, Karki D. 2011. Phylogeography of microbial phototrophs in the dry valleys of the high Himalayas and Antarctica. *Proceedings of the Royal Society B: Biological Sciences* 278:702-708.

- Schmidt SK, Darcy JL. 2014. Phylogeny of ulotrichalean algae from extreme high-altitude and high-latitude ecosystems. *Polar Biology*:doi:10.1007/s00300-014-1631-6.
- Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9(7):671-675.
- Schultz J, Müller T, Achtziger M, Seibel PN, Dandekar T, Wolf M. 2006. The internal transcribed spacer 2 database - a web server for (not only) low level phylogenetic analyses. *Nucleic Acids Research* 34:704-707.
- Sciuto K, Lewis LA, Verleyen E, Moro I, La Rocca N. 2015. *Chodatodesmus australis* sp. nov. (Scenedesmaceae, Chlorophyta) from Antarctica, with the emended description of the genus *Chodatodesmus*, and circumscription of *Flechtneria rotunda* gen. et sp. nov. *Journal of Phycology*:doi:10.1111/jpy.12355-14-227.
- Seibel PN, Muller T, Dandekar T, Schultz J, Wolf M. 2006. 4SALE - a tool for synchronous RNA sequence and secondary structure alignment and editing. *BMC Bioinformatics* 7(498):doi:10.1186/1471-2105-7-498.
- Seibel PN, Muller T, Dandekar T, Wolf M. 2008. Synchronous visual analysis and editing of RNA sequence and secondary structure alignments using 4SALE. *BMC Research Notes* 1(91):doi:10.1186/1756-0500-1-91.
- Selig C, Wolf M, Müller T, Dandekar T, Schultz J. 2008. The ITS2 Database II: homology modelling RNA structure for molecular systematics. *Nucleic Acids Research* 36:377-380.
- Shade A, Hogan CS, Klimowicz AK, Linske M, McManus PS, Handelsman J. 2012. Culturing captures members of the soil rare biosphere. *Environmental Microbiology* 14(9):2247-2252.
- Sharma NK, Rai AK, Singh S, Brown RMJ. 2007. Airborne algae: their present status and relevance. *Journal of Phycology* 43(4):615-627.
- Sharma NK, Rai AK. 2010. Biodiversity and biogeography of microalgae: progress and pitfalls. *Environmental Reviews* 19:doi:10.1139/A10-020.
- Shekhovtseva OG. 2014. Soil algae in urban ecosystems of the city of Mariupol. *Gruntoznavstvo* 15(1-2):88-92.
- Shi XL, Marie D, Jardillier L, Scanlan DJ, Vaultot D. 2009. Groups without cultured representatives dominate eukaryotic picophytoplankton in the oligotrophic South East Pacific Ocean. *PLoS ONE* 4(10):e7657.
- Shields LM, Durrell LW. 1964. Algae in relation to soil fertility. *The Botanical Review* 30(1):92-128.
- Shiraishi F, Bissett A, de Beer D, Reimer A, Arp G. 2008a. Photosynthesis, respiration and exopolymer calcium-binding in biofilm calcification (Westerhöfer and Deinschwanger Creek, Germany). *Geomicrobiology Journal* 25(2):83-94.
- Shiraishi F, Reimer A, Bissett A, de Beer D, Arp G. 2008b. Microbial effects on biofilm calcification, ambient water chemistry and stable isotope records in a highly supersaturated setting (Westerhöfer Bach, Germany). *Palaeogeography, Palaeoclimatology, Palaeoecology* 262(1):91-106.
- Shtina EA, Gollerbach MM. 1976. Ecology of soil algae. Nauka. 143 pp.
- Shtina EA. 1990. Soil algae as ecological indicators. *Botanical Journal*(4):441-452.
- Shukla SP, Kvíderová J, Elster J. 2011. Nutrient requirements of polar *Chlorella*-like species. *Czech Polar Reports* 1(1):1-10.
- Sivakumar G, Jeong K, Lay JO. 2014. Bioprocessing of *Stichococcus bacillaris* strain siva2011. *Biotechnology for Biofuels* 7(62):doi:10.1186/1754-6834-7-62.

- Škaloud P, Jiří N, Radochová B, Kubínová L. 2005. Confocal microscopy of chloroplast morphology and ontogeny in three strains of *Dictyochloropsis* (Trebouxiophyceae, Chlorophyta). *Phycologia* 44(3):261-269.
- Škaloud P. 2009. Species composition and diversity of aero-terrestrial algae and cyanobacteria of the Boreč Hill ventaroles. *Fottea* 9(1):65–80.
- Škaloud P, Rindi F. 2013. Ecological differentiation of cryptic species within an asexual protist morphospecies: a case study of filamentous green alga *Klebsormidium* (Streptophyta). *Journal of Eukaryotic Microbiology* 60(4):350-362.
- Škaloud P, Lukešová A, Malavasi V, Ryšánek D, Hřčková K, Rindi F. 2014a. Molecular evidence for the polyphyletic origin of low pH adaptation in the genus *Klebsormidium* (Klebsormidiophyceae, Streptophyta). *Plant Ecology and Evolution* 147(3):333-345.
- Škaloud P, Škaloudová M, Procházková A, Němcová Y. 2014b. Morphological delineation and distribution patterns of four newly described species within the *Synura petersenii* species complex (Chrysophyceae, Stramenopiles). *European Journal of Phycology* 49(2):213-229.
- Škaloud P, Steinová J, Řídká T, Vančurová L, Peksa O. 2015. Assembling the challenging puzzle of algal biodiversity: species delimitation within the genus *Asterochloris* (Trebouxiophyceae, Chlorophyta). *Journal of Phycology* 51:507–527.
- Šlapeta J, Moreira D, López-García P. 2005. The extent of protist diversity: insights from molecular ecology of freshwater eukaryotes. *Proceedings of the Royal Society B: Biological Sciences* 272(1576):2073-2081.
- Šlapeta J, López-García P, Moreira D. 2006. Global dispersal and ancient cryptic species in the smallest marine eukaryotes. *Molecular Biology and Evolution* 23(1):23-29.
- Slocombe SP, Zhang Q, Ross M, Anderson A, Thomas NJ, Lapresa Á, Rad-Menéndez C, Campbell CN, Black KD, Stanley MS, Day JG. 2015. Unlocking nature's treasure-chest: screening for oleaginous algae. *Scientific Reports* 5:doi:10.1038/srep09844.
- Sluiman HJ, Guihal C. 2008. Assessing phylogenetic affinities and species delimitations in *Klebsormidiales* (Streptophyta): Nuclear-encoded rDNA phylogenies and its secondary structure models in *Klebsormidium*, *Hormidiella* and *Entransia*. *Journal of Phycology* 44:183-195.
- Smith H, Hardy P, Leith I, Spaul V, Twelves E. 1974. A biological survey of St. Paul's Rocks in the equatorial Atlantic Ocean. *Biological Journal of the Linnean Society* 6(2):89-96.
- Smith S, Abed R, Garcia-Pichel F. 2004. Biological soil crusts of sand dunes in Cape Cod National Seashore, Massachusetts, USA. *Microbial Ecology* 48(2):200-208.
- Soare LC, Dobrescu CM. 2010. Preliminary data on edaphic algae in the city of Pitești (Romania). *Analele Universității din Oradea - Fascicula Biologie*. p. 186-189.
- Song H, Zhang Q, Liu G, Hu Z. 2015. *Polulichloris henanensis* gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a novel subaerial coccoid green alga. *Phytotaxa* 218(2):137-146.
- Spitzer K. 2013. Biodiversity of terrestrial green algae in tropical mountain rain forest in Podocarpus National Park (Ecuador) using a culture dependent approach. MSc thesis, University of Duisburg-Essen. 122 pp.
- Spitzer K, Faßhauer F, Hodač L, Friedl T. 2014. Biodiversity of terrestrial green algae in a tropical mountain rain forest of Ecuador (Podocarpus National Park). Poster. Book of Abstracts of the 44th Annual Meeting of the Ecological Society of Germany, Austria and Switzerland. Hildesheim (Germany). p. 240.

- Stamatakis A, Hoover P, Rougemont J. 2008. A rapid bootstrap algorithm for the RAxML Web servers. *Systematic Biology* 57(5):758-771.
- Stanek-Tarkowska J, Noga T. 2012. Diversity of diatoms (Bacillariophyceae) in the soil under traditional tillage and reduced tillage. *Inżynieria Ekologiczna* 30:287-296.
- Starks TL, Shubert LE, Trainor FR. 1981. Ecology of soil algae: a review. *Phycologia* 20:65-80.
- Stibal M, Šabacká M, Kaštovská K. 2006. Microbial communities on glacier surfaces in Svalbard: impact of physical and chemical properties on abundance and structure of cyanobacteria and algae. *Microbial Ecology* 52(4):644-654.
- Strimmer K, Von Haeseler A. 1996. Quartet puzzling: a quartet maximum-likelihood method for reconstructing tree topologies. *Molecular Biology and Evolution* 13(7):964-969.
- Sumbali G, Mehrotra R. 2009. *Principles of Microbiology*. Tata McGraw-Hill. 926 pp.
- Sun Z, Li G, Wang C, Jing Y, Zhu Y, Zhang S, Liu Y. 2014. Community dynamics of prokaryotic and eukaryotic microbes in an estuary reservoir. *Scientific Reports* 4(6966):doi:10.1038/srep06966.
- Suutari M, Majaneva M, Fewer D, Voirin B, Aiello A, Friedl T, Chiarello A, Blomster J. 2010. Molecular evidence for a diverse green algal community growing in the hair of sloths and a specific association with *Trichophilus welckeri* (Chlorophyta, Ulvophyceae). *BMC Evolutionary Biology* 10(1):86.
- Taib N, Mangot J-F, Domaizon I, Bronner G, Debroas D. 2013. Phylogenetic affiliation of SSU rRNA genes generated by massively parallel sequencing: new insights into the freshwater protist diversity. *PLoS ONE* 8(3):e58950.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28(10):2731-2739.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30(12):2725-2729.
- Tell G. 1976. Algas del suelo de los alrededores de Buenos Aires (República Argentina). *Darwiniana* 20(3-4):491-548.
- Temraleeva A, Pinskii D, Patova E, Spirina E. 2011. The use of algae-cyanobacterial communities for the assessment of lead pollution of gray forest soils. *Eurasian Soil Science* 44(3):326-331.
- ter Braak CJF, Šmilauer P. 2002. CANOCO reference manual and CanoDraw for Windows user's guide: software for canonical community ordination (version 4.5). Microcomputer Power. 500 pp.
- Terlizzi DE, Karlander EP. 1979. Soil algae from a Maryland serpentine formation. *Soil Biology and Biochemistry* 11(2):205-207.
- Thüs H, Muggia L, Pérez-Ortega S, Favero-Longo SE, Joneson S, O'Brien H, Nelsen MP, Duque-Thüs R, Grube M, Friedl T. 2011. Revisiting photobiont diversity in the lichen family Verrucariaceae (Ascomycota). *European Journal of Phycology* 46(4):399-415.
- Trainor FR. 1970. Survival of algae in a desiccated soil. *Phycologia* 9(2):111-113.
- Trainor FR. 1985. Survival of algae in a desiccated soil: a 25 year study. *Phycologia* 24(1):79-82.
- Trainor FR, Gladych R. 1995. Survival of algae in a desiccated soil: a 35-year study. *Phycologia* 34(3):191-192.
- Trenkwalder H. 1975. Neue Bodenalgae aus Föhrenwäldern im Raum von Brixen (Südtirol, Italien). *Berichte des naturwissenschaftlich-medizinischen Vereins in Innsbruck* 62:7-19.

- Treves H, Raanan H, Finkel OM, Berkowicz SM, Keren N, Shotland Y, Kaplan A. 2013. A newly isolated *Chlorella* sp. from desert sand crusts exhibits a unique resistance to excess light intensity. *FEMS Microbiology Ecology* 86(3):373-380.
- Trześcińska M, Pawlik-Skowrońska B. 2008. Soil algal communities inhabiting zinc and lead mine spoils. *Journal of Applied Phycology* 20(4):341-348.
- Tschermak-Woess E. 1988. New and known taxa of *Chlorella* (Chlorophyceae): Occurrence as lichen phycobionts and observations on living dictyosomes. *Plant Systematics and Evolution* 159(1-2):123-139.
- Türk R, Gärtner G. 2001. Biological soil crusts of the subalpine, alpine and nival areas in the Alps. *Ecological Studies* 150:67-73.
- Turmel M, Otis C, Lemieux C. 2015. Dynamic evolution of the chloroplast genome in the Green algal classes Pedinophyceae and Trebouxiophyceae. *Genome Biology and Evolution* 7:2062–2082.
- Veselá J. 2006. Benthic algal communities and their ecology in sandstone periodically desiccated brook in National Park Bohemian Switzerland (Czech Republic). *Czech Phycology* 6:99-110.
- Veselá J, Urbánková P, Cerná K, Neustupa J. 2012. Ecological variation within traditional diatom morphospecies: diversity of *Frustulia rhomboides* sensu lato (Bacillariophyceae) in European freshwater habitats. *Phycologia* 51(5):552-561.
- Viles HA. 2012. Microbial geomorphology: a neglected link between life and landscape. *Geomorphology* 157–158:6-16.
- Vinogradova OM, Darienko TM. 2008. Terrestrial algae of hypersaline environments of the Central Syvash islands (Kherson Region, Ukraine). *Biologia* 63(6):813-823.
- Viprey M, Guillou L, Ferréol M, Vaultot D. 2008. Wide genetic diversity of picoplanktonic green algae (Chloroplastida) in the Mediterranean Sea uncovered by a phylum-biased PCR approach. *Environmental Microbiology* 10(7):1804-1822.
- Vishnivetskaya TA. 2009. Viable Cyanobacteria and Green algae from the permafrost darkness. In: Margesin R, editor. *Permafrost Soils*. Berlin, Heidelberg: Springer-Verlag. p. 73-84.
- Vyverman W, Verleyen E, Wilmotte A, Hodgson DA, Willems A, Peeters K, Van de Vijver B, De Wever A, Leliaert F, Sabbe K. 2010. Evidence for widespread endemism among Antarctic micro-organisms. *Polar Science* 4:103-113.
- Wallner J. 1934. Über die Beteiligung kalkablagernder Pflanzen bei der Bildung südbayerischer Tuffe. *Bibliotheca Botanica* 110:1–30.
- Watanabe S. 1983. New and Interesting Green Algae from Soils of some Asian and Oceanian Regions. *Archiv für Protistenkunde* 127(3):223-270.
- Watanabe S, Nakayama T. 2007. Ultrastructure and phylogenetic relationships of the unicellular green algae *Ignatius tetrasporus* and *Pseudocharacium americanum* (Chlorophyta). *Phycological Research* 55:doi:10.1111/j.1440-1835.2006.00439.x.
- Wheeler DL, Barrett T, Benson DA, Bryant SH, Canese K, Chetvernin V, Church DM, DiCuccio M, Edgar R, Federhen S. 2007. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Research* 35:D5-D12.
- Winsborough BM, Golubić S. 1987. The role of diatoms in stromatolite growth: two examples from modern freshwater settings. *Journal of Phycology* 23(2):195-201.
- Wojtuń B, Samecka-Cymerman A, Kolon K, Kempers A, Skrzypek G. 2013. Metals in some dominant vascular plants, mosses, lichens, algae, and the biological soil crust in various types of terrestrial tundra, SW Spitsbergen, Norway. *Polar Biology* 36(12):1799-1809.

- Wolf M, Achtziger M, Schultz J, Dandekar T, Müller T. 2005. Homology modeling revealed more than 20,000 rRNA internal transcribed spacer 2 (ITS2) secondary structures. *RNA* 11:1616-1623.
- Wolf M, Ruderisch B, Dandekar T, Schultz J, Müller T. 2008. ProfDistS: (profile-) distance based phylogeny on sequence - structure alignments. *Bioinformatics* 24(20):2401-2402.
- Wolf M, Chen S, Song J, Ankenbrand M, Muller T. 2013. Compensatory base changes in ITS2 secondary structures correlate with the biological species concept despite intragenomic variability in ITS2 sequences-a proof of concept. *PLoS ONE* 8(6):e66726.
- Wong FKY, Lacap DC. 2010. Hypolithic microbial community of quartz pavement in the high-altitude tundra of Central Tibet. *Environmental Microbiology* 60:730-739.
- Xiao X, Sogge H, Lagesen K, Tooming-Klunderud A, Jakobsen KS, Rohrlack T. 2014. Use of high throughput sequencing and light microscopy show contrasting results in a study of phytoplankton occurrence in a freshwater environment. *PLoS ONE* 9(8):e106510.
- Yamamoto M, Nishikawa T, Kajitani H, Kawano S. 2007. Patterns of asexual reproduction in *Nannochloris bacillaris* and *Marvania geminata* (Chlorophyta, Trebouxiophyceae). *Planta* 226(4):917-927.
- Zancan S, Trevisan R, Paoletti MG. 2006. Soil algae composition under different agro-ecosystems in North-Eastern Italy. *Agriculture, Ecosystems and Environment* 112:1-12.
- Zenova G, Shtina E. 1990. Soil Algae: Tutorial. Moscow State University. 80 pp.
- Zenova G, Shtina E, Dedysh S, Glagoleva O, Likhacheva A, Gracheva T. 1995. Ecological relations of algae in biocenoses. *Microbiology* 64(2):121-133.
- Zhan A, Hulak M, Sylvester F, Huang X, Adebayo AA, Abbott CL, Adamowicz SJ, Heath DD, Cristescu ME, MacIsaac HJ. 2013. High sensitivity of 454 pyrosequencing for detection of rare species in aquatic communities. *Methods in Ecology and Evolution* 4(6):558-565.
- Zhang J, Huss VA, Sun X, Chang K, Pang D. 2008. Morphology and phylogenetic position of a trebouxiophycean green alga (Chlorophyta) growing on the rubber tree, *Hevea brasiliensis*, with the description of a new genus and species. *European Journal of Phycology* 43(2):185-193.
- Zheng Y, Li T, Yu X, Bates PD, Dong T, Chen S. 2013. High-density fed-batch culture of a thermotolerant microalga *Chlorella sorokiniana* for biofuel production. *Applied Energy* 108:281-287.
- Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Research* 31(13):3406-3415.

Appendix | Chapter 1

Supporting Figures

Figure S1. Microphotographs of mixed cultures (Archaeplastida).

Figure S2. Microphotographs of mixed cultures (Archaeplastida, Stramenopiles, Cryptophyta).

Figure S3. Diatoms in mixed cultures.

Figure S4. Cyanobacteria in mixed cultures.

Figure S5. Diversity in mixed cultures as inferred by light-microscopy.

Figure S6. Geographic evidence of 25 most common Green algal morphotypes.

Supporting Tables

Table S1. List of all analyzed isolates.

Table S2a. List of all detected species belonging to Trebouxiophyceae.

Table S2b. List of all detected species belonging to Chlorophyceae, other green algae and Stramenopiles.

Table S3. List of the closest GenBank-relatives of our detected species.

Table S4a. List of all analyzed full and partial 18S rDNA sequences.

Table S4b. Distribution of the green algal species across the sampling sites.

Table S5. General morphological characteristics of the new isolates.

Table S6a. Diversity of morphospecies in soil drill cores.

Table S6b. Diversity of morphospecies in topsoil samples.

Table S7. Frequency of the morphospecies in soil drill cores.




Figure S1. Microphotographs of mixed cultures (Archaeplastida). (a) Characiochloridaceae, *Chlamydropodium* (HEG2); (b) Characiochloridaceae, *Chlamydropodium* (HEG9); (c) cf. *Tetracystis* (HEG9); (d) Actinochloridaceae (HEW1); (e) Actinochloridaceae (HEW1); (f) Chlorosarcinaceae, cf. *Desmotetra* (HEG6); (g) Scenedesmaceae, *Scenedesmus* cf. *soil* (HEG7); (h) Selenastraceae, *Monoraphidium terrestre* (HEG9); (i) cf. *Keratococcus* (HEG7); (j) *Podohedra* (HEG7); (k) cf. *Geminella* (HEW3); (l) *Scotinosphaera* (HEG9). Scale bars = 10 μ m.




Figure S2. Microphotographs of mixed cultures (Archaeplastida, Stramenopiles, Cryptophyta). (a) cf. *Pseudendoclonium* (HEG7); (b) cf. *Dilabifilum* (SEG8); (c) cf. *Chaetophora* (HEG9); (d) *Cosmarium* (HEG9); (e) *Cosmarium* (HEG9); (f) cf. *Klebsormidium* (HEG9); (g) *Interfilum* cf. *terricola* (HEW); (h) Rhodophyta, *Porphyridium* (HEG9); (i) *Cryptomonas* (HEG9); (j) *Tribonema* (HEG9); (k) cf. *Ellipsoidion* (HEG6); (l) cf. *Monallantus* (HEW4). Scale bars = 10 μ m.




Figure S3. Diatoms in mixed cultures. (a) Monoraphid diatoms, cf. *Planothidium* (HEG9); (b)-(k) Symmetrical biraphid diatoms; (b) *Humidophila* (HEG6); (c) *Luticola* (HEG9); (d) cf. *Eolimna* (HEW3); (e) cf. *Mayamaea* (HEW3); (f) *Pinnularia* (HEW6); (g) *Pinnularia* (HEW9); (h) *Pinnularia* (HEW); (i) *Pinnularia* (HEG8); (j) *Stauroneis* (HEG9); (k) cf. *Neidium* (HEW4); (l) Surirelloid diatoms, *Surirella* (HEW); (m)-(o) Nitzschoid diatoms; (m) *Nitzschia* (HEG4); (n) *Nitzschia* (HEG9); (o) *Hantzschia* (HEG6). Scale bars = 5 μ m.




Figure S4. Cyanobacteria in mixed cultures. (a) cf. *Pseudanabaena* (HEG9); (b) cf. *Leptolyngbya* (HEW4); (c) cf. *Leptolyngbya* (HEG9); (d) cf. *Limnothrix* (HEG6); (e) *Pseudanabaenales* (HEG7); (f) *Phormidium* (HEG7); (g) *Nostoc* (HEG7); (h) *Nostoc* (HEG7); (i) *Nostoc* (HEG7); (j) *Cylindrospermum* (HEG6); (k) cf. *Tolypothrix* (HEG2); (l) cf. *Stigonema* (HEG9). Scale bars = 10 μ m.




Figure S5. Diversity in mixed cultures as inferred by light-microscopy. (a) Localization of the three German Biodiversity Exploratories. (b) Comparison of alpha-diversities (= numbers of morphotypes) of soil algae in grasslands and forests. Each bar represents pooled data from nine to twelve plots (e.g., HEG(1-9) includes observations from all Hainich grassland plots); (c) Comparison of alpha-diversities among the land-uses (ext=extensive, man=managed, int=intensive). Each bar includes pooled data from multiple plots (the number is given in parentheses). (d) Redundancy analysis (RDA) of algal communities changing along environmental gradients (pH, C_{org}=organic carbon, N_{tot}=total nitrogen). Dots (forests) and squares (grasslands) represent algal communities which were pooled within each land-use category (e.g., ALBe includes observations from plots AEW7-AEW9; e=extensively used plots). A subset of 22 most frequently observed morphotypes are shown (black vectors). (e) Non-metric multidimensional scaling (NMDS) of algal communities in multiple subsamples within a plot. Each dot (forest subsample) and square (grassland subsample) represents data recorded during three seasons. Within each plot (HEG8, HEG9, HEW5, HEW6) three subsamples were analyzed (NE=north-east plot edge, MID=middle point, SW=south-west plot edge, MIX=pooled sample combining all three subsamples within a plot; sampling scheme is shown above).




Table S1. List of all analyzed isolates.

| Species | Identifier | GPS | Exploratory | Plot | Habitat | Sampling | 18S | ITS2 |
|--------------------------------------|---------------------------|--------------------------------------|-------------|------|-----------|--------------|------|------|
| <i>Acutodesmus rubescens</i> | LH08SG8041 | N53° 6' 50.294" E14° 1' 1.559" | SCH | SEG8 | grassland | drill core | 1656 | - |
| <i>Bracteacoccus cohaerens</i> | LH10HG9034 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1722 | - |
| <i>Bracteacoccus cohaerens</i> (cf.) | LH08SG2015 | N53° 5' 21.505" E13° 58' 48.169" | SCH | SEG2 | grassland | drill core | 2056 | - |
| <i>Chlamydomonas gerloffii</i> (cf.) | LH08SW5031 | N53° 3' 25.321" E13° 53' 7.318" | SCH | SEW5 | forest | drill core | 1711 | - |
| <i>Chlamydomonas rapa</i> | LH08SG1077 | N53° 5' 14.712" E13° 58' 10.717" | SCH | SEG1 | grassland | drill core | 1694 | - |
| <i>Chlamydomonas rapa</i> | LH08SG9055 | N53° 5' 53.455" E13° 36' 45.241" | SCH | SEG9 | grassland | drill core | 1660 | - |
| <i>Chlamydomonas rapa</i> (cf.) | LH10HG1027 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | soil surface | 1710 | 290 |
| <i>Chlamydomonas typica</i> (cf.) | LH08SG9022 | N53° 5' 53.455" E13° 36' 45.241" | SCH | SEG9 | grassland | drill core | 1262 | - |
| <i>Chlamydropodium vacuolatum</i> | LH10HG1013 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | soil surface | 1728 | 287 |
| <i>Chlorococcum minutum</i> (cf.) | LH08AG701K | N48° 23' 29.116" E9° 22' 36.65" | ALB | AEG7 | grassland | drill core | 793 | - |
| <i>Chlorococcum minutum</i> (cf.) | LH08AW5056 (=SAG 2479) | N48° 25' 10.626" E9° 24' 52.854" | ALB | AEW5 | forest | drill core | 1707 | - |
| <i>Chlorococcum minutum</i> (cf.) | LH08AW5107 | N48° 25' 10.626" E9° 24' 52.854" | ALB | AEW5 | forest | drill core | 1662 | - |
| <i>Chlorococcum minutum</i> (cf.) | LH08AW5111 | N48° 25' 10.626" E9° 24' 52.854" | ALB | AEW5 | forest | drill core | 1346 | - |
| <i>Chlorococcum sphacosum</i> | LH10HG3113 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | soil surface | 1713 | - |
| <i>Coelastrella multistriata</i> | LH08AG2003 | N48° 22' 36.686" E9° 28' 22.023" | ALB | AEG2 | grassland | drill core | 535 | - |
| <i>Coelastrella multistriata</i> | LH08AW4118 | N48° 23' 56.755" E9° 14' 41.378" | ALB | AEW4 | forest | drill core | 1208 | - |
| <i>Coelastrella multistriata</i> | LH10HG7083 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 2117 | - |
| <i>Coelastrella multistriata</i> | LH10HG7097 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 512 | - |
| <i>Coelastrella multistriata</i> | LH10HG7100 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 700 | - |
| <i>Coelastrella multistriata</i> | LH10HG7102 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 411 | - |
| <i>Coelastrella multistriata</i> | LH10HG8109 | N51° 16' 16.527" E10° 25' 4.6" | HAI | HEG8 | grassland | soil surface | 454 | - |
| <i>Coelastrella</i> sp. | LH10HG1009 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | soil surface | 1770 | - |
| <i>Coelastrella</i> sp. | LH10HG1033 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | soil surface | 1622 | - |
| <i>Coelastrella</i> sp. | LH10HG2087 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 470 | - |
| <i>Coelastrella</i> sp. | LH10HG2098 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 2206 | - |
| <i>Coelastrella</i> sp. | LH10HG2P01 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 589 | - |
| <i>Coelastrella</i> sp. | LH10HG2P12 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 599 | - |
| <i>Coelastrella</i> sp. | LH10HG5136 | N51° 12' 57.22" E10° 19' 21.096" | HAI | HEG5 | grassland | soil surface | 1313 | - |
| <i>Coelastrella</i> sp. | LH10HG6035 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1511 | - |
| <i>Coelastrella</i> sp. | LH10HG6060 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1616 | 266 |
| <i>Coelastrella</i> sp. | LH10HG7017 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1683 | - |
| <i>Coelastrella</i> sp. | LH10HG7018 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1429 | 297 |
| <i>Coelastrella</i> sp. | LH10HG7023 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1637 | - |
| <i>Coelastrella</i> sp. | LH10HG7030 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1506 | 298 |
| <i>Coelastrella</i> sp. | LH10HG9130 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1053 | - |
| <i>Desmotetra stigmatica</i> | LH08SG2049 | N53° 5' 21.505" E13° 58' 48.169" | SCH | SEG2 | grassland | drill core | 1667 | - |

Table S1. (continuation)

| Species | Identifier | GPS | Exploratory | Plot | Habitat | Sampling | 18S | ITS2 |
|---------------------------------------|---------------------------|--------------------------------------|-------------|------|-----------|--------------|------|------|
| <i>Desmotetra stigmatica</i> | LH10HG6P18 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 494 | - |
| <i>Heterochlamydomonas</i> sp. | LH08AG2004 | N48° 22' 36.686" E9° 28' 22.023" | ALB | AEG2 | grassland | drill core | 1699 | - |
| <i>Jenufa</i> sp. | LH08AW8035 (=SAG 2383) | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1700 | - |
| <i>Jenufa</i> sp. | LH08AW8098 | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1277 | - |
| <i>Oogamochlamys</i> sp.(I) | LH08SG8047 | N53° 6' 50.294" E14° 1' 1.559" | SCH | SEG8 | grassland | drill core | 1654 | - |
| <i>Oogamochlamys</i> sp.(II) | LH08AW1069 (=SAG 2476) | N48° 28' 41.063" E9° 20' 3.877" | ALB | AEW1 | forest | drill core | 1279 | - |
| <i>Pseudomuriella aurantiaca</i> | LH10HG2039 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1656 | - |
| <i>Pseudomuriella aurantiaca</i> | LH10HG9038 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 2081 | - |
| <i>Stephanosphaerinia</i> sp. | LH10HG6108 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1719 | - |
| <i>Tatrensinia</i> sp.(I) | LH08SW7115 | N53° 6' 26.453" E13° 41' 39.908" | SCH | SEW7 | forest | drill core | 1716 | - |
| <i>Tatrensinia</i> sp.(II) | LH10HG7016 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1733 | - |
| <i>Tatrensinia</i> sp.(II) | LH10HG9131 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 527 | - |
| <i>Pedinomonas minor</i> (cf.) | LH08SG2033 | N53° 5' 21.505" E13° 58' 48.169" | SCH | SEG2 | grassland | drill core | 1489 | - |
| <i>Auxenochlorella protothecoides</i> | LH10HG5119 | N51° 12' 57.22" E10° 19' 21.096" | HAI | HEG5 | grassland | soil surface | 1711 | - |
| <i>Auxenochlorella protothecoides</i> | LH10HG6096 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1725 | - |
| <i>Auxenochlorella protothecoides</i> | LH10HG7124 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1720 | - |
| <i>Auxenochlorella</i> sp. | LH08AW4103 (=SAG 2478) | N48° 23' 56.755" E9° 14' 41.378" | ALB | AEW4 | forest | drill core | 1655 | 321 |
| <i>Chlorella mirabilis</i> (cf.) | LH08AG9040 | N48° 23' 40.815" E9° 30' 10.053" | ALB | AEG9 | grassland | drill core | 1648 | 314 |
| <i>Chlorella mirabilis</i> (cf.) | LH10HG6139 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1683 | - |
| <i>Chlorella vulgaris</i> | LH08HG1081 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | drill core | 1687 | 300 |
| <i>Chlorella vulgaris</i> | LH08HG2013 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | drill core | 1692 | - |
| <i>Chlorella vulgaris</i> | LH08HG2065 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | drill core | 1695 | - |
| <i>Chlorella vulgaris</i> | LH08HG2083 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | drill core | 1676 | - |
| <i>Chlorella vulgaris</i> | LH08HG2091 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | drill core | 1695 | - |
| <i>Chlorella vulgaris</i> | LH08HG2096 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | drill core | 1678 | - |
| <i>Chlorella vulgaris</i> | LH08HG4032 | N51° 6' 48.104" E10° 26' 10.249" | HAI | HEG4 | grassland | drill core | 1671 | 300 |
| <i>Chlorella vulgaris</i> | LH08HG4088 | N51° 6' 48.104" E10° 26' 10.249" | HAI | HEG4 | grassland | drill core | 1692 | 300 |
| <i>Chlorella vulgaris</i> | LH08HG5074 | N51° 12' 57.22" E10° 19' 21.096" | HAI | HEG5 | grassland | drill core | 1695 | 300 |
| <i>Chlorella vulgaris</i> | LH08HG5082 | N51° 12' 57.22" E10° 19' 21.096" | HAI | HEG5 | grassland | drill core | 1692 | 300 |
| <i>Chlorella vulgaris</i> | LH08HW6087 | N51° 16' 3.791" E10° 14' 21.762" | HAI | HEW6 | forest | drill core | 303 | - |
| <i>Chlorella vulgaris</i> | LH08HW9060 | N51° 7' 48.871" E10° 22' 52.139" | HAI | HEW9 | forest | drill core | 1686 | - |
| <i>Chlorella vulgaris</i> | LH08HW9094 | N51° 7' 48.871" E10° 22' 52.139" | HAI | HEW9 | forest | drill core | 1686 | 300 |
| <i>Chlorella vulgaris</i> | LH08SG1071 | N53° 5' 14.712" E13° 58' 10.717" | SCH | SEG1 | grassland | drill core | 1695 | 300 |
| <i>Chlorella vulgaris</i> | LH08SG3006 | N53° 6' 10.204" E13° 59' 8.519" | SCH | SEG3 | grassland | drill core | 1695 | 300 |
| <i>Chlorella vulgaris</i> | LH10HG2049 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1720 | 300 |
| <i>Chlorella vulgaris</i> | LH10HG3088 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | soil surface | 1674 | - |

Table S1. (continuation)

| Species | Identifier | GPS | Exploratory | Plot | Habitat | Sampling | 18S | ITS2 |
|---|---------------------------|--------------------------------------|-------------|------|-----------|--------------|------|------|
| <i>Chlorella vulgaris</i> | LH10HG4093 | N51° 6' 48.104" E10° 26' 10.249" | HAI | HEG4 | grassland | soil surface | 1724 | - |
| <i>Chlorella vulgaris</i> | LH10HG6014 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1719 | 300 |
| <i>Chlorella vulgaris</i> | LH10HG6052 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1735 | 300 |
| <i>Chlorella vulgaris</i> | LH10HG6099 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1665 | - |
| <i>Chlorella vulgaris</i> | LH10HG9075 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1720 | 300 |
| <i>Chlorella vulgaris</i> (cf.) | LH10HG2081 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1690 | 305 |
| <i>Chlorella vulgaris</i> (cf.) | LH10HG7072 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1448 | - |
| <i>Chloroidium ellipsoideum</i> (cf.) | LH08AG8046 | N48° 25' 21.505" E9° 29' 31.649" | ALB | AEG8 | grassland | drill core | 1658 | - |
| <i>Chloroidium ellipsoideum</i> (cf.) | LH08AG9062 | N48° 23' 40.815" E9° 30' 10.053" | ALB | AEG9 | grassland | drill core | 1706 | - |
| <i>Chloroidium ellipsoideum</i> (cf.) | LH10HG6086 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1677 | - |
| <i>Chloroidium ellipsoideum</i> (cf.) | LH10HG7132 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1632 | - |
| <i>Chloroidium ellipsoideum</i> (cf.) | LH10HG9105 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1724 | - |
| <i>Chloroidium saccharophilum</i> | LH08AW8042 | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1694 | - |
| <i>Chloroidium saccharophilum</i> | LH10HG5089 | N51° 12' 57.22" E10° 19' 21.096" | HAI | HEG5 | grassland | soil surface | 568 | - |
| <i>Chloroidium saccharophilum</i> | LH10HG7062 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1719 | - |
| <i>Coccomyxa viridis</i> | LH08AW8039 (=SAG 2483) | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1731 | - |
| <i>Coccomyxa simplex</i> | LH08SG9051 | N53° 5' 53.455" E13° 36' 45.241" | SCH | SEG9 | grassland | drill core | 1409 | - |
| <i>Coccomyxa</i> sp. | LH08AW1017 | N48° 28' 41.063" E9° 20' 3.877" | ALB | AEW1 | forest | drill core | 1732 | - |
| <i>Dictyochloropsis splendida</i> | LH08AW3050 | N48° 24' 44.145" E9° 21' 20.127" | ALB | AEW3 | forest | drill core | 1722 | - |
| <i>Diplosphaera</i> sp.(I) | LH08HW8075 | N51° 21' 20.852" E10° 31' 1.083" | HAI | HEW8 | forest | drill core | 1824 | 326 |
| <i>Diplosphaera</i> sp.(II) | LH08AG9089 | N48° 23' 40.815" E9° 30' 10.053" | ALB | AEG9 | grassland | drill core | 1660 | 334 |
| <i>Lobosphaera bisecta</i> (cf.) | LH10HG3P15 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | soil surface | 601 | - |
| <i>Lobosphaera irregularis</i> (cf.) | LH08AW3064 | N48° 24' 44.145" E9° 21' 20.127" | ALB | AEW3 | forest | drill core | 1731 | - |
| <i>Lobosphaera irregularis</i> (cf.) | LH08SW5063 | N53° 3' 25.321" E13° 53' 7.318" | SCH | SEW5 | forest | drill core | 705 | - |
| <i>Muriella terrestris</i> (cf.) | LH08SG3009 | N53° 6' 10.204" E13° 59' 8.519" | SCH | SEG3 | grassland | drill core | 1692 | 312 |
| <i>Muriella terrestris</i> (cf.) | LH10HG1070 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | soil surface | 1767 | - |
| <i>Muriella terrestris</i> (cf.) | LH10HG1076 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | soil surface | 1732 | 312 |
| <i>Muriella terrestris</i> (cf.) | LH10HG7118 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1675 | - |
| <i>Muriella terrestris</i> (cf.) | LH10HG8050 | N51° 16' 16.527" E10° 25' 4.6" | HAI | HEG8 | grassland | soil surface | 1733 | 311 |
| <i>Muriella terrestris</i> (cf.) | LH10HG9077 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1729 | - |
| <i>Nannochloris</i> sp. | LH08SG3102 | N53° 6' 10.204" E13° 59' 8.519" | SCH | SEG3 | grassland | drill core | 802 | - |
| <i>Nannochloris</i> sp. | LH08SG8030 | N53° 6' 50.294" E14° 1' 1.559" | SCH | SEG8 | grassland | drill core | 1648 | - |
| <i>Nannochloris</i> sp. | LH10HG6095 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1721 | - |
| <i>Navichloris fusiformis</i> (provisional denomination) | LH08AW3007 (=SAG 2477) | N48° 24' 44.145" E9° 21' 20.127" | ALB | AEW3 | forest | drill core | 1694 | - |
| <i>Neocystis brevis</i> | LH08AG9012 | N48° 23' 40.815" E9° 30' 10.053" | ALB | AEG9 | grassland | drill core | 1673 | - |
| <i>Neocystis brevis</i> | LH08AW8001 (=SAG 2480) | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1735 | - |

Table S1. (continuation)

| Species | Identifier | GPS | Exploratory | Plot | Habitat | Sampling | 18S | ITS2 |
|--|---------------------------|--------------------------------------|-------------|------|-----------|--------------|------|------|
| <i>Neocystis brevis</i> | LH08HW6059 | N51° 16' 3.791" E10° 14' 21.762" | HAI | HEW6 | forest | drill core | 1713 | - |
| <i>Neocystis brevis</i> | LH08HW6108 | N51° 16' 3.791" E10° 14' 21.762" | HAI | HEW6 | forest | drill core | 1687 | - |
| <i>Neocystis brevis</i> | LH08SG2072 | N53° 5' 21.505" E13° 58' 48.169" | SCH | SEG2 | grassland | drill core | 1696 | - |
| <i>Neocystis brevis</i> | LH10HG4082 | N51° 6' 48.104" E10° 26' 10.249" | HAI | HEG4 | grassland | soil surface | 357 | - |
| <i>Neocystis brevis</i> | LH10HG9054 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1731 | - |
| <i>Neocystis brevis</i> | LH10HG9080 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1736 | 295 |
| <i>Neocystis brevis</i> | LH10HG9P02 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 543 | - |
| <i>Pseudostichococcus monallantoides</i> | LH08AG7097 | N48° 23' 29.116" E9° 22' 36.65" | ALB | AEG7 | grassland | drill core | 392 | - |
| <i>Pseudostichococcus monallantoides</i> | LH10HG2066 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1709 | - |
| <i>Pseudostichococcus monallantoides</i> | LH10HG3045 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | soil surface | 1737 | 277 |
| <i>Pseudostichococcus</i> sp. | LH08SW8044 | N53° 11' 30.47" E13° 55' 49.216" | SCH | SEW8 | forest | drill core | 1661 | 285 |
| <i>Stichococcus</i> sp.(I) | LH08AG9028 | N48° 23' 40.815" E9° 30' 10.053" | ALB | AEG9 | grassland | drill core | 1245 | - |
| <i>Stichococcus</i> sp.(I) | LH08AW8025 | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1712 | - |
| <i>Stichococcus</i> sp.(II) | LH08AG1100 | N48° 23' 52.818" E9° 20' 31.152" | ALB | AEG1 | grassland | drill core | 573 | - |
| <i>Stichococcus</i> sp.(II) | LH08SG5057 | N53° 6' 26.83" E14° 0' 1.885" | SCH | SEG5 | grassland | drill core | 1695 | 302 |
| <i>Stichococcus</i> sp.(II) | LH08SG5067 | N53° 6' 26.83" E14° 0' 1.885" | SCH | SEG5 | grassland | drill core | 1659 | - |
| <i>Stichococcus</i> sp.(II) | LH08SG5079 | N53° 6' 26.83" E14° 0' 1.885" | SCH | SEG5 | grassland | drill core | 1692 | 302 |
| <i>Stichococcus</i> sp.(II) | LH08SG5090 | N53° 6' 26.83" E14° 0' 1.885" | SCH | SEG5 | grassland | drill core | 1683 | 302 |
| <i>Stichococcus</i> sp.(II) | LH08SG5092 | N53° 6' 26.83" E14° 0' 1.885" | SCH | SEG5 | grassland | drill core | 785 | - |
| <i>Stichococcus</i> sp.(II) | LH10HG2063 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1721 | 302 |
| <i>Stichococcus</i> sp.(II) | LH10HG2067 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1721 | - |
| <i>Stichococcus</i> sp.(II) | LH10HG3128 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | soil surface | 1721 | - |
| <i>Stichococcus</i> sp.(II) | LH10HG7090 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1719 | 302 |
| <i>Stichococcus</i> sp.(II) | LH10HG7092 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1710 | - |
| <i>Stichococcus</i> sp.(II) | LH10HG9068 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1722 | - |
| <i>Stichococcus</i> sp.(III) | LH08AG7010 | N48° 23' 29.116" E9° 22' 36.65" | ALB | AEG7 | grassland | drill core | 1660 | 274 |
| <i>Stichococcus</i> sp.(III) | LH10HG6110 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | soil surface | 1724 | - |
| <i>Stichococcus</i> sp.(IV) | LH08SG1073 | N53° 5' 14.712" E13° 58' 10.717" | SCH | SEG1 | grassland | drill core | 1696 | 325 |
| <i>Stichococcus</i> sp.(V) | LH08SW1099 | N52° 54' 3.05" E13° 50' 46.921" | SCH | SEW1 | forest | drill core | 1653 | 310 |
| <i>Stichococcus</i> sp.(VI) | LH08AW8023 (=SAG 2482) | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1722 | 311 |
| <i>Stichococcus</i> sp.(VI) | LH08AW8104 | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1687 | 311 |
| <i>Stichococcus</i> sp.(VII) | LH08AW8002 (=SAG 2481) | N48° 22' 57.322" E9° 22' 56.584" | ALB | AEW8 | forest | drill core | 1730 | - |
| Unidentified Chlorellaceae (I) | LH08AG1034 | N48° 23' 52.818" E9° 20' 31.152" | ALB | AEG1 | grassland | drill core | 1681 | 274 |
| Unidentified Chlorellaceae (I) | LH10HG7073 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1409 | - |
| Unidentified Chlorellaceae (II) | LH08SG2053 | N53° 5' 21.505" E13° 58' 48.169" | SCH | SEG2 | grassland | drill core | 1672 | - |
| Unidentified Chlorellaceae (II) | LH08SG3029 | N53° 6' 10.204" E13° 59' 8.519" | SCH | SEG3 | grassland | drill core | 798 | - |

Table S1. (continuation)

| Species | Identifier | GPS | Exploratory | Plot | Habitat | Sampling | 18S | ITS2 |
|--|---------------------------|--------------------------------------|-------------|------|-----------|--------------|------|------|
| Unidentified Chlorellaceae (II) | LH10HG2094 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1719 | - |
| Unidentified Chlorellaceae (II) | LH10HG3135 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | soil surface | 1718 | - |
| Unidentified Chlorellaceae (II) | LH10HG7071 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1732 | - |
| Unidentified Chlorellaceae (II) | LH10HG9020 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1617 | 299 |
| Unidentified Chlorellaceae (II) | LH10HG9026 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | soil surface | 1716 | 274 |
| Unidentified Chlorellaceae (III) | LH08SG3078 | N53° 6' 10.204" E13° 59' 8.519" | SCH | SEG3 | grassland | drill core | 1611 | 299 |
| Unidentified Chlorellaceae (III) | LH08SG3093 | N53° 6' 10.204" E13° 59' 8.519" | SCH | SEG3 | grassland | drill core | 1678 | 299 |
| Unidentified Chlorellaceae (IV) | LH10HG709K | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1685 | - |
| <i>Xylochloris</i> sp. | LH08AG7024 (=SAG 2382) | N48° 23' 29.116" E9° 22' 36.65" | ALB | AEG7 | grassland | drill core | 1719 | - |
| <i>Pseudendocloniopsis botryoides</i> | LH08HW9058 | N51° 7' 48.871" E10° 22' 52.139" | HAI | HEW9 | forest | drill core | 2058 | - |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08AG1113 | N48° 23' 52.818" E9° 20' 31.152" | ALB | AEG1 | grassland | drill core | 2038 | - |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08AG7011 | N48° 23' 29.116" E9° 22' 36.65" | ALB | AEG7 | grassland | drill core | 2236 | - |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08AG9045 | N48° 23' 40.815" E9° 30' 10.053" | ALB | AEG9 | grassland | drill core | 2093 | - |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08HW1038 | N51° 11' 7.278" E10° 19' 25.036" | HAI | HEW1 | forest | drill core | 1882 | 274 |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08HW4109 | N51° 22' 10.134" E10° 31' 59.979" | HAI | HEW4 | forest | drill core | 2234 | - |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08HW5021 | N51° 15' 49.961" E10° 14' 27.448" | HAI | HEW5 | forest | drill core | 1827 | - |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08HW9005 | N51° 7' 48.871" E10° 22' 52.139" | HAI | HEW9 | forest | drill core | 1736 | - |
| <i>Klebsormidium</i> <i>dissectum/elegans</i> (cf.) | LH08HW9106 | N51° 7' 48.871" E10° 22' 52.139" | HAI | HEW9 | forest | drill core | 2211 | 274 |
| <i>Klebsormidium flaccidum</i> (cf.) | LH08HW5061 | N51° 15' 49.961" E10° 14' 27.448" | HAI | HEW5 | forest | drill core | 2064 | 274 |
| <i>Klebsormidium flaccidum</i> (cf.) | LH08SG3027 | N53° 6' 10.204" E13° 59' 8.519" | SCH | SEG3 | grassland | drill core | 1817 | - |
| <i>Klebsormidium flaccidum</i> (cf.) | LH10HG2056 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | soil surface | 1777 | 274 |
| <i>Klebsormidium flaccidum</i> (cf.) | LH10HG7028 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | soil surface | 1794 | 275 |
| <i>Asterosiphon</i> sp. | LH10HG3064 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | drill core | 1780 | - |
| <i>Botrydiopsalean</i> sp. | LH08AW1076 | N48° 28' 41.063" E9° 20' 3.877" | ALB | AEW1 | forest | soil surface | 1763 | - |
| <i>Botrydiopsalean</i> sp. | LH08SG2K53 | N53° 5' 21.505" E13° 58' 48.169" | SCH | SEG2 | grassland | soil surface | 619 | - |
| <i>Botrydiopsis callosa</i> (cf.) | LH08AW4043 | N48° 23' 56.755" E9° 14' 41.378" | ALB | AEW4 | forest | soil surface | 1753 | - |
| <i>Heterococcus</i> sp. | LH10HG2140 | N51° 0' 2.696" E10° 25' 48.036" | HAI | HEG2 | grassland | drill core | 1734 | - |
| <i>Heterococcus</i> sp. | LH10HG9085 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | drill core | 1734 | - |
| <i>Heterococcus chodatii</i> (cf.) | LH10HG9111 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | drill core | 1693 | - |
| <i>Heterococcus chodatii</i> (cf.) | LH10HG9126 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | drill core | 627 | - |
| <i>Heterococcus caespitosus</i> (cf.) | LH08AG2020 | N48° 22' 36.686" E9° 28' 22.023" | ALB | AEG2 | grassland | soil surface | 1690 | - |
| <i>Heterothrix sessile</i> | LH10HG5079 | N51° 12' 57.22" E10° 19' 21.096" | HAI | HEG5 | grassland | drill core | 1791 | - |
| <i>Heterothrix sessile</i> | LH10HG9037 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | drill core | 1568 | - |
| <i>Heterothrix</i> sp. | LH08SG5052 | N53° 6' 26.83" E14° 0' 1.885" | SCH | SEG5 | grassland | soil surface | 1779 | - |
| <i>Heterothrix</i> sp. | LH10HG7061 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | drill core | 1806 | - |

Table S1. (continuation)

| Species | Identifier | GPS | Exploratory | Plot | Habitat | Sampling | 18S | ITS2 |
|--------------------------------------|------------|--------------------------------------|-------------|------|-----------|--------------|------|------|
| <i>Xanthonema bristolianum</i> (cf.) | LH08HW9018 | N51° 7' 48.871" E10° 22' 52.139" | HAI | HEW9 | forest | soil surface | 1752 | - |
| <i>Xanthonema bristolianum</i> (cf.) | LH10HG6059 | N51° 12' 53.766" E10° 23' 28.395" | HAI | HEG6 | grassland | drill core | 1780 | - |
| <i>Xanthonema exile</i> (cf.) | LH10HG7078 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | drill core | 1783 | - |
| <i>Xanthonema</i> sp. | LH10HG1K69 | N50° 58' 17.934" E10° 24' 19.306" | HAI | HEG1 | grassland | drill core | 1716 | - |
| <i>Xanthonema</i> sp. | LH10HG3065 | N50° 59' 53.129" E10° 25' 58.616" | HAI | HEG3 | grassland | drill core | 1783 | - |
| <i>Xanthonema</i> sp. | LH10HG7029 | N51° 16' 24.897" E10° 24' 37.485" | HAI | HEG7 | grassland | drill core | 1782 | - |
| <i>Xanthonema</i> sp. | LH10HG8112 | N51° 16' 16.527" E10° 25' 4.6" | HAI | HEG8 | grassland | drill core | 374 | - |
| <i>Xanthonema</i> sp. | LH10HG9031 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | drill core | 1782 | - |
| <i>Xanthonema</i> sp. | LH10HG9058 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | drill core | 1782 | - |
| <i>Xanthonema</i> sp. | LH10HW9129 | N51° 7' 48.871" E10° 22' 52.139" | HAI | HEW9 | forest | drill core | 1575 | - |
| <i>Eustigmatos</i> sp. | LH10HG5036 | N51° 12' 57.22" E10° 19' 21.096" | HAI | HEG5 | grassland | drill core | 1764 | - |
| <i>Eustigmatos</i> sp. | LH10HG9133 | N51° 13' 26.031" E10° 22' 50.834" | HAI | HEG9 | grassland | drill core | 1691 | - |

Legend. ALB=Schwäbische Alb, HAI=Hainich-Dün, SCH=Schorfheide-Chorin.

Table S2a. List of all detected species belonging to Trebouxiophyceae.

| Species | Clade | Representative isolates and SAG-strains | GenBank |
|--|---------------------|---|----------|
| <i>Auxenochlorella protothecoides</i> | Chlorellaceae | LH10HG6096 | - |
| <i>Auxenochlorella</i> sp. | Chlorellaceae | LH08AW4103 (=SAG 2478) | KP081390 |
| <i>Chlorella mirabilis</i> (cf.) | <i>Prasiola</i> | LH08AG9040 | - |
| <i>Chlorella mirabilis</i> (cf.) | <i>Prasiola</i> | LH10HG6139 | - |
| <i>Chlorella vulgaris</i> | Chlorellaceae | LH08HG1081 | - |
| <i>Chlorella vulgaris</i> | Chlorellaceae | LH08SG3006 | - |
| <i>Chlorella vulgaris</i> (cf.) | Chlorellaceae | LH10HG2081 (partial 18S) | - |
| <i>Chlorella vulgaris</i> (cf.) | Chlorellaceae | LH10HG7072 (partial 18S) | - |
| <i>Chloroidium ellipsoideum</i> (cf.) | <i>Watanabea</i> | LH10HG9105 | - |
| <i>Chloroidium saccharophilum</i> | <i>Watanabea</i> | LH10HG7062 | - |
| <i>Coccomyxa viridis</i> | <i>Botryococcus</i> | LH08AW8039 (=SAG 2483) | KP081391 |
| <i>Coccomyxa simplex</i> | <i>Botryococcus</i> | LH08SG9051 (partial 18S) | - |
| <i>Coccomyxa</i> sp. | <i>Botryococcus</i> | LH08AW1017 | KP081392 |
| <i>Dictyochloropsis splendida</i> | incertae sedis | LH08AW3050 (= LH-aw3050) | JQ988930 |
| <i>Diplosphaera</i> sp.(I) | <i>Prasiola</i> | LH08HW8075 | - |
| <i>Diplosphaera</i> sp.(II) | <i>Prasiola</i> | LH08AG9089 | - |
| <i>Lobosphaera bisecta</i> (cf.) | <i>Lobosphaera</i> | LH10HG3P15 (partial 18S) | - |
| <i>Lobosphaera irregularis</i> (cf.) | <i>Lobosphaera</i> | LH08AW3064 | KP081398 |
| <i>Muriella terrestris</i> (cf.) | Chlorellaceae | LH08SG3009 | - |
| <i>Nannochloris</i> sp. | Chlorellaceae | LH08SG8030 | - |
| <i>Navichloris fusiformis</i> (proposed taxon) | incertae sedis | LH08AW3007 (=SAG 2477) | KP081399 |
| <i>Neocystis brevis</i> | <i>Neocystis</i> | LH10HG9080 | - |
| <i>Pseudostichococcus monallantoides</i> | <i>Prasiola</i> | LH10HG3045 | - |
| <i>Pseudostichococcus</i> sp. | <i>Prasiola</i> | LH08SW8044 | - |
| <i>Stichococcus</i> sp.(I) | <i>Prasiola</i> | LH08AW8025 | KP081397 |
| <i>Stichococcus</i> sp.(II) | <i>Prasiola</i> | LH08SG5057 | - |
| <i>Stichococcus</i> sp.(III) | <i>Prasiola</i> | LH10HG6110, LH08AG7010 | - |
| <i>Stichococcus</i> sp.(IV) | <i>Prasiola</i> | LH08SG1073 | - |
| <i>Stichococcus</i> sp.(V) | <i>Prasiola</i> | LH08SW1099 | - |
| <i>Stichococcus</i> sp.(VI) | <i>Prasiola</i> | LH08AW8023 (=SAG 2482) | KP081395 |
| <i>Stichococcus</i> sp.(VII) | <i>Prasiola</i> | LH08AW8002 (=SAG 2481) | KP081394 |
| Unidentified Chlorellaceae (I) | Chlorellaceae | LH08AG1034 | - |
| Unidentified Chlorellaceae (II) | Chlorellaceae | LH10HG9020 | - |
| Unidentified Chlorellaceae (III) | Chlorellaceae | LH08SG3093 | - |
| Unidentified Chlorellaceae (IV) | Chlorellaceae | LH10HG709K | - |
| <i>Xylochloris</i> sp. | incertae sedis | LH08AG7024 (= SAG 2382) | JQ988942 |

Table S2b. List of all detected species belonging to Chlorophyceae, other green algae and Stramenopiles.

| Species | Clade | Representative isolates and SAG-strains | GenBank |
|--|-------------------|---|----------|
| <i>Acutodesmus rubescens</i> | Sphaeropleales | LH08SG8041 | - |
| <i>Bracteacoccus cohaerens</i> | Sphaeropleales | LH10HG9034 | - |
| <i>Bracteacoccus cohaerens</i> (cf.) | Sphaeropleales | LH08SG2015 | - |
| <i>Chlamydomonas gerloffii</i> (cf.) | Chlamydomonadales | LH08SW5031 | - |
| <i>Chlamydomonas rapa</i> | Chlamydomonadales | LH08SG1077 | - |
| <i>Chlamydomonas rapa</i> (cf.) | Chlamydomonadales | LH10HG1027 | - |
| <i>Chlamydomonas typica</i> (cf.) | Chlamydomonadales | LH08SG9022 (partial 18S) | - |
| <i>Chlamydropodium vacuolatum</i> | Chlamydomonadales | LH10HG1013 | - |
| <i>Chlorococcum minutum</i> (cf.) | Chlamydomonadales | LH08AW5056 (=SAG 2479) | KP081402 |
| <i>Chlorococcum sphacosum</i> | Chlamydomonadales | LH10HG3113 | - |
| <i>Coelastrella multistriata</i> | Sphaeropleales | LH10HG7083 | - |
| <i>Coelastrella</i> sp. | Sphaeropleales | LH10HG2098 (+LH10HG7018) | - |
| <i>Desmotetra stigmatica</i> | Chlamydomonadales | LH08SG2049 | - |
| <i>Heterochlamydomonas</i> sp. | Chlamydomonadales | LH08AG2004 | - |
| <i>Jenufa</i> sp. | incertae sedis | LH08AW8035 (= SAG 2383) | JQ988933 |
| <i>Oogamochlamys</i> sp.(I) | Chlamydomonadales | LH08SG8047 | - |
| <i>Oogamochlamys</i> sp.(II) | Chlamydomonadales | LH08AW1069 (= SAG 2476; partial 18S) | KP081401 |
| <i>Pseudomuriella aurantiaca</i> | Sphaeropleales | LH10HG2039 | - |
| <i>Stephanosphaerinia</i> sp. | Chlamydomonadales | LH10HG6108 | - |
| <i>Tatrensinia</i> sp.(I) | Chlamydomonadales | LH08SW7115 | - |
| <i>Tatrensinia</i> sp.(II) | Chlamydomonadales | LH10HG7016 (+LH10HG9131) | - |
| <i>Pseudendocloniopsis botryoides</i> (cf.) | Ulotrichales | LH08HW9058 | - |
| <i>Pedinomonas minor</i> (cf.) | Pedinophyceae | LH08SG2033 (partial 18S) | - |
| <i>Klebsormidium dissectum/elegans</i> (cf.) | Streptophyta | LH08HW9106 | - |
| <i>Klebsormidium flaccidum</i> (cf.) | Streptophyta | LH10HG2056 | - |
| <i>Asterosiphon</i> sp. | incertae sedis | LH10HG3064 | - |
| <i>Botrydiopsalean</i> sp. | Botrydiopsalean | LH08AW1076 | - |
| <i>Botrydiopsis callosa</i> (cf.) | Botrydiopsalean | LH08AW4043 | - |
| <i>Eustigmatos</i> sp. | Eustigmatophyceae | LH10HG9133, LH10HG5036 | - |
| <i>Heterococcus caespitosus</i> (cf.) | Chlorellidialean | LH08AG2020 | - |
| <i>Heterococcus chodatii</i> (cf.) | Chlorellidialean | LH10HG9111 | - |
| <i>Heterococcus</i> sp. | Chlorellidialean | LH10HG9085, LH10HG2140 | - |
| <i>Heterothrix sessile</i> | Tribonematalean | LH10HG5079 | - |
| <i>Heterothrix</i> sp. | Tribonematalean | LH10HG7061, LH08SG5052 | - |
| <i>Xanthonema bristolianum</i> (cf.) | Tribonematalean | LH08HW9018 | - |
| <i>Xanthonema exile</i> (cf.) | Tribonematalean | LH10HG7078 | - |
| <i>Xanthonema</i> sp. | Tribonematalean | LH10HG9058, LH10HG7029 | - |

Table S3. List of the closest GenBank-relatives of our detected species.

| Representative isolate(s) in 18S tree | Closest GenBank relatives; authentic strain* | GenBank | Similarity % | Habitat of origin | Land |
|--|---|----------|--------------|---------------------------|------|
| LH10HG6096 | <i>Auxenochlorella protothecoides</i> SAG 211-7a* | X56101 | 100.00 | terrestrial/sap | DE |
| SAG 2478) | <i>Auxenochlorella protothecoides</i> SAG 211-7a* | X56101 | 97.46 | terrestrial/sap | DE |
| LH08AG9040 | <i>Chlorella mirabilis</i> SAG 38.88* | X74000 | 99.88 | terrestrial/soil | RU |
| LH10HG6139 | <i>Chlorella mirabilis</i> SAG 38.88* | X74000 | 99.82 | terrestrial/soil | RU |
| LH08HG1081 | <i>Chlorella vulgaris</i> SAG 211-11b* | FM205832 | 99.94 | freshwater | NL |
| LH08SG3006 | <i>Chlorella vulgaris</i> SAG 211-11b* | FM205832 | 100.00 | freshwater | NL |
| LH10HG2081 (PS) | <i>Chlorella vulgaris</i> SAG 211-11b* | FM205832 | 99.75 | freshwater | NL |
| LH10HG7072 (PS) | <i>Chlorella vulgaris</i> SAG 211-11b* | FM205832 | 99.82 | freshwater | NL |
| LH10HG9105 | <i>Chloroidium ellipsoideum</i> SAG 3.95* | FM946012 | 99.88 | n/a | n/a |
| LH10HG7062 | <i>Chloroidium saccharophilum</i> SAG 211-9a* | FM946000 | 100.00 | terrestrial/sap | DE |
| SAG 2483 | <i>Coccomyxa viridis</i> CCALA 306 | AM167525 | 100.00 | terrestrial/lichen | CZ |
| LH08SG9051 (PS) | <i>Coccomyxa simplex</i> SAG 216-8* | HQ317304 | 100.00 | freshwater | RO |
| LH08AW1017 | <i>Coccomyxa</i> sp. KN-2011-T3 | HE586515 | 99.94 | terrestrial/lichen | ID |
| LH08AW3050 | <i>Dictyochloropsis splendida</i> CAUP H8601 | GU017662 | 100.00 | terrestrial/soil/fumarole | CZ |
| LH08HW8075 | <i>Chlorella sphaerica</i> SAG 11.88* | AJ416105 | 99.94 | terrestrial/lichen | NZ |
| LH08AG9089 | <i>Chlorella sphaerica</i> SAG 11.88* | AJ416105 | 99.82 | terrestrial/lichen | NZ |
| LH10HG3P15 (PS) | <i>Myrmecia bisecta</i> SAG 2043* | Z47209 | 100.00 | terrestrial/soil | IT |
| LH08AW3064 | <i>Myrmecia irregularis</i> CCAP 221/8 | HQ902935 | 100.00 | n/a | n/a |
| LH08SG3009 | <i>Muriella terrestris</i> ASIB V38 | AB012845 | 99.94 | terrestrial/soil | IT |
| LH08SG8030 | <i>Nannochloris bacillaris</i> | AB080300 | 99.94 | n/a | n/a |
| SAG 2477) | <i>Leptochlorella</i> sp. clone QE17 | FJ790649 | 95.54 | terrestrial/epilithic | CN |
| LH10HG9080 | <i>Neocystis brevis</i> CAUP D 802* | JQ920360 | 100.00 | terrestrial/soil | CH |
| LH10HG3045 | <i>Stichococcus mirabilis</i> CCAP 379/3 | AJ311638 | 99.82 | n/a | n/a |
| LH08SW8044 | <i>Stichococcus mirabilis</i> CCAP 379/3 | AJ311638 | 100.00 | n/a | n/a |
| LH08AW8025 | <i>Trebouxia</i> sp. UR47/4 | AY762604 | 100.00 | terrestrial/facade | DE |
| LH08SG5057 | <i>Trebouxia</i> sp. UR55/3 | AY762606 | 100.00 | terrestrial/facade | DE |
| LH10HG6110, LH08AG7010 | <i>Stichococcus deasonii</i> UTEX 1706* | DQ275460 | 99.70 | terrestrial/soil | USA |
| LH08SG1073 | <i>Stichococcus</i> sp.4 WB47 | KF144240 | 100.00 | freshwater/biofilm | DE |
| LH08SW1099 | <i>Stichococcus deasonii</i> UTEX 1706* | DQ275460 | 98.90 | terrestrial/soil | USA |
| SAG 2482 | <i>Stichococcus</i> sp.2 D4-2A | KF144238 | 100.00 | freshwater/biofilm | DE |
| SAG 2481 | <i>Stichococcus</i> K4-4 | AB055866 | 100.00 | n/a | n/a |
| LH08AG1034 | <i>Marvania</i> sp. WB67 | KF144207 | 100.00 | freshwater/biofilm | DE |
| LH10HG9020 | <i>Nannochloris</i> sp. JL-4-6 | AY195983 | 100.00 | freshwater | USA |
| LH08SG3093 | <i>Nannochloris</i> sp. Ant-1 | EF440182 | 100.00 | terrestrial/permafrost | AQ |
| LH10HG709K | <i>Nannochloris</i> sp. AS 2-10 | AY195968 | 97.76 | freshwater | USA |
| SAG 2382 | <i>Xylochloris irregularis</i> CAUP H7801* | EU105209 | 96.47 | terrestrial/epixylic | SG |
| LH08SG8041 | <i>Scenedesmus rubescens</i> CCAP 232/1 | X74002 | 100.00 | n/a | n/a |
| LH10HG9034 | <i>Bracteacoccus cohaerens</i> UTEX 1272* | GQ985406 | 100.00 | terrestrial/soil | USA |
| LH08SG2015 | <i>Bracteacoccus cohaerens</i> UTEX 1272* | GQ985406 | 99.94 | terrestrial/soil | USA |
| LH08SW5031 | <i>Chlamydomonas gerloffii</i> CCAP 11/72* | FR865610 | 99.82 | freshwater | CZ |
| LH08SG1077 | <i>Chlamydomonas rapa</i> SAG 48.72* | U70790 | 100.00 | freshwater/plankton | SK |
| LH10HG1027 | <i>Chlamydomonas rapa</i> SAG 48.72* | U70790 | 99.94 | freshwater/plankton | SK |
| LH08SG9022 (PS) | <i>Chlamydomonas typica</i> SAG 61.72* | AB701557 | 99.92 | terrestrial/soil | USA |
| LH10HG1013 | <i>Chlorococcum robustum</i> Kr 86 30 | AY122332 | 100.00 | n/a | n/a |

Table S3. (continuation)

| Representative isolate(s) in 18S tree | Closest GenBank relatives; authentic strain* | GenBank | Similarity % | Habitat of origin | Land |
|--|---|----------|--------------|-----------------------|------|
| SAG 2479 | <i>Chlorococcum minutum</i> SAG 21.95 | JN968585 | 99.77 | terrestrial/soil | IT |
| LH10HG3113 | <i>Chlorococcum sphacosum</i> SAG 66.80* | JN968580 | 100.00 | terrestrial/soil | USA |
| LH10HG7083 | <i>Coelastrella multistriata</i> Hanagata C6-2 | AB012846 | 100.00 | terrestrial/epixylic | JP? |
| LH10HG2098 (+LH10HG7018) | <i>Scenedesmus</i> sp. KGU Y002 | AB742453 | 100.00 | n/a | n/a |
| LH08SG2049 | <i>Desmotetra stigmatica</i> UTEX 962* | DQ009760 | 100.00 | n/a | n/a |
| LH08AG2004 | <i>Heterochlamydomonas rugosa</i> SAG 45.86 | AF367859 | 99.76 | freshwater | UK |
| SAG 2383) | <i>Jenufa minuta</i> CAUP H 8102* | HM563744 | 98.21 | terrestrial/epixylic | SG |
| LH08SG8047 | <i>Oogamochlamys gigantea</i> SAG 21.72* | AJ410468 | 98.41 | terrestrial/soil | USA |
| SAG 2476 (PS) | <i>Chlamydomonas</i> sp. CCAP 11/159 | FR865553 | 97.69 | freshwater | USA |
| LH10HG2039 | <i>Pseudomuriella aurantiaca</i> SAG 249-1* | X91268 | 100.00 | terrestrial/soil | CH |
| LH10HG6108 | <i>Chlorococcales</i> sp. VII3 | FJ946904 | 99.59 | freshwater | AQ |
| LH08SW7115 | Uncultured Haematococcaceae clone Amb_18S_582 | EF023273 | 99.77 | terrestrial/soil | NL |
| LH10HG7016 (+LH10HG9131) | <i>Chlorococcum</i> cf. <i>tatrense</i> CCCryo 101-99 | AF514407 | 99.71 | n/a | n/a |
| LH08HW9058 | <i>Pseudocloniopsis botryoides</i> SAG 465-1* | AJ416103 | 99.88 | freshwater | CH |
| LH08SG2033 (PS) | <i>Pedinomonas minor</i> SAG 1965-3 | HE610132 | 99.80 | freshwater/plankton | SK |
| LH08HW9106 | <i>Klebsormidium dissectum</i> SAG 2155* | EF372518 | 100.00 | terrestrial/soil | FR |
| LH10HG2056 | <i>Klebsormidium flaccidum</i> SAG 7.91 | EU434019 | 100.00 | freshwater | RU |
| LH10HG3064 | <i>Asterosiphon dichotomus</i> UTEX LB 2066 | AM490829 | 98.54 | n/a | n/a |
| LH08AW1076 | <i>Chlorellidium pyrenoidosum</i> PAB 785 | AJ579338 | 99.31 | soil(?) | AQ |
| LH08AW4043 | <i>Botrydiopsis callosa</i> SAG 30.83 | AJ579340 | 99.29 | soil | IT |
| LH10HG9133, LH10HG5036 | <i>Eustigmatos magna</i> CCMP 387 | U41051 | 99.82 | terrestrial/soil | NZ |
| LH08AG2020 | <i>Heterococcus protonematoideus</i> SAG 835-9 | AJ579334 | 99.94 | terrestrial/soil | CH |
| LH10HG9111 | <i>Heterococcus chodatii</i> SAG 835-3* | AM490822 | 100.00 | terrestrial/subaerial | CH |
| LH10HG9085, LH10HG2140 | <i>Xanthophyceae</i> sp. IX3 | FJ946906 | 99.59 | freshwater | AQ |
| LH10HG5079 | <i>Heterothrix sessile</i> IBSG-V28 | AM490818 | 100.00 | n/a | n/a |
| LH10HG7061, LH08SG5052 | <i>Heterothrix</i> sp.1 ACOL A1 | AM491612 | 99.26-99.31 | n/a | n/a |
| LH08HW9018 | <i>Xanthonema bristolianum</i> CCALA 516 | AM490819 | 100.00 | terrestrial/snow | SK |
| LH10HG7078 | <i>Xanthonema exile</i> PAB 395 | AM491615 | 99.94 | n/a | n/a |
| LH10HG9058, LH10HG7029 | <i>Xanthonema exile</i> PAB 395 | AM491615 | 99.89+99.77 | n/a | n/a |

Legend. AQ=Antarctica, CH=Switzerland, CN=China, CZ=Czech Republic, DE=Germany, FR=France, ID=Indonesia, IT=Italy, JP=Japan, NL=Netherlands, NZ=New Zealand, RO=Romania, RU=Russia, SG=Singapore, SK=Slovakia, UK=United Kingdom, USA=United States of America.

Table S4a. List of all analyzed full and partial (=PS) 18S rDNA sequences.

| Species | Representative isolates (in phylogenetic trees) | Similar full 18S (OTU/0.00-level) | Similar partial 18S (OTU/0.00-level) |
|--|--|---|--|
| <i>Auxenochlorella protothecoides</i> | LH10HG6096 | LH10HG7124, LH10HG5119 | |
| <i>Auxenochlorella</i> sp. | LH08AW4103 | | |
| <i>Chlorella mirabilis</i> (cf.) | LH08AG9040 | | |
| <i>Chlorella mirabilis</i> (cf.) | LH10HG6139 | | |
| <i>Chlorella vulgaris</i> | LH08HG1081 | | |
| <i>Chlorella vulgaris</i> | LH08SG3006 | LH10HG6014, LH08HG2065, LH08HG2013, LH08HG5074, LH08HW9060, LH10HG2049, LH08HG4088, LH10HG6052, LH10HG9075, LH08HW9094, LH08SG1071, LH08HG5082, LH08HG2083, LH08HG2091, LH08HG2096, LH08HG4032, LH10HG4093, LH10HG6099, LH10HG3088 | LH08HW6087 |
| <i>Chlorella vulgaris</i> (cf.) | LH10HG7072 (PS) | | |
| <i>Chlorella vulgaris</i> (cf.) | LH10HG2081 (PS) | | |
| <i>Chloroidium ellipsoideum</i> (cf.) | LH10HG9105 | LH10HG6086, LH10HG7132, LH08AG8046, LH08AG9062 | |
| <i>Chloroidium saccharophilum</i> | LH10HG7062 | LH08AW8042 | LH10HG5089 |
| <i>Coccomyxa viridis</i> | SAG 2483 | | |
| <i>Coccomyxa simplex</i> | LH08SG9051 (PS) | | |
| <i>Coccomyxa</i> sp. | LH08AW1017 | | |
| <i>Dictyochloropsis splendida</i> | LH08AW3050 | | |
| <i>Diplosphaera</i> sp.(I) | LH08HW8075 | | |
| <i>Diplosphaera</i> sp.(II) | LH08AG9089 | | |
| <i>Lobosphaera bisecta</i> (cf.) | LH10HG3P15 (PS) | | |
| <i>Lobosphaera irregularis</i> (cf.) | LH08AW3064 | | LH08SW5063 |
| <i>Muriella terrestris</i> (cf.) | LH08SG3009 | LH10HG1070, LH10HG8050, LH10HG1076, LH10HG7118, LH10HG9077 | |
| <i>Nannochloris</i> sp. | LH08SG8030 | LH10HG6095 | LH08SG3102 |
| <i>Navichloris fusiformis</i> (proposed taxon) | SAG 2477 | | |
| <i>Neocystis brevis</i> | LH10HG9080 | LH08SG2072, LH08AG9012, LH08AW8001, LH08HW6108, LH08HW6059, LH10HG4082, LH10HG9054 | LH10HG9P02 |
| <i>Pseudostichococcus monallantoides</i> | LH10HG3045 | LH10HG2066 | LH08AG7097 |
| <i>Pseudostichococcus</i> sp. | LH08SW8044 | | |
| <i>Stichococcus</i> sp.(I) | LH08AW8025 | | LH08AG9028 |
| <i>Stichococcus</i> sp.(II) | LH08SG5057 | LH10HG7092, LH08SG5067, LH08SG5079, LH08SG5090, LH10HG2067, LH10HG2063, LH10HG9068, LH10HG3128, LH10HG7090 | LH08AG1100, LH08SG5092 |
| <i>Stichococcus</i> sp.(III) | LH10HG6110, LH08AG7010 | | |
| <i>Stichococcus</i> sp.(IV) | LH08SG1073 | | |
| <i>Stichococcus</i> sp.(V) | LH08SW1099 | | |
| <i>Stichococcus</i> sp.(VI) | SAG 2482 | LH08AW8104 | |
| <i>Stichococcus</i> sp.(VII) | SAG 2481 | | |
| Unidentified Chlorellaceae (I) | LH08AG1034 | | LH10HG7073 |
| Unidentified Chlorellaceae (II) | LH10HG9020 | LH08SG2053, LH10HG7071, LH10HG9026, LH10HG2094, LH10HG3135 | LH08SG3029 |
| Unidentified Chlorellaceae (III) | LH08SG3093 | LH08SG3078 | |
| Unidentified Chlorellaceae (IV) | LH10HG709K | | |
| <i>Xylochloris</i> sp. | SAG 2382 | | |
| <i>Acutodesmus rubescens</i> | LH08SG8041 | | |
| <i>Bracteacoccus cohaerens</i> | LH10HG9034 | | |
| <i>Bracteacoccus cohaerens</i> (cf.) | LH08SG2015 | | |
| <i>Chlamydomonas gerloffii</i> (cf.) | LH08SW5031 | | |
| <i>Chlamydomonas rapa</i> | LH08SG1077 | LH08SG9055 | |

Table S4a. (continuation)

| Species | Representative isolates (in phylogenetic trees) | Similar full 18S (OTU/0.00-level) | Similar partial 18S (OTU/0.00-level) |
|--|--|---|---|
| <i>Chlamydomonas rapa</i> (cf.) | LH10HG1027 | | |
| <i>Chlamydomonas typica</i> (cf.) | LH08SG9022 (PS) | | |
| <i>Chlamydomonium vacuolatum</i> | LH10HG1013 | | |
| <i>Chlorococcum minutum</i> (cf.) | SAG 2479 | LH08AW5107 | LH08AG701K, LH08AW5111 |
| <i>Chlorococcum sphacosum</i> | LH10HG3113 | | |
| <i>Coelastrella multistriata</i> | LH10HG7083 | | LH10HG7102, LH10HG7100, LH10HG8109, LH10HG7097, LH08AG2003, LH08AW4118 |
| <i>Coelastrella</i> sp. | LH10HG2098 | LH10HG1009, LH10HG1033, LH10HG6060, LH10HG7017, LH10HG7023 | LH10HG2087, LH10HG7018, LH10HG6035, LH10HG7030, LH10HG9130, LH10HG2P12, LH10HG2P01, LH10HG5136 LH10HG6P18 |
| <i>Desmotetra stigmatica</i> | LH08SG2049 | | |
| <i>Heterochlamydomonas</i> sp. | LH08AG2004 | | |
| <i>Jenufa</i> sp. | SAG 2383 | | LH08AW8098 |
| <i>Oogamochlamys</i> sp.(I) | LH08SG8047 | | |
| <i>Oogamochlamys</i> sp.(II) | SAG 2476 | | |
| <i>Pseudomuriella aurantiaca</i> | LH10HG2039 | LH10HG9038 | |
| <i>Stephanosphaerina</i> sp. | LH10HG6108 | | |
| <i>Tatrensinia</i> sp.(I) | LH08SW7115 | | |
| <i>Tatrensinia</i> sp.(II) | LH10HG7016 | | LH10HG9131 |
| <i>Pseudendocloniopsis botryoides</i> (cf.) | LH08HW9058 | | |
| <i>Pedinomonas minor</i> (cf.) | LH08SG2033 (PS) | | |
| <i>Klebsormidium dissectum/elegans</i> (cf.) | LH08HW9106 | LH08AG9045, LH08HW9005, LH08AG1113, LH08HW1038, LH08AG7011, LH08HW4109, LH08HW5021 | |
| <i>Klebsormidium flaccidum</i> (cf.) | LH10HG2056 | LH08HW5061, LH10HG7028, LH08SG3027 | |
| <i>Asterosiphon</i> sp. | LH10HG3064 | | |
| <i>Botrydiopsalean</i> sp. | LH08AW1076 | | LH08SG2K53 |
| <i>Botrydiopsis callosa</i> (cf.) | LH08AW4043 | | |
| <i>Heterococcus caespitosus</i> (cf.) | LH08AG2020 | | |
| <i>Heterococcus chodatii</i> (cf.) | LH10HG9111 | | LH10HG9126 |
| <i>Heterococcus</i> sp. | LH10HG9085, LH10HG2140 | | |
| <i>Heterothrix sessile</i> | LH10HG5079 | LH10HG9037 | |
| <i>Heterothrix</i> sp. | LH10HG7061, LH08SG5052 | | |
| <i>Xanthonema bristolianum</i> (cf.) | LH08HW9018 | LH10HG6059 | |
| <i>Xanthonema exile</i> (cf.) | LH10HG7078 | | |
| <i>Xanthonema</i> sp. | LH10HG9058, LH10HG7029 | LH10HG3065, LH10HG9031, LH10HG1K69 | LH10HG8112, LH10HW9129 |
| <i>Eustigmatos</i> sp. | LH10HG9133, LH10HG5036 | | |

Table S4b. Distribution of the green algal species across the sampling sites.

| Species | Representative isolates (in phylogenetic trees) | Drill core (2008) SCH | Drill core (2008) ALB | Drill core (2008) HAI | Topsoil (2010) HAI | I | M | E |
|--|--|-----------------------------|-----------------------------|-----------------------------|--------------------------|---|---|---|
| <i>Auxenochlorella protothecoides</i> | LH10HG6096 | | | | HEG5/6/7 | | 1 | 1 |
| <i>Auxenochlorella</i> sp. | LH08AW4103 | | AEW4 | | | | 1 | |
| <i>Chlorella mirabilis</i> (cf.) | LH08AG9040 | | AEG9 | | | | | 1 |
| <i>Chlorella mirabilis</i> (cf.) | LH10HG6139 | | | | HEG6 | | 1 | |
| <i>Chlorella vulgaris</i> | LH08HG1081 | | | HEG1 | | 1 | | |
| <i>Chlorella vulgaris</i> | LH08SG3006 | SEG1/3 | | HEG2/4/5, HEW6/9 | HEG2/3/4/6/9 | 1 | 1 | 1 |
| <i>Chlorella vulgaris</i> (cf.) | LH10HG7072 (PS) | | | | HEG7 | | | 1 |
| <i>Chlorella vulgaris</i> (cf.) | LH10HG2081 (PS) | | | | HEG2 | 1 | | |
| <i>Chloroidium ellipsoideum</i> (cf.) | LH10HG9105 | | AEG8/9 | | HEG6/7/9 | | | 1 |
| <i>Chloroidium saccharophilum</i> | LH10HG7062 | | AEW8 | | HEG5/7 | | 1 | 1 |
| <i>Coccomyxa viridis</i> | SAG 2483 | | AEW8 | | | | | 1 |
| <i>Coccomyxa simplex</i> | LH08SG9051 (PS) | SEG9 | | | | | | 1 |
| <i>Coccomyxa</i> sp. | LH08AW1017 | | AEW1 | | | 1 | | |
| <i>Dictyochloropsis splendida</i> | LH08AW3050 | | AEW3 | | | 1 | | |
| <i>Diplosphaera</i> sp.(I) | LH08HW8075 | | | HEW8 | | | | 1 |
| <i>Diplosphaera</i> sp.(II) | LH08AG9089 | | AEG9 | | | | | 1 |
| <i>Lobosphaera bisecta</i> (cf.) | LH10HG3P15 (PS) | | | | HEG3 | 1 | | |
| <i>Lobosphaera irregularis</i> (cf.) | LH08AW3064 | SEW5 | AEW3 | | | 1 | 1 | |
| <i>Muriella terrestris</i> (cf.) | LH08SG3009 | SEG3 | | | HEG1/7/8/9 | 1 | | 1 |
| <i>Nannochloris</i> sp. | LH08SG8030 | SEG3/8 | | | HEG6 | 1 | 1 | 1 |
| <i>Navichloris fusiformis</i> (proposed taxon) | SAG 2477 | | AEW3 | | | 1 | | |
| <i>Neocystis brevis</i> | LH10HG9080 | SEG2 | AEG9, AEW8 | HEW6 | HEG4/9 | 1 | 1 | 1 |
| <i>Pseudostichococcus monallantoides</i> | LH10HG3045 | | AEG7 | | HEG2/3 | 1 | | 1 |
| <i>Pseudostichococcus</i> sp. | LH08SW8044 | SEW8 | | | | | | 1 |
| <i>Stichococcus</i> sp.(I) | LH08AW8025 | | AEG9/AEW8 | | | | | 1 |
| <i>Stichococcus</i> sp.(II) | LH08SG5057 | SEG5 | AEG1 | | HEG2/3/7/9 | 1 | 1 | 1 |
| <i>Stichococcus</i> sp.(III) | LH10HG6110, LH08AG7010 | | AEG7 | | HEG6 | | 1 | 1 |
| <i>Stichococcus</i> sp.(IV) | LH08SG1073 | SEG1 | | | | 1 | | |
| <i>Stichococcus</i> sp.(V) | LH08SW1099 | SEW1 | | | | 1 | | |
| <i>Stichococcus</i> sp.(VI) | SAG 2482 | | AEW8 | | | | | 1 |
| <i>Stichococcus</i> sp.(VII) | SAG 2481 | | AEW8 | | | | | 1 |
| Unidentified Chlorellaceae (I) | LH08AG1034 | | AEG1 | | HEG7 | 1 | | 1 |
| Unidentified Chlorellaceae (II) | LH10HG9020 | SEG2/3 | | | HEG2/3/7/9 | 1 | | 1 |
| Unidentified Chlorellaceae (III) | LH08SG3093 | SEG3 | | | | 1 | | |
| Unidentified Chlorellaceae (IV) | LH10HG709K | | | | HEG7 | | | 1 |
| <i>Xylochloris</i> sp. | SAG 2382 | | AEG7 | | | | | 1 |
| <i>Acutodesmus rubescens</i> | LH08SG8041 | SEG8 | | | | | | 1 |
| <i>Bracteacoccus cohaerens</i> | LH10HG9034 | | | | HEG9 | | | 1 |
| <i>Bracteacoccus cohaerens</i> (cf.) | LH08SG2015 | SEG2 | | | | 1 | | |
| <i>Chlamydomonas gerloffii</i> (cf.) | LH08SW5031 | SEW5 | | | | | 1 | |
| <i>Chlamydomonas rapa</i> | LH08SG1077 | SEG1/9 | | | | 1 | | 1 |
| <i>Chlamydomonas rapa</i> (cf.) | LH10HG1027 | | | | HEG1 | 1 | | |
| <i>Chlamydomonas typica</i> (cf.) | LH08SG9022 (PS) | SEG9 | | | | | | 1 |
| <i>Chlamydomonium vacuolatum</i> | LH10HG1013 | | | | HEG1 | 1 | | |
| <i>Chlorococcum minutum</i> (cf.) | SAG 2479 | | AEG7/AEW5 | | | | 1 | 1 |
| <i>Chlorococcum sphacosum</i> | LH10HG3113 | | | | HEG3 | 1 | | |
| <i>Coelastrella multistriata</i> | LH10HG7083 | | AEG2/AEW4 | | HEG7/8 | 1 | 1 | 1 |
| <i>Coelastrella</i> sp. | LH10HG2098 | | | | HEG1/2/5/6/7/9 | 1 | 1 | 1 |
| <i>Desmotetra stigmatica</i> | LH08SG2049 | SEG2 | | | HEG6 | 1 | 1 | |
| <i>Heterochlamydomonas</i> sp. | LH08AG2004 | | AEG2 | | | 1 | | |
| <i>Jenufa</i> sp. | SAG 2383 | | AEW8 | | | | | 1 |
| <i>Oogamochlamys</i> sp.(I) | LH08SG8047 | SEG8 | | | | | | 1 |
| <i>Oogamochlamys</i> sp.(II) | SAG 2476 | | AEW1 | | | 1 | | |

Table S4b. (continuation)

| Species | Representative isolates (in phylogenetic trees) | Drill core (2008) SCH | Drill core (2008) ALB | Drill core (2008) HAI | Topsoil (2010) HAI | I | M | E |
|--|--|-----------------------------|-----------------------------|-----------------------------|--------------------------|---|---|---|
| <i>Pseudomuriella aurantiaca</i> | LH10HG2039 | | | | HEG2/9 | 1 | 0 | 1 |
| <i>Stephanosphaeria</i> sp. | LH10HG6108 | | | | HEG6 | | 1 | |
| <i>Tatrensinia</i> sp.(I) | LH08SW7115 | SEW7 | | | | | | 1 |
| <i>Tatrensinia</i> sp.(II) | LH10HG7016 | | | | HEG7/9 | | | 1 |
| <i>Pseudendothia botryoides</i> (cf.) | LH08HW9058 | | | HEW9 | | | | 1 |
| <i>Pedinomonas minor</i> (cf.) | LH08SG2033 (PS) | SEG2 | | | | 1 | | |
| <i>Klebsormidium dissectum/elegans</i> (cf.) | LH08HW9106 | | AEG1/7/9 | HEW1/4/5/9 | | 1 | 1 | 1 |
| <i>Klebsormidium flaccidum</i> (cf.) | LH10HG2056 | SEG3 | | HEW5 | HEG2/7 | 1 | 1 | 1 |
| <i>Asterosiphon</i> sp. | LH10HG3064 | | | | HEG3 | 1 | | |
| <i>Botrydiopsis</i> sp. | LH08AW1076 | SEG2 | AEW1 | | | 1 | | |
| <i>Botrydiopsis callosa</i> (cf.) | LH08AW4043 | | AEW4 | | | | 1 | |
| <i>Heterococcus caespitosus</i> (cf.) | LH08AG2020 | | AEG2 | | | 1 | | |
| <i>Heterococcus chodatii</i> (cf.) | LH10HG9111 | | | | HEG9 | | | 1 |
| <i>Heterococcus</i> sp. | LH10HG9085, LH10HG2140 | | | | HEG2/9 | 1 | | 1 |
| <i>Heterothrix sessile</i> | LH10HG5079 | | | | HEG5/9 | | 1 | 1 |
| <i>Heterothrix</i> sp. | LH10HG7061, LH08SG5052 | SEG5 | | | HEG7 | | | 1 |
| <i>Xanthonema bristolianum</i> (cf.) | LH08HW9018 | | | HEW9 | HEG6 | | 1 | |
| <i>Xanthonema exile</i> (cf.) | LH10HG7078 | | | | HEG7 | | | 1 |
| <i>Xanthonema</i> sp. | LH10HG9058, LH10HG7029 | | | | HEG1/3/7/8/9, HEW9 | 1 | | 1 |
| <i>Eustigmatos</i> sp. | LH10HG9133, LH10HG5036 | | | | HEG5/9 | 1 | | 1 |

Legend. PS=partial sequence; Land-use intensity: E=extensive, M=managed, I=intensive; Exploratory: ALB=Schwäbische Alb, HAI=Hainich-Dün, SCH=Schorfheide-Chorin.

Table S5. General morphological characteristics of the new isolates.

| Representative isolates | Morphotype assignation | Cell shape | Cell size | Figure |
|-------------------------|--|--------------------------------|-------------------------------|-----------------------|
| LH10HG6096 | <i>Chlorella</i> | spherical | Ø=2.9-6.5 µm | Figure 7k |
| SAG 2478 | <i>Chlorella</i> | spherical | Ø=2.2-5.2 µm | Figure 7l |
| LH08AG9040 | <i>Chlorella</i> -like | spherical | Ø=3.0-6.4 µm | Figure 7a |
| LH10HG6139 | <i>Chlorella</i> -like | spherical | n/a | n/a |
| LH08HG1081 | <i>Chlorella</i> | spherical/ellipsoidal | Ø=2.3-5.2 µm | Figure 7b |
| LH08SG3006 | <i>Chlorella</i> | spherical/ellipsoidal | n/a | n/a |
| LH10HG2081 (PS) | <i>Chlorella</i> | spherical/ellipsoidal | Ø=2.5-6.2 µm | Figure 7c |
| LH10HG7072 (PS) | <i>Chlorella</i> | spherical/ellipsoidal | n/a | n/a |
| LH10HG9105 | <i>Chloroidium</i> | elliptical | l=4.8-7.8 µm, w=3.3-5.3 µm | Figure 6i |
| LH10HG7062 | <i>Chloroidium</i> | elliptical | l=5.6-9.1 µm, w=4.0-6.2 µm | Figure 6h |
| SAG 2483 | <i>Coccomyxa</i> | fusiform (asymmetrical) | l=4.7-8.4 µm, w=1.8-3.6 µm | Figure 6f |
| LH08SG9051 (PS) | <i>Coccomyxa</i> | fusiform (asymmetrical) | l=4.4-7.8 µm, w=2.0-4.3 µm | Figure 6e |
| LH08AW1017 | <i>Coccomyxa</i> | fusiform (asymmetrical) | n/a | n/a |
| LH08AW3050 | <i>Dictyochloropsis</i> | spherical | Ø=7.7-39.4 µm | Figure 6l |
| LH08HW8075 | <i>Diplosphaera</i> | broadly oval/cylindric | l=3.2-6.2 µm, w=1.9-4.4 µm | Figure 8b (Chapter 3) |
| LH08AG9089 | <i>Diplosphaera</i> | broadly oval/cylindric | n/a | n/a |
| LH10HG3P15 (PS) | <i>Lobosphaera</i> | spherical | Ø=6.2-9.6 µm | Figure 6j |
| LH08AW3064 | <i>Lobosphaera</i> | spherical | n/a | n/a |
| LH08SG3009 | <i>Chlorella</i> | spherical | Ø=1.7-4.2 µm | Figure 3b (Chapter 3) |
| LH08SG8030 | <i>Chlorella</i> | spherical | Ø=2.2-3.9 µm | Figure 7e (Chapter 3) |
| SAG 2477 | <i>Coccomyxa</i> -like | fusiform (symmetrical) | l=8.3-15.3 µm, w=3.6-7.8 µm | Figure 6g |
| LH10HG9080 | <i>Neocystis</i> | broadly oval | l=4.8-8.7 µm, w=2.7-6.6 µm | Figure 6d |
| LH10HG3045 | <i>Stichococcus</i> | cylindrical | n/a | n/a |
| LH08SW8044 | <i>Stichococcus</i> | cylindrical | l=3.4-9.8 µm; w=1.6-3.3 µm | Figure 8d (Chapter 3) |
| LH08AW8025 | <i>Stichococcus</i> | cylindrical | l=1.7-19.8 µm, w=1.4-3.6 µm | Figure 6a |
| LH08SG5057 | <i>Stichococcus</i> | cylindrical | l=4.5-9.8 µm, w=1.9-3.3 µm | Figure 6b |
| LH10HG6110, LH08AG7010 | <i>Stichococcus</i> | cylindrical | n/a | n/a |
| LH08SG1073 | <i>Stichococcus</i> | cylindrical | n/a | n/a |
| LH08SW1099 | <i>Stichococcus</i> / <i>Diplosphaera</i> | cylindrical/packages | l=3.0-6.3 µm; w=1.0-4.2 µm | Figure 8j (Chapter 3) |
| SAG 2482 | <i>Stichococcus</i> | cylindrical | l=3.8-7.2 µm, w=1.9-6.3 | Figure 6c |
| SAG 2481 | <i>Stichococcus</i> | cylindrical | l=3.1-8.7 µm; w=1.6-2.9 µm | Figure 8e (Chapter 3) |
| LH08AG1034 | <i>Chlorella</i> | spherical | Ø=2.4-4.5 µm | Figure 7g |
| LH10HG9020 | <i>Chlorella</i> | spherical | Ø=2.4-4.8 µm | Figure 7h |
| LH08SG3093 | <i>Chlorella</i> | spherical | Ø=2.3-3.8 µm | Figure 3d (Chapter 3) |
| LH10HG709K | <i>Chlorella</i> | spherical/ellipsoidal | l=3.1-5.1 µm, w=2.0-3.7 µm | Figure 7j |
| SAG 2382 | <i>Trebouxia</i> -like | spherical | l=4.7-12.9 µm, w=3.1-10.6 µm | Figure 6k |
| LH08SG8041 | <i>Scenedesmus</i> -like | ellipsoidal, obtuse cell poles | Ø=3.9-7.4 µm | Figure 3g |
| LH10HG9034 | <i>Bracteacoccus</i> -like | spherical | Ø=5.4-12.3 µm | Figure 3i |
| LH08SG2015 | <i>Bracteacoccus</i> -like | spherical | n/a | n/a |
| LH08SW5031 | <i>Chlamydomonas</i> | oval flagellated | l=10.9-16.4 µm, w=4.3-10.1 µm | Figure 4c |
| LH08SG1077 | <i>Chlamydomonas</i> | oval flagellated | n/a | n/a |
| LH10HG1027 | <i>Chlamydomonas</i> | oval flagellated | l=6.1-9.9 µm, w=2.7-5.1 µm | Figure 4d |
| LH08SG9022 (PS) | <i>Chlamydomonas</i> | oval flagellated | l=6.7-12.2 µm, w=4.0-8.0 µm | Figure 4f |

Table S5. (continuation)

| Representative isolates | Morphotype assignation | Cell shape | Cell size | Figure |
|-------------------------|----------------------------|--------------------------------|-------------------------------|--------------------------|
| LH10HG1013 | <i>Chlorococcum</i> | spherical/oval | n/a | n/a |
| SAG 2479 | <i>Chlorococcum</i> | spherical/oval | l=5.6-9.5 µm, w=5.6-9.5 µm | Figure 4k |
| LH10HG3113 | <i>Chlorococcum</i> | spherical/oval | l=5.8-9.3 µm, w=2.6-6.3 µm | Figure 4j |
| LH10HG7083 | <i>Scenedesmus</i> -like | ellipsoidal, obtuse cell poles | Ø=3.5-7.0 µm | Figure 3e |
| LH10HG2098 LH10HG7018 | <i>Scenedesmus</i> -like | ellipsoidal, obtuse cell poles | l=6.8-12.5 µm, w=7.0-10.9 µm | n/a |
| LH08SG2049 | <i>Chlorosarcina</i> -like | spherical | Ø=3.5-6.6 µm | Figure 3f Figure 4g |
| LH08AG2004 | <i>Chlamydomonas</i> | oval flagellated | l=6.0-10.5 µm, w=3.1-7.5 µm | Figure 4e |
| LH08AW8035 (= SAG 2383) | <i>Bracteacoccus</i> -like | spherical | Ø=2.9-7.3 µm | Figure 4l |
| LH08SG8047 | <i>Chlamydomonas</i> | oval flagellated | Ø=5.1-8.9 µm | Figure 4a |
| SAG 2476 (PS) | <i>Chlamydomonas</i> | oval flagellated | l=9.1-14.5 µm, w=5.1-10.0 µm | Figure 4b |
| LH10HG2039 | <i>Bracteacoccus</i> -like | spherical | Ø=3.6-8.2 µm | Figure 3h |
| LH10HG6108 | <i>Chlorococcum</i> | spherical/oval | n/a | n/a |
| LH08SW7115 | <i>Chlorococcum</i> | spherical/oval | l=8.3-17.8 µm, w=5.2-13.3 µm | Figure 4h |
| LH10HG7016 LH10HG913 | <i>Chlorococcum</i> | spherical/oval | Ø=7.9-13.1 µm | n/a Figure 4i |
| LH08HW9058 | <i>Pseudendocloniopsis</i> | sphaerical, obtuse cell pole | Ø=4.7-9.5 µm | Figure 3d |
| LH08SG2033 (PS) | <i>Chlamydomonas</i> -like | oval flagellated | l=2.9-5.6 µm, w=2.5-4.8 µm | Figure 3c |
| LH08HW9106 | <i>Klebsormidium</i> | trichal | l=4.2-12.8 µm, w=4.1-6.2 µm | Figure 3b |
| LH10HG2056 | <i>Klebsormidium</i> | trichal | l=6.4-19.7 µm, w=4.5-7.0 µm | Figure 3a |
| LH10HG3064 | <i>Asterosiphon</i> | spherical/siphonal | Ø=5.4-12.1 µm | Figure 11k |
| LH08AW1076 | <i>Botrydiopsis</i> | sphaerical | Ø=4.9-10.2 µm | Figure 11i |
| LH08AW4043 | <i>Botrydiopsis</i> | sphaerical | Ø=5.1-8.8 µm | Figure 11j |
| LH10HG9133, LH10HG5036 | <i>Eustigmatos</i> | spherical | Ø=6.1-13.1 µm | Figure 11l n/a |
| LH08AG2020 | <i>Heterococcus</i> | heterotrichal | n/a | n/a |
| LH10HG9111 | <i>Heterococcus</i> | heterotrichal | l=5.2-12.4 µm, w=4.1-13.2 µm | Figure 11g, h |
| LH10HG9085, LH10HG2140 | <i>Heterococcus</i> | heterotrichal | n/a | Figure 11f |
| LH10HG5079 | <i>Xanthonema</i> | trichal | n/a | n/a |
| LH10HG7061, LH08SG5052 | <i>Xanthonema</i> | trichal | l=10.2-25.4 µm, w=2.7-10.7 µm | Figure 11d Figure 11e |
| LH08HW9018 | <i>Xanthonema</i> | trichal | l=3.3-15.3 µm, w=2.9-5.7 µm | Figure 11c |
| LH10HG7078 | <i>Xanthonema</i> | trichal | l=4.2-11.2 µm, w=3.3-6.0 µm | Figure 11b |
| LH10HG9058, LH10HG7029 | <i>Xanthonema</i> | trichal | l=5.7-12.1 µm, w=3.4-6.5 µm | Figure 11a, n/a |

Legend. PS=partial sequence.

Table S6a. Diversity of morphospecies in soil drill cores.

| Soil horizon | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h | A-h |
|--------------------------------|-------|-------|-------|-------|-------|--------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Exploratory | H | H | H | H | H | H | A | A | A | A | A | A | S | S | S | S | S | S | S |
| Landscape/plot Nr. | G/1-3 | G/4-6 | G/7-9 | W/1-3 | W/4-6 | W/7-12 | G/1-3 | G/4-6 | G/7-9 | W/1-3 | W/4-6 | W/7-9 | G/1-3 | G/4-6 | G/7-9 | W/1-3 | W/4-6 | W/7-9 | W/7-9 |
| Land-use | E | M | I | E | M | I | E | M | I | E | M | I | E | M | I | E | M | I | I |
| Subsample | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX | MX |
| <i>Bracteacoccus</i> | - | - | - | 1 | 1 | - | 1 | 1 | - | - | - | - | 1 | 1 | - | - | - | - | - |
| <i>Characiochloridaceae</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Chlamydomonas</i> | 1 | 1 | - | - | 1 | 1 | - | - | 1 | 1 | - | 1 | 1 | 1 | 1 | - | 1 | 1 | 1 |
| <i>Chlorococcum</i> | 1 | 1 | 1 | - | 1 | 1 | - | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | - | 1 | - | - |
| <i>Chlorosarcinaceae</i> | - | - | 1 | - | - | - | 1 | - | 1 | - | - | - | - | 1 | - | - | - | - | - |
| <i>Monoraphidium terrestre</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Radiococcaceae</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Scenedesmeceae</i> | - | 1 | - | - | 1 | - | 1 | - | 1 | - | 1 | - | 1 | 1 | 1 | - | - | - | - |
| <i>Chlorella</i> | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | - |
| <i>Chloroidium</i> | - | - | - | - | - | 1 | - | - | - | - | 1 | - | - | 1 | - | - | - | - | - |
| <i>Coccomyxa</i> | - | 1 | - | 1 | 1 | 1 | - | - | - | 1 | - | 1 | 1 | 1 | 1 | - | - | - | - |
| <i>Dictyochloropsis</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Keratococcus</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Lobosphaera</i> | - | - | 1 | - | 1 | 1 | 1 | - | - | 1 | - | 1 | 1 | - | 1 | - | - | 1 | - |
| <i>Muriella</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Neocystis</i> | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Stichococcus</i> | - | 1 | - | 1 | 1 | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| <i>Dilabifilum</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Kentrosphaera</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Pseudendocloniopsis</i> | 1 | 1 | - | - | - | 1 | - | - | - | - | - | - | 1 | - | - | - | - | - | - |
| <i>Pseudendoclonium</i> | - | - | - | - | - | 1 | - | - | 1 | - | 1 | 1 | - | - | - | - | - | - | - |
| <i>Cosmarium</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Klebsormidium</i> | - | - | - | - | 1 | 1 | 1 | - | 1 | - | 1 | 1 | 1 | 1 | - | 1 | - | 1 | - |
| <i>Porphyridium</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Diadesmis</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Hantzschia</i> | 1 | 1 | 1 | - | 1 | - | - | - | - | - | - | - | 1 | - | 1 | - | - | - | - |
| <i>Mayamaea</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Navicula</i> | 1 | 1 | 1 | - | 1 | - | - | 1 | - | - | - | - | 1 | 1 | 1 | - | - | - | - |
| <i>Nitzschia</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Pinnularia</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Stauroneis</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Surirella</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Eustigmatos</i> | - | 1 | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - |
| <i>Heterococcus</i> | - | - | - | - | 1 | - | 1 | - | - | 1 | - | 1 | 1 | 1 | - | - | - | - | - |
| <i>Tribonema</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Xanthonema</i> | - | - | - | - | - | 1 | 1 | - | - | 1 | - | - | - | - | - | - | - | - | - |
| <i>Xanth. coccal</i> | 1 | 1 | 1 | - | 1 | - | 1 | - | 1 | - | - | 1 | 1 | 1 | - | 1 | - | - | - |
| <i>Cryptomonas</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>CylindrospERMUM</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Leptolyngbya</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Nostoc</i> | 1 | 1 | - | - | 1 | - | - | 1 | 1 | 1 | - | - | - | 1 | 1 | - | - | - | - |
| <i>Phormidium</i> | 1 | 1 | 1 | - | 1 | - | 1 | - | 1 | 1 | - | - | 1 | 1 | 1 | - | - | - | - |
| <i>Plectonema</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Pseudanabaena</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| <i>Stigonema</i> | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |

Legend. A-h=drill core; O-h=topsoil. H=Hainich-Dün; A=Schwäbische Alb; S=Schorfheide-Chorin. G=grassland; W=forest. E=extensive; M=managed; I=intensive. MX=mixed.

Table S6b. Diversity of morphospecies in topsoil samples.

| Soil horizon | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | O-h | |
|----------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|
| Exploratory | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | H | |
| Landscape/plot Nr. | G7 | G7 | G7 | G7 | G8 | G8 | G8 | G8 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | G9 | |
| Land-use | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | E | |
| Subsample | MD | NE | SW | MX | MD | NE | SW | MX | MD | NE | SW | MX | MD | NE | SW | MX | MD | NE | SW | MX | MD | NE | SW | MX |
| Bracteacoccus | - | 1 | - | 1 | - | - | 1 | - | 1 | - | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Characiochloridaceae | - | 1 | - | - | - | - | - | - | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | 1 | - | - | - | |
| Chlamydomonas | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| Chlorococcum | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | 1 | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | |
| Chlorosarcinaceae | - | - | - | 1 | - | - | - | - | 1 | 1 | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Monoraphidium terrestre | - | - | - | - | - | - | - | - | - | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | - | |
| Radiococcaceae | 1 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Scenedesmaceae | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | 1 | - | - | - | - | - | - | - | - | - | |
| Chlorella | - | 1 | 1 | - | 1 | 1 | - | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | - | |
| Chloroidium | - | - | - | - | 1 | - | 1 | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | - | |
| Coccomyxa | - | - | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | 1 | 1 | - | - | |
| Dictyochloropsis | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Keratococcus | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Lobosphaera | - | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | 1 | - | |
| Muriella | 1 | - | - | 1 | - | 1 | 1 | - | 1 | 1 | 1 | 1 | - | 1 | - | - | - | - | - | - | - | - | - | |
| Neocystis | - | - | - | - | 1 | - | - | - | 1 | - | 1 | - | - | - | - | - | - | - | - | 1 | - | - | - | |
| Stichococcus | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | 1 | 1 | 1 | - | 1 | - | - | - | |
| Dilabifilum | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Kentrosphaera | - | - | - | - | - | - | - | 1 | - | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Pseudendocloniopsis | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Pseudendoclonium | - | - | - | - | - | - | 1 | - | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Cosmarium | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Klebsormidium | - | 1 | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | |
| Porphyridium | - | - | - | - | - | - | - | - | - | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Diadesmis | - | 1 | - | 1 | - | - | 1 | - | 1 | - | 1 | - | - | - | - | - | - | - | - | - | - | - | - | |
| Hantzschia | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Mayamaea | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | 1 | - | |
| Navicula | - | - | 1 | - | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| Nitzschia | - | - | - | - | 1 | 1 | - | 1 | - | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Pinnularia | - | - | - | - | - | - | - | - | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Stauroneis | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Surirella | - | - | - | - | - | - | - | - | - | 1 | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Eustigmatos | - | 1 | - | - | - | - | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | 1 | - | - | - | - | - | - | - | |
| Heterococcus | 1 | 1 | - | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | |
| Tribonema | 1 | - | - | - | 1 | - | - | - | - | 1 | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Xanthonema | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | 1 | - | |
| Xanth. coccal | 1 | 1 | - | 1 | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | 1 | 1 | - | - | - | - | - | - | |
| Cryptomonas | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Cylindrospermum | - | - | - | 1 | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | |
| Leptolyngbya | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Nostoc | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Phormidium | 1 | - | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Plectonema | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Pseudanabaena | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |
| Stigonema | - | - | - | - | - | - | - | - | - | - | - | 1 | - | - | - | - | - | - | - | - | - | - | - | |

Legend. A-h=drill core; O-h=topsoil. H=Hainich-Dün; A=Schwäbische Alb; S=Schorfheide-Chorin. G=grassland; W=forest. E=extensive; M=managed; I=intensive. MX=mixed; MD=middle; NE=NorthEast; SW=SouthWest.

Table S7. Frequency of the morphospecies in soil drill cores.

| Morphotype and the authority | Nr. obs | % of samples | Microphotographs | Isolate/mixed culture |
|---|---------|--------------|----------------------------------|-----------------------|
| <i>Bracteacoccus</i> Tereg | 12 | 32 | Figure 3i | isolate |
| <i>Characiochloridaceae</i> Skuja | 6 | 16 | Supplementary Figure 1a-b | mixed culture |
| <i>Chlamydomonas</i> Ehrenberg | 32 | 84 | Figure 4a-f | isolate |
| <i>Chlorococcum</i> Meneghini | 31 | 82 | Figure 4h-k | isolate |
| <i>Chlorosarcinaceae</i> Bourrelly ex Groover & Bold | 8 | 21 | Supplementary Figure 1f | isolate |
| <i>Monoraphidium terrestre</i> (Bristol) Krienitz | 2 | 5 | Supplementary Figure 1h | mixed culture |
| <i>Radiococcaceae</i> Fott & Komárek | 1 | 3 | n/a | mixed culture |
| <i>Scenedesmeceae</i> Oltmanns | 21 | 55 | Supplementary Figure 1g | isolate |
| <i>Chlorella</i> Beijerinck | 24 | 63 | Figure 7a-c | isolate |
| <i>Chloroidium</i> Nadson | 6 | 16 | Figure 6h-i | isolate |
| <i>Coccomyxa</i> Schmidle | 24 | 63 | Figure 6e-g | isolate |
| <i>Dictyochloropsis</i> Geitler em. Tschermak-Woess | 1 | 3 | Figure 6l | isolate |
| <i>Keratococcus</i> Pascher | 11 | 29 | Supplementary Figure 1i-j | mixed culture |
| <i>Lobosphaera</i> Reisinger | 20 | 53 | Figure 6j | isolate |
| <i>Muriella</i> J. B. Petersen | 9 | 24 | Figure 7d-i | isolate |
| <i>Neocystis</i> F. Hindák | 6 | 16 | Figure 6d | isolate |
| <i>Stichococcus</i> Nägeli | 30 | 79 | Figure 6a-c | isolate |
| <i>Dilabifilum</i> Tschermak-Woess | 1 | 3 | Supplementary Figure 2b | mixed culture |
| <i>Kentrosphaera</i> Borzi | 4 | 11 | Supplementary Figure 1l | mixed culture |
| <i>Pseudendoctoniopsis</i> Vischer | 15 | 39 | Figure 3d | isolate |
| <i>Pseudendoctonidium</i> Wille | 9 | 24 | Supplementary Figure 2a | isolate |
| <i>Cosmarium</i> Corda | 1 | 3 | Supplementary Figure 2d-e | mixed culture |
| <i>Klebsormidium</i> Silva, Mattox & Blackwell | 23 | 61 | Figure 3a-b | isolate |
| <i>Porphyridium</i> Nägeli | 3 | 8 | Supplementary Figure 2h | mixed culture |
| <i>Diadisma</i> contenta (Grunow) D. G. Mann | 5 | 13 | Supplementary Figure 3b | mixed culture |
| <i>Hantzschia amphioxys</i> (Ehrenberg) Grunow | 18 | 47 | Supplementary Figure 3o | mixed culture |
| <i>Mayamaea</i> H. Lange-Bertalot | 13 | 34 | Supplementary Figure 3e | mixed culture |
| <i>Navicula</i> Bory de Saint-Vincent | 24 | 63 | Supplementary Figure 3a,c-d | mixed culture |
| <i>Nitzschia</i> Hassall | 6 | 16 | Supplementary Figure 3m-n | mixed culture |
| <i>Pinnularia</i> Ehrenberg | 4 | 11 | Supplementary Figure 3f-i | mixed culture |
| <i>Stauroneis</i> Ehrenberg | 1 | 3 | Supplementary Figure 3j | mixed culture |
| <i>Surirella</i> Turpin | 2 | 5 | Supplementary Figure 3l | mixed culture |
| <i>Eustigmatos</i> D. J. Hibberd | 10 | 26 | Figure 11l | isolate |
| <i>Heterococcus</i> Chodat | 24 | 63 | Figure 11f-h | isolate |
| <i>Tribonema</i> Derbès & Solier | 4 | 11 | Supplementary Figure 2j | mixed culture |
| <i>Xanthonema</i> Silva | 20 | 53 | Figure 11a-e | isolate |
| <i>Xanth. coccal</i> | 22 | 58 | Figure 11i-j; Suppl. Figure 2k-l | Isolate; mix.cul. |
| <i>Cryptomonas</i> Ehrenberg | 1 | 3 | Supplementary Figure 2i | mixed culture |
| <i>Cylindrocapsa</i> F. T. Kützing ex É. Bornet & C. Flahault | 2 | 5 | Supplementary Figure 4j | mixed culture |
| <i>Leptolyngbya</i> Anagnostidis & Komárek | 12 | 32 | Supplementary Figure 4b-e | mixed culture |
| <i>Nostoc</i> Vaucher ex Bornet & Flahault | 20 | 53 | Supplementary Figure 4g-i | mixed culture |
| <i>Phormidium</i> Kützing ex Gomont | 21 | 55 | Supplementary Figure 4f | mixed culture |
| <i>Plectonema</i> Thuret ex Gomont | 1 | 3 | n/a | mixed culture |
| <i>Pseudanabaena</i> Lauterborn | 1 | 3 | Supplementary Figure 4a | mixed culture |
| <i>Stigonema</i> Agardh ex Bornet et Flahault | 1 | 3 | Supplementary Figure 4l | mixed culture |

Appendix | Chapter 2

Supporting Tables

Table S1. Species of green algae recovered from creek biofilms.

Table S2. Green algal species recovered from creek biofilms and their main morphological features.

Table S3. Percentage identities (PID) to closest GenBank-relatives.

Table S4. Sequences representing additional groups and lineages shown as collapsed triangles in **Figures 1-3**.

Table S1. Species of green algae recovered from creek biofilms.

| Species | Isolate(s) studied in detail | Sequence accession numbers (original identifiers) | Additional isolates | Sequence accession numbers |
|---|----------------------------------|--|--|--|
| <i>Chlorella</i> sp. | RK52 D11-2 | KF144182 KF144172 | DB1-5, DB1-10 DB2-4, DB2-7 DB3-2, DB6-28 DB6-33, DB14-17 DB14-21, DB14-23 DB14-12, SAG 2391 | KF144171, KF144173 KF144177-KF144181 KF144174-KF144176 |
| <i>Coccomyxa</i> cf. <i>pringsheimii</i> | WB28 WB32 | KF144225 | | KF144223 KF144224 (DB14-14) |
| <i>Coccomyxa</i> cf. <i>simplex</i> | WB40 | KF144226 | | |
| <i>Elliptochloris subsphaerica</i> | WB5-D1e | KF144205 | WB27 | KF144204 |
| <i>Marvania</i> sp. | WB67 | KF144207 | | |
| <i>Muriella terrestris</i> | D6-DB2 | KF144209 | SAG 2390 | |
| <i>Neocystis</i> cf. <i>mucosa</i> | SAG 2405 | KF144212 (WB21) | | |
| <i>Stichococcus bacillaris</i> | WB13 WB74 | KF144231 KF144232 | | |
| <i>Stichococcus</i> cf. <i>deasonii</i> | WB38 | KF144233 | | |
| <i>Stichococcus mirabilis</i> | WB69 | KF144234 | | |
| <i>Stichococcus</i> sp.1 | DB6-27 WB65 SAG 2407 | KF144235 KF144236 KF144237 (WB68) | | |
| <i>Stichococcus</i> sp.2 | D4-2A | KF144238 | | |
| <i>Stichococcus</i> sp.3 | SAG 2408 | KF144239 (WB8) | | |
| <i>Stichococcus</i> sp.4 | SAG 2406 | KF144240 (WB47) | | |
| <i>Acutodesmus obliquus</i> | D22-6-2B | KF144164 | | |
| <i>Bracteacoccus aerius</i> -relative | SAG 2403 | KF144165 (WB18) | | |
| <i>Bracteacoccus</i> sp. | DB9-3 | KF144166 | | |
| <i>Chlamydomonas</i> sp. | RK68 | KF144168 | DB6-21 | KF144167 |
| <i>Chlamydomonium</i> sp. | SAG 2402 RK50 | KF144169 (RK41) KF144170 | | |
| <i>Chlorococcum sphacosum</i> | SAG 2398 | KF144183 (GRK6-DB1) | | |
| <i>Chlorococcum ellipsoideum</i> -relatives1 | GRK6-DB5 | KF144184 | | |
| <i>Chlorococcum ellipsoideum</i> -relatives2 | SAG 2400 GRK6-DB6 SAG 2401 | KF144186 (GRK7-WB4) KF144185 KF144187 (GRK7-WB5) | | |
| <i>Chlorococcum minutum</i> -relative | SAG 2399 | KF144188 (GRK6-DB3) | | |
| <i>Desmodesmus</i> cf. <i>armatus</i> | RK43 SAG 2395 | KF144195 KF144191 (D7-6-1B) | DB6-23, DB6-26 DB6-30, SAG 2389 | KF144192-KF144194 |
| <i>Monoraphidium terrestre</i> cf. <i>dybowskii</i> | SAG 2393 | KF144208 (D21-6-5B) | | |
| <i>Mychonastes</i> cf. <i>homosphaera</i> | RK48 | KF144210 | | |
| <i>Mychonastes</i> sp. | DB6-29 | KF144211 | | |
| <i>Pseudomuriella</i> cf. <i>schumacherensis</i> | RK3 | KF144227 | | |
| <i>Scenedesmaceae</i> sp. | RK49 | KF144228 | | |
| <i>Desmochloris</i> cf. <i>halophila</i> | SAG 2397 | KF144190 (DB1-9) | | |
| <i>Dilabifilum printzii</i> | WB41 | KF144201 | DB15-6, WB3 WB24, WB31 WB43, WB59 | KF144197, KF144200 KF144199, KF144196 KF144202, KF144203 |
| <i>Hazenia mirabilis</i> | SAG 2396 | KF144206 (D9-14B) | | |
| <i>Pseudendocloniopsis botryoides</i> | SAG 2394 DB6-19 | KF144213 (D2-6-1A) KF144219 | DB14-6 DB14-11 DB14-24 DB14-28 DB14-45 WB4 WB45 | KF144218 KF144214-KF144217 KF144221 KF144220 |
| <i>Pseudendoclonium akinetum</i> | SAG 2404 | KF144222 (WB20) | | |

Legend. The 74 unialgal isolates, their affiliation to a certain class, their strain designations and NCBI accession numbers of their 18S rRNA gene sequences are listed. The 45 strains used as exemplar for detailed examination are listed separately from those additional strains representing the same species. When an isolate was accessioned by the SAG culture collection, the corresponding SAG strain number is given and the original strain designation is in brackets next to its sequence accession number.

Table S2. Green algal species recovered from creek biofilms and their main morphological features.

| Species | Figure | Morphological features |
|--|---------------|--|
| <i>Chlorella</i> sp. | Figure 5c, 5d | Cells globose or ellipsoidal with smooth cell walls. Single parietal chloroplast with single pyrenoid. Reproduction by 2-8 autospores per cell, released by rupture of parental cell wall. Cell diameter 4.5-8.0 µm. |
| <i>Coccomyxa</i> cf. <i>pringsheimii</i> | Figure 5i | Heteropolar cells oval, fusiform or egg-shaped with smooth cell walls. Single parietal chloroplast without pyrenoids. Cell diameter: 4.4-8.0 µm. |
| <i>Coccomyxa</i> cf. <i>simplex</i> | n/a | Heteropolar cells oval, fusiform or egg-shaped with smooth cell walls. Single parietal chloroplast without pyrenoids. Cell length: 4.6-8.3 µm. |
| <i>Elliptochloris subsphaerica</i> | Figure 5h | Young cells cylindric, slightly inflexed, mature cells ellipsoidal to globose. Single parietal chloroplast with single pyrenoid. Autosporangia ellipsoidal with several cylindric autospores. Cell diameter 7.0-10.5 µm. |
| <i>Marvania</i> sp. | Figure 5f | Cells spherical, with 2 parietal chloroplasts, pyrenoids absent. Cell diameter 2.5-4.2 µm. |
| <i>Muriella terrestris</i> | Figure 5e | Cells spherical, with 2 or more parietal chloroplasts without a pyrenoid. Asexual reproduction by autospores, released by rupture of parental cell wall. Cell diameter: 2.4-4.4 µm. |
| <i>Neocystis</i> cf. <i>mucosa</i> | Figure 5g | Cells broadly oval with single parietal chloroplast without a pyrenoid. Asexual reproduction by autospores, 2-4 per autosporangium, released by rupture of the cell wall, remaining visible in the colony. Cell diameter: 2.9-5.7 µm. |
| <i>Stichococcus bacillaris</i> | Figure 5a | Few-celled fragmenting filaments or individual cells. Cells cylindrical or elongate, sometimes slightly oval with single parietal chloroplast without pyrenoids. Cell length: 4.1-17.4 µm. |
| <i>Stichococcus</i> cf. <i>deasonii</i> | n/a | Cylindrical elongate cells with a single parietal chloroplast containing a single starch-enveloped pyrenoid, cell length: 4.1-8.9 µm. |
| <i>Stichococcus mirabilis</i> | n/a | For description see <i>Stichococcus bacillaris</i> above, cell length: 3.2-9.5 µm. |
| <i>Stichococcus</i> sp.1 | n/a | For description see <i>Stichococcus bacillaris</i> above, cell length: 2.9-11.9 µm. |
| <i>Stichococcus</i> sp.2 | n/a | For description see <i>Stichococcus</i> cf. <i>deasonii</i> above, cell length: 6.2-11.4 µm. |
| <i>Stichococcus</i> sp.3 | n/a | For description see <i>Stichococcus</i> cf. <i>deasonii</i> above, cells sometimes flexed. Cell length: 6.2-11.4 µm. |
| <i>Stichococcus</i> sp.4 | Figure 5b | For description see <i>Stichococcus</i> cf. <i>deasonii</i> above, cell length: 3.7-9.5 µm. |
| <i>Acutodesmus obliquus</i> | Figure 6h | Fusiform cells with acute cell poles, without bristles. Cell length: 6.2-10.9 µm. |
| <i>Bracteacoccus</i> sp. | Figure 7b | Sphaerical cells with several to many parietal lens-shaped chloroplasts without pyrenoids. Cell diameter: 4.9-10.0 µm. |
| <i>Chlamydomonas</i> sp. | Figure 6f | Spherical cells with anterior contractile vacuole and two isokont anterior flagella. Cells contain single cup-shaped chloroplast with one basal pyrenoid and an eyespot at the cell anterior. Cell length: 7.2-11.1 µm. |
| <i>Chlorococcum</i> cf. <i>sphaerosum</i> | Figure 6a | Cells spherical to ellipsoidal with smooth cell walls. One parietal chloroplast per cell containing single pyrenoid. Biflagellated zoospores remain ellipsoidal after motility ceases (<i>Chlamydomonas</i> -type). Cell diameter: 7.0-10.5 µm. |
| <i>Chlorococcum ellipsoideum</i> -relatives1 | Figure 6c | Cells spherical to ellipsoidal with smooth cell walls. One parietal chloroplast per cell containing single pyrenoid. Cell diameter: 8.5-11.7 µm. |
| <i>Chlorococcum ellipsoideum</i> -relatives2 | Figure 6d, 6e | Spherical cells with anterior contractile vacuole, possessing two isokont anterior flagella. Single band-shaped chloroplast per cell with a single lateral pyrenoid and an eyespot at the cell anterior. Cell length: 5.8-11.8 µm. |
| <i>Desmodesmus</i> cf. <i>armatus</i> | Figure 6g | Ellipsoidal cells with obtuse cell poles and bristles. Cells form 4-8-celled coenobia. Cell length: 6.1-17.6 µm. |
| <i>Mychonastes</i> cf. <i>homosphaera</i> | Figure 7c | Spherical <i>Chlorella</i> -like cells with single chloroplasts without a pyrenoid. Cell diameter: 1.8-3.3 µm. |
| <i>Mychonastes</i> sp. | Figure 7b | For description see <i>Mychonastes</i> cf. <i>homosphaera</i> above, cell diameter: 1.8-3.3 µm. |
| <i>Pseudomuriella</i> cf. <i>schumacherensis</i> | Figure 7a | Spherical cells containing chloroplast (parietal saucer-shaped in young cells or divided into several segments in mature cells. Pyrenoids are absent. Cell diameter: 3.7-8.1 µm. |
| <i>Scenedesmeaceae</i> sp. | Figure 6i | For description see <i>Desmodesmus</i> cf. <i>armatus</i> above, cell length: 5.3-8.4 µm. |
| <i>Dilabifilum printzii</i> | Figure 7i | For description see <i>Desmodesmus</i> cf. <i>armatus</i> above, cell length: 5.0-10.1 µm. |
| <i>Hazenia mirabilis</i> | Figure 7e | Individual cells contain single parietal chloroplast with a pyrenoid. Thalli heterotrichous, consisting of uniseriate, irregularly branched filaments. Cell length: 7.6-21.4 µm. |
| <i>Pseudendocloniopsis botryoides</i> | Figure 7f, 7g | Globose cells form roundish, irregular packets or heterotrichous thalli. Cells with single parietal chloroplast and a pyrenoid. Cell diameter: 4.0-11.1 µm. |
| <i>Pseudendoclonium akinetum</i> | Figure 7h | Globose cells forming uniseriate filaments, irregularly branched. Cells containing one parietal chloroplast with one or two pyrenoids. Cell length: 5.6-10.2 µm. |

Table S3. Percentage identities (PID) to closest GenBank-relatives.

| Identification of taxa based on 18S rDNA | Strain(s) | Closest relative(s) accession number(s) | Percentage identity | Habitat |
|---|--------------------------------|--|------------------------|-------------------------------|
| <i>Acutodesmus obliquus</i> | D22-6-2B | FR865726 | 100 | freshwater |
| <i>Bracteacoccus</i> | DB9-3 | U63103 | 99.59 | air-borne dust |
| <i>Bracteacoccus aerius</i> -relative | SAG 2403 | U63101 | 98.59 | air-borne dust |
| <i>Chlamydomonas</i> sp. | RK68 | X53904 | 98.65 | freshwater |
| <i>Chlamydomonas</i> sp. | RK68 | JN903974 | 99.13 | freshwater |
| <i>Chlamydomonas</i> sp. | RK68 | AY781664 | 99.18 | freshwater |
| <i>Chlamydomodium vacuolatum</i> | RK50+SAG 2402 | M63001 | 98.77 | n/a |
| <i>Chlamydomodium vacuolatum</i> | RK50+SAG 2402 | GQ122367 | 99.82 | freshwater |
| <i>Chlorella</i> sp. | RK52, D11-2 | GQ487223 | 99.71 | freshwater |
| <i>Chlorella</i> sp. | RK52/D11-2 | FR865683, AY197628, FJ946888, DQ377324 | 99.90 | freshwater |
| <i>Chlorococcum</i> cf. <i>sphacosum</i> | SAG 2398 | JN968580/JN968584 | 99.94 | freshwater |
| <i>Chlorococcum ellipsoideum</i> -relative1 | GRK6-DB5 | U70586 | 98.45 | terrestrial (soil) |
| <i>Chlorococcum ellipsoideum</i> -relative2 | SAG 2401/SAG 2400 /GRK6-DB6 | U70586 | 98.33 | terrestrial (soil) |
| <i>Chlorococcum minutum</i> -relative | SAG 2399 | JN968585 | 97.65 | terrestrial (soil) |
| <i>Coccomyxa</i> cf. <i>pringsheimii</i> | WB28 | AY762603 | 99.61 | terrestrial (phycobiont) |
| <i>Coccomyxa</i> cf. <i>simplex</i> | WB40 | FJ648514 | 99.83 | freshwater |
| <i>Desmochloris</i> sp. | SAG 2397 | FM882216/AB049416 | 99.30 | freshwater |
| <i>Desmodesmus</i> cf. <i>armatus</i> | SAG 2395/RK43 | FR865727 | 99.71 | freshwater |
| <i>Dilabifilum printzii</i> | WB41 (PS) | KF144198 | 100 | freshwater |
| <i>Elliptochloris</i> cf. <i>subsphaerica</i> | WB5-D1e | FJ648515, FJ648518 | 99.94 | terrestrial (roof tile) |
| <i>Hazenia</i> sp. | SAG 2396 | AF387156 | 99.94 | freshwater |
| <i>Marvania</i> | WB67 | AF124336 | 97.83 | freshwater |
| <i>Marvania</i> | WB67 | EF440182 | 99.30 | terrestrial (permafrost) |
| <i>Marvania</i> | WB67 | AY195983 | 99.60 | freshwater |
| <i>Monoraphidium terrestre</i> cf. <i>dybowskii</i> | SAG 2393 | Y16939 | 99.54 | freshwater |
| <i>Muriella terrestris</i> | D6-DB2 | AB012845 | 99.94 | terrestrial (soil) |
| <i>Muriella terrestris</i> | D6-DB2 | AY195969 | 100 | freshwater |
| <i>Mychonastes</i> cf. <i>homosphaera</i> | RK48 | GU799581 | 99.42 | freshwater |
| <i>Mychonastes</i> sp. | DB6-29 | AY543066 | 99.94 | freshwater |
| <i>Neocystis</i> cf. <i>mucosa</i> | SAG 2405 | HQ287929 | 99.56 | terrestrial (soil) |
| <i>Neocystis</i> cf. <i>mucosa</i> | SAG 2405 | HM565928 | 99.71 | n/a |
| <i>Pseudendocloniopsis botryoides</i> | SAG 2394 | AJ416102 | 99.76 | terrestrial (soil) |
| <i>Pseudendocloniopsis botryoides</i> | SAG 2394 | FR865755 | 99.94 | freshwater |
| <i>Pseudendoclonium akinetum</i> | SAG 2404 | AM087962 | 99.81 | freshwater |
| <i>Pseudendoclonium akinetum</i> | SAG 2404 | AM109906 | 99.75 | freshwater |
| <i>Pseudendoclonium akinetum</i> | SAG 2404 | DQ011230 | 99.94 | freshwater |
| <i>Pseudendoclonium basilense</i> | D9-14B | Z47996 | 99.82 | freshwater |
| <i>Pseudomuriella</i> cf. <i>schumacherensis</i> | RK3 | FR865690/X91268 | 99.01 | freshwater/terrestrial (soil) |
| <i>Pseudomuriella</i> cf. <i>schumacherensis</i> | RK3 | HQ292768 | 99.54 | terrestrial (epiphytic) |
| <i>Stichococcus bacillaris</i> | WB74 | AJ311637 | 99.88 | freshwater |
| <i>Stichococcus bacillaris</i> | WB13 | AJ311637 | 100 | freshwater |
| <i>Stichococcus</i> cf. <i>deasonii</i> | WB38 | DQ275460 | 99.74 | terrestrial (soil) |
| <i>Stichococcus mirabilis</i> | WB69 | AJ311638 | 100 | n/a |
| <i>Stichococcus</i> sp.1 | DB6-27 | AB055866 | 99.71 | n/a |
| <i>Stichococcus</i> sp.1 | DB6-27, WB65 | AB055866 | 100 | n/a |
| <i>Stichococcus</i> sp.2 | D4-2A | DQ275460 | 99.48 | terrestrial (soil) |
| <i>Stichococcus</i> sp.3 | SAG 2408 | DQ275460 | 98.60 | terrestrial (soil) |
| <i>Stichococcus</i> sp.4 | SAG 2406 | DQ275460 | 99.01 | terrestrial (soil) |

Legend. For each species, their closest available neighbouring sequences and pairwise sequence similarity (isolate/neighbouring sequence; in percentage calculated from p-distances) and the habitat from which the closest neighbours have been isolated are listed. The species are in two groups, i.e. one with high similarities with neighbouring sequences (99.9 and 100.0 %) so that they represent the same species as the isolate, the other with lower sequence similarities so that the neighbouring sequences do not represent the same species as the isolate. "n.a." represents for "not applicable" where origin is not known.

Table S4. Sequences representing additional groups and lineages shown as collapsed triangles in **Figures 1-3**.

| Figure | Taxon | GenBank accession number(s) |
|----------|---|---|
| Figure 1 | Chlorodendrophyceae | X68484, X70802 |
| | Chlorophyceae | DQ078295, AY779858, M62861, M32703, COU83125 |
| | <i>Choricystis/Botryococcus</i> | AY195970, AY762605, AY197640, AJ581912 |
| | <i>Desmococcus</i> | AJ431571, EU434017 |
| | <i>Hegewaldia</i> | FM205843, FM205844 |
| | <i>Lobosphaera</i> clade | AB006051, EU878372, EU878374 |
| | Oocystaceae | AF387154, AF228686, AF228691, AF228690, AF228689 |
| | <i>Parachlorella</i> -clade | AB176664, AB037085, Y17470, X56105, FM205845 |
| | <i>Prasiola/Prasiolopsis/Trichophilus</i> | AJ416106, EF200524, AY762600, AY762601 |
| | <i>Pseudochlorella/Koliella</i> | AB006050, X63520, U18520, AF514410 |
| | <i>Pseudomarvania</i> | AB087559, FJ896222 |
| | Trebouxiales clade | Z28971, M62995, AB080310, Z21551 |
| | Ulvophyceae | AB183610, AY278217 |
| | <i>Watanabea</i> clade | AJ439401, X73991, FM946000, EU346910, X73998, EF595524 |
| Figure 2 | <i>Arenicolinia</i> | AB218701, AY271673, AF513371 |
| | Chaetopeltidales | U83126, U83125, U83124 |
| | Chaetophorales | AF182820, AF182816, AF182821, D86499 |
| | <i>Characiosiphonia</i> | AF395437, AF395435 |
| | Chlorodendrophyceae | X68484, X70802 |
| | <i>Chlorogonia</i> | AJ410443, U70589 |
| | <i>Chloromonadinia</i> | U57696, AJ410448, U57697 |
| | <i>Crucicarteria</i> | D86501, U70595 |
| | <i>Dunaliellinia</i> | EU589200, DQ009743 |
| | <i>Golenkinia</i> | AF499924, AF499923 |
| | <i>Hafniomonas</i> | AB248251, D86496 |
| | <i>Jenufa</i> | HM563743, HM563744, AB257660, AB257659 |
| | <i>Moewusinia</i> | U70782, AY220571 |
| | <i>Monadinia</i> | AF514404, U57694 |
| | Nephroselmidophyceae | X75565, X74754 |
| | Oedogoniales | DQ018735, U83135, U83132 |
| | <i>Oogamochlamydia</i> | AJ410461, AJ410464, AJ410466, AB175836, AB175837 |
| | <i>Phacotinia</i> | AY009898, AF395438 |
| | <i>Polytomina</i> | U22941, AJ781310, U13984, U13985 |
| | <i>Radicarteria</i> | AF182817, AF182819 |
| | Sphaeropleales | see Fig.3 - Chlorophyceae which are not included in collapsed triangles |
| | <i>Tatrensinia</i> | AF514407, EF023280, AF514411 |
| | Trebouxiphyceae | AY762602, AB017435 |
| | <i>Treubarinia</i> | AY008844, U73471, U73474, AF277651 |
| | Ulvophyceae | AY278217, AB183610 |
| Figure 3 | Chaetopeltidales | U83126, U83125, U83124 |
| | Chaetophorales | AF182820, AF182816, AF182821, D86499 |
| | Chlamydomonadales | see Fig.2 - Chlorophyceae which are not included in collapsed triangles |
| | Chlorodendrophyceae | X68484, X70802 |
| | <i>Golenkinia</i> | AF499924, AF499923 |
| | Hydrodictyceae | AY663044, AY663041, AY779858, M62997 |
| | <i>Jenufa</i> | HM563743, HM563744, AB257660, AB257659 |
| | Neochloridaceae | AF288361, AF288364, M62861, AJ581914, M63000 |
| | Nephroselmidophyceae | X75565, X74754 |
| | Oedogoniales | DQ018735, U83135, U83132 |
| | Sphaeropleaceae | AF302770, AJU73469 |
| | Trebouxiphyceae | AB017435, AY762602 |
| | Ulvophyceae | AY278217, AB183610 |

Appendix | Chapter 3

Supporting Figures

Figure S1. ITS2 secondary structure analysis of *Chlorella vulgaris* and *C. pituita*.

Figure S2. Analysis of compensatory base changes (CBCs) in ITS2 sequences of *Chlorella*-like species.

Figure S3. 18S ML phylogeny of *Stichococcus*-like species.

Figure S4. ITS2 secondary structure analysis of *Stichococcus*-like species.

Figure S5. Neighbor-joining tree of *Stichococcus*-like species based on ITS2 sequences/secondary structures.

Figure S6. Microphotographs of representatives of the *Prasiola* clade.

Supporting Tables

Table S1. List of all analyzed accessions (isolates, strains and clones) of the polar *Chlorella*-like species and relatives.

Table S2. List of all analyzed accessions (isolates, strains and clones) of the *Stichococcus*-like species and relatives.

Table S3. List of the analyzed *Stichococcus*-like clades and their representative strains.




Figure S1. ITS2 secondary structure analysis of *Chlorella vulgaris* and *C. pituita*. (a) Polymorphic nucleotide sites detected in 13 ribotypes of *C. vulgaris* (ribotype A = SAG 211-11b) and in *C. pituita*. (b) ITS2 secondary structure model of *C. vulgaris* SAG 211-11b. Compensatory base changes (CBCs) between *C. vulgaris* and *C. pituita* (a, b, c, e) are marked by black triangles. Additional CBCs (d, f) among intraspecific ribotypes of *C. vulgaris* are shown as well. Bold letters represent nucleotide sites conserved in *C. vulgaris* and *C. pituita*.




Figure S2. Analysis of compensatory and hemi-compensatory base changes (CBCs, hCBCs) in ITS2 sequences of *Chlorella*-like species. (a) Neighbor-net analysis of *Chlorella vulgaris* SAG 211-11b (ribotype A), other *C. vulgaris* ribotypes (B-O) and *C. pituita* ACO1 311 (Fig. S1a). The numbers close to the solid lines represent bootstrap support values; the numbers close to the dashed lines represent CBCs between two sequences (the respective helices (Fig. S1b) are given by Roman numerals). (b)-(d) Consensus secondary structures of closely related accessions of the *Chlorellaceae*: (b) *Chlorella* sp.1 and *C. sp.2*; neighbor-joining tree (based on ITS2 sequences and secondary structures) and consensus ITS2 secondary structure of accessions closely related to the polar strains. (c) *Muriella terrestris*. (d) *Marvania* relatives. Nucleotide positions (represented as dots) within the ITS2 secondary structures are colored as follows: grey dots=conserved positions, orange dots=nucleotide substitutions and black dots=nucleotide deletions. Grey numbers between sequences in the NJ-trees give nucleotide differences and compensatory base changes (CBCs), respectively.




Figure S3. 18S ML phylogeny of *Stichococcus*-like species. Sequences of authentic strains are marked by a '<' sign. The numbers next to branches indicate statistical support values (maximum-likelihood bootstraps (ML)/Bayesian posterior probabilities (BI)); the clades of particular interest were additionally tested via maximum parsimony (MP) and bio-neighbor-joining (NJ) and the statistic support values are given in the following order: ML/BI/MP/NJ. Assignations into operational taxonomic units are based on sequence similarities $\geq 99.5\%$ (= OTU) and $\geq 99.0\%$ (= OTU).




Figure S4. ITS2 secondary structure analysis of *Stichococcus*-like species. (a) Polymorphic nucleotide sites detected in 33 ribotypes of *Stichococcus*-like species and allies. (b) ITS2 secondary structure model of *S. bacillaris* SAG 379-1b. Bold letters represent nucleotide sites which are conserved across all 56 analyzed ITS2 accessions.




Figure S5. Neighbor-joining tree of *Stichococcus*-like species based on ITS2 sequences/secondary structures. Thick lines indicate bootstrap support values ≥ 80 . Compensatory base changes (CBCs) detected within clades are given on branches as white numbers in black circles; the grey numbers above CBCs give nucleotide differences detected within clades. The polar and tropical accessions are blue and green colored, respectively.




Figure S6. Microphotographs of representatives of the *Prasiola* clade. (a) '*Chlorella*' *mirabilis* SAG 38.88; (b) '*Chlorella*' *sphaerica* SAG 11.88; (c) *Diplosphaera* sp. SAG 2049; (d) '*Desmococcus*' *spinocystis* SAG 2067; (e) *Desmococcus olivaceus* SAG 1.92; (f) *Prasiolopsis ramosa* SAG 26.82; (g) *Prasiola crista* SAG 43.96; (h) *Pseudomarvania ampulliformis* SAG 2047; (i) *Stichococcus jenerensis* SAG 2138; (j) *Pseudostichococcus monallantoides* SAG 380-1; (k) *Stichococcus fragilis* SAG 379-4; (l) *Stichococcus deasonii* SAG 2139.

Table S1. List of all analyzed accessions (isolates, strains and clones) of the polar *Chlorella*-like species and relatives.

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|-------------------------|-----------------------------|----------------|------------|------------|----------------------------------|---------------------------|--|
| <i>Chlorella</i> s.str. | <i>Chlorella pituita</i> | ACOI 311 | GQ176852 | GQ176853 | - | PT (Mira) | freshwater (trout nursery) |
| <i>Chlorella</i> s.str. | <i>Chlorella pituita</i> | ACOI 856 | FM205855 | FM205856 | - | PT (Manteigas) | freshwater |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> | 3GSG1R K22 | this study | n/a | N50°52'0.000" E10°50'0.000" | DE (Castle Gleichen) | subaerial (epilithic; sandstone) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> | CCAP 211/79 | FR865683 | FR865683 | - | UK (Edinburgh) | waste solvent biofilter |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> | DB1-10 | KF144171 | this study | - | DE (Deinschwanger Bach) | semi-terrestrial (creek biofilm) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> | HEG7SWWB K4396 | this study | n/a | N51°16'24.897" E10°24'37.485" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> | L1 | this study | n/a | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> | L5 | this study | n/a | N79°08'0.000" W80°30'0.000" | CA (Ellesmere Island) | soil (close to river) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> | NJ-7 | DQ377323 | n/a | - | AQ (Zhongshan Station) | terrestrial (wet rocks) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> A | CCAP 211/11B | n/a | AY591507 | - | NL (Delft) | freshwater |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> A | CCAP 254/5 | FR865696 | FR865697 | - | US (Bloomington, Indiana) | freshwater (aquarium) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> A | SAG 211-11b | FM205832 | FM205832 | - | NL (Delft) | freshwater |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | CCAP 211/110 | FN298918 | FN298918 | - | US (n/a) | freshwater (<i>Paramecium</i> symbiont) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | CCAP 211/74 | FR865682 | FR865682 | - | UK (Cumbria) | freshwater (Esthwaite Water) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | CCAP 211/80 | FM205853 | FM205853 | - | DE (Elsnigk) | freshwater (Molkerteich) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | CCAP 211/81 | FM205854 | FM205854 | - | DE (Trebichau) | freshwater (Salzteich) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH08HG4032 | this study | this study | N51°6'48.104" E10°26'10.249" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH08HG4088 | this study | this study | N51°6'48.104" E10°26'10.249" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH08HG5074 | this study | this study | N51°12'57.220" E10°19'21.096" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH08HW9094 | this study | this study | N51°7'48.871" E10°22'52.139" | DE (Hainich-Dün) | soil (forest) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH08SG1071 | this study | this study | N53°5'14.712" E13°58'10.717" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH08SG3006 | this study | this study | N53°6'10.204" E13°59'8.519" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH10HG2049 | this study | this study | N51°0'2.696" E10°25'48.036" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH10HG6014 | this study | this study | N51°12'53.766" E10°23'28.395" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH10HG6019 | this study | this study | N51°12'53.766" E10°23'28.395" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | LH10HG9075 | this study | this study | N51°13'26.031" E10°22'50.834" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> B | NB-1 | n/a | KC840685 | - | CN? | n/a |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> E | SAG 9.88 | n/a | AY591500 | - | ES (Madrid) | freshwater (waste-water) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> F | CCAP 211/11P | FR865658 | FR865658 | - | SE (Lund) | freshwater (pond in park) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> F | SAG 211-11p | n/a | AY591505 | - | SE (Lund) | freshwater (pond in park) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> G | TW-1 | n/a | KC840684 | - | CN(?) | n/a |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> H | CCAP 211/82 | FM205855 | FM205855 | - | DE (Micheln) | freshwater |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> H | LH08HG5082 | this study | this study | N51°12'57.220" E10°19'21.096" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> H | LH10HG6052 | this study | this study | N51°12'53.766" E10°23'28.395" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> I | CCAP 211/63 | FR865681 | FR865681 | - | UK (Cambridge) | freshwater (Peterhouse Ditch) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> I | LH08HG1081 | this study | this study | N50°58'17.934" E10°24'19.306" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> J | CCAP 211/109 | FN298917 | FN298917 | - | US | freshwater (<i>Paramecium</i> symbiont) |

Table S1. (continuation)

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|---------------------------|-------------------------------------|-----------------|------------|------------|----------------------------------|-------------------------|---|
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> K | D2 | JX185297 | JX185298 | - | CN(?) | n/a |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> L | LH10HG2067 | this study | this study | N51°0'2.696" E10°25'48.036" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> L | LH10HG2081 | this study | this study | N51°0'2.696" E10°25'48.036" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> M | L4 | this study | this study | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> N | L6 | this study | this study | N79°08'0.000" W80°30'0.000" | CA (Ellesmere Island) | soil (moraine) |
| <i>Chlorella</i> s.str. | <i>Chlorella vulgaris</i> O | LH10HG1069 | this study | this study | N50°58'17.934" E10°24'19.306" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp. | KNUA034 | KM243327 | KM243327 | - | AQ | freshwater |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp.2 | L3 | this study | this study | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp.1 | N9 | this study | this study | N79°58'0.000" E11°21'0.000" | NO (Svalbard) | soil (deglaciated) |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp. | NDem 9/21 | AY197628 | n/a | - | US | freshwater |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp. | NJ-18 | DQ377324 | n/a | - | AQ (Zhongshan Station) | terrestrial (wet rocks) |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp. | NMX37N | JF767012 | n/a | N41.48 E112.63 | CN? | freshwater |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp. | RK52 | KF144182 | n/a | - | DE (Deinschwanger Bach) | freshwater biofilm |
| <i>Chlorella</i> sp. | <i>Chlorella</i> sp. | VPL9-6 | FJ946888 | n/a | - | AQ (Trinity Peninsula) | freshwater |
| <i>Chlorella</i> sp. | Chlorellaceae | MCWWW9 | KP204587 | KP204587 | - | CA (Nova Scotia) | freshwater (wastewater) |
| <i>Chlorella</i> sp. | Chlorellaceae | N505T04 | GU942201 | GU942201 | - | CN (South China Sea) | marine |
| <i>Chlorella</i> sp. | <i>Micractinium</i> <i>inerimum</i> | NLP-F014 | KF597304 | KF597304 | - | KR | freshwater (wastewater) |
| Chlorellaceae | Chlorellaceae | L24 | this study | n/a | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Hindakia</i> | <i>Hindakia fallax</i> | CCAP 222/29 | GQ487223 | GQ487223 | - | - | - |
| <i>Hindakia</i> | <i>Hindakia tetrachotoma</i> | CCAP 222/80 | GQ487233 | GQ487233 | - | - | - |
| <i>Micractinium</i> | <i>Micractinium pusillum</i> | CCAP 248/5 | FM205836 | FM205836 | - | - | - |
| <i>Muriella</i> | Chlorophyta | clone WIM107 | AM114820 | n/a | - | - | - |
| <i>Muriella</i> | <i>Muriella</i> sp. | AS 2-4 | AY195969 | n/a | - | - | - |
| <i>Muriella</i> | <i>Muriella terrestris</i> | AEG6B K4797 | this study | n/a | N48°23'52.818" E9°20'31.152" | DE (Schwäbische Alb) | soil (grassland) |
| <i>Muriella</i> | <i>Muriella terrestris</i> | ASIB V38 | AB012845 | n/a | - | - | - |
| <i>Muriella</i> | <i>Muriella terrestris</i> | clone Ant 8/104 | n/a | JN653521 | - | AQ | soil (permafrost) |
| <i>Muriella</i> | <i>Muriella terrestris</i> | D6-DB2 | KF144209 | n/a | - | - | - |
| <i>Muriella</i> | <i>Muriella terrestris</i> | L20 | this study | this study | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Muriella</i> | <i>Muriella terrestris</i> | LH08SG3009 | this study | this study | N53°6'10.204" E13°59'8.519" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Muriella</i> | <i>Muriella terrestris</i> | LH10HG9077 | this study | n/a | N51°13'26.031" E10°22'50.834" | DE (Hainich-Dün) | soil (grassland) |
| <i>Muriella</i> | <i>Muriella terrestris</i> | N5 | this study | this study | S67°34'0.000" W68°08'0.000" | AQ (Adelaide Island) | soil (rookeries) |
| <i>Nannochloris</i> -like | <i>Marvania geminata</i> | SAG 12.88 | AF124336 | KM020037 | - | - | - |
| <i>Nannochloris</i> -like | <i>Chlorella</i> sp. | 193-GA188 | EU282456 | n/a | - | RU (Siberia, Kolyma) | soil (permafrost) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | clone Ant 8/117 | n/a | JN653529 | - | AQ | soil (permafrost) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative | HEG9 B K25502 | this study | n/a | N51°13'26.031" E10°22'50.834" | DE (Hainich-Dün) | soil (grassland) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | L13 | this study | n/a | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | L14 | this study | n/a | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | L15 | this study | n/a | S67°36'0.000" W68°15'0.000" | AQ (Anchorage Island) | soil (bare ground) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | L23 | this study | n/a | S67°32'0.000" W68°07'0.000" | AQ (Killingbeck Island) | endozoic (gut content; <i>Cryptopygus</i>) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | L32 | this study | n/a | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |

Table S1. (continuation)

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|---------------------------|---|-------------|------------|------------|---------------------------------|-------------------------|------------------|
| <i>Nannochloris</i> -like | <i>Marvania</i> relative2 | LH08AG1034 | this study | this study | N48°23'52.818" E9°20'31.152" | DE (Schwäbische Alb) | soil (grassland) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | LH08SG2053 | this study | n/a | N53°5'21.505" E13°58'48.169" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | LH08SG3078 | this study | this study | N53°6'10.204" E13°59'8.519" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | LH08SG3093 | this study | this study | N53°6'10.204" E13°59'8.519" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Nannochloris</i> -like | <i>Marvania</i> relative1 | LH10HG2094 | this study | n/a | N51°0'2.696" E10°25'48.036" | DE (Hainich-Dün) | soil (grassland) |
| <i>Nannochloris</i> -like | <i>Marvania</i> sp. | WB67 | KF144207 | n/a | - | - | - |
| <i>Nannochloris</i> -like | <i>Nannochloris</i> <i>coccoides</i> | CCAP 251/1b | AB080301 | n/a | - | - | - |
| <i>Nannochloris</i> -like | <i>Nannochloris</i> sp. | An-1 | EF440182 | n/a | - | - | - |
| <i>Nannochloris</i> -like | <i>Nannochloris</i> sp. | JL 4-6 | AY195983 | n/a | - | - | - |

Legend. AQ=Antarctica, CA=Canada, CN=China, DE=Germany, ES=Spain, KR=South Korea, NL=Netherlands, NO=Norway, PT=Portugal, RU=Russia, SE=Sweden, UK=United Kingdom, US=United States of America.

Table S2. List of all analyzed accessions (isolates, strains and clones) of the *Stichococcus*-like species and relatives.

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|----------------------------|------------------------------------|-----------------|------------|------------|----------------------------------|------------------------------|--|
| <i>Chlorella mirabilis</i> | <i>Chlorella mirabilis</i> | KFFB12-1 | this study | this study | S4°6'49.852" W78°58'1.012" | EC (Bombuscaro) | subaerial (litter) |
| <i>Chlorella mirabilis</i> | <i>Chlorella mirabilis</i> | L10 | this study | n/a | S62°10'0.000" W58°30'0.000" | AQ (King George Island) | soil (deglaciated) |
| <i>Chlorella mirabilis</i> | <i>Chlorella mirabilis</i> | LH10HG6139 | this study | n/a | N51°12'53.766" E10°23'28.395" | DE (Hainich-Dün) | soil (grassland) |
| <i>Chlorella mirabilis</i> | <i>Chlorella mirabilis</i> | Andreyeva 748-I | X74000 | n/a | - | RU | tundra |
| <i>Chlorella mirabilis</i> | <i>Chlorella mirabilis</i> | LH8AG9040 | this study | this study | N48°23'40.815" E9°30'10.053" | DE (Schwäbische Alb) | soil (grassland) |
| <i>Chlorella mirabilis</i> | <i>Chlorella mirabilis</i> | SEG5B K5318 | this study | n/a | N53°6'26.830" E14°0'1.885" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Desmococcus</i> | <i>Desmococcus endolithicus</i> | clone EN5JG | KR054372 | n/a | - | - | - |
| <i>Desmococcus</i> | <i>Desmococcus endolithicus</i> | SAG 25.92 | EU434026 | this study | - | AQ (Marie Bird Land) | subaerial (chasmoeendolithic) |
| <i>Desmococcus</i> | <i>Desmococcus olivaceus</i> | SAG 1.92 | EU434017 | KM020049 | - | AT (Vienna) | subaerial (epixylic; tree bark) |
| <i>Diplosphaera</i> | <i>Stichococcus</i> sp. | BCP-NB2-VF10 | AF513370 | n/a | - | - | - |
| <i>Diplosphaera</i> | <i>Stichococcus</i> sp. | BCP-SRS2-VF14 | AY377441 | n/a | - | - | - |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> | clone rtCF18sti | EF591011 | n/a | - | - | freshwater |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> | MBIC10465 | AB183601 | n/a | - | - | - |
| <i>Diplosphaera</i> | <i>Diplosphaera chodatii</i> | FontaineM3-1 | n/a | JX645020 | - | AT (Waldaist) | subaerial (<i>Dermatocarpus</i> symbiont) |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | A07020 | n/a | HM237336 | - | CN (Shapotou desert) | subaerial (<i>Endocarpus</i> symbiont) |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | AEW9B K460 | this study | n/a | N48°22'09.600" E9°24'54.800" | DE (Schwäbische Alb) | soil (forest) |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | Z07020 | n/a | HM237335 | - | CN (Shapotou desert) | subaerial (<i>Endocarpus</i> symbiont) |
| <i>Diplosphaera</i> | <i>Stichococcus chlorellioides</i> | BCP-CNP2-VF11B | AY271675 | n/a | - | - | - |
| <i>Diplosphaera</i> | <i>Stichococcus</i> sp. | clone QE28 | FJ790654 | n/a | - | - | - |
| <i>Diplosphaera</i> | <i>Chlorella sphaerica</i> | SAG 11.88 | AJ416105 | this study | - | NZ (Wawaira Scenic Reserve) | subaerial (phycobiont) |
| <i>Diplosphaera</i> | <i>Stichococcus chodatii</i> | UTEX 1177 | AB055867 | n/a | - | - | - |
| <i>Diplosphaera</i> | <i>Diplosphaera chodatii</i> | UTEX 1177 | AB055867 | HQ129931 | - | US (Texas) | soil (blackland prairie soil) |
| <i>Diplosphaera</i> | <i>Diplosphaera mucosa</i> | SAG 48.86 | KF673370 | n/a | - | AQ (Princess Elizabeth Land) | subaerial (moss) |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | J4028B | JN573864 | n/a | - | - | - |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | KS120SM6L | this study | this study | S3°58'31.436" W79°4'17.341" | EC (San Francisco) | subaerial (epiphytic; leaves) |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | KS135SM6L | n/a | this study | S3°58'31.436" W79°4'17.341" | EC (San Francisco) | subaerial (epiphytic; leaves) |

Table S2. (continuation)

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|--------------------------------|---|----------------|------------|------------|----------------------------------|-------------------------|--|
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | LH08AG9089 | this study | this study | N48°23'40.815" E9°30'10.053" | DE (Schwäbische Alb) | soil (grassland) |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | LH08HW8075 | this study | this study | N51°21'20.852" E10°31'1.083" | DE (Hainich-Dün) | soil (forest) |
| <i>Diplosphaera</i> | <i>Diplosphaera</i> sp. | SAG 2049 | n/a | this study | S8°16'25.200" E115°09'57.900" | ID (Bali, Lake Bratan) | freshwater |
| <i>Pseudomarvania</i> | <i>Pseudomarvania aerophytica</i> | CAUP H 7301 | FJ896222 | n/a | - | - | - |
| <i>Pseudomarvania</i> | <i>Pseudomarvania ampulliformis</i> | SAG 2047 | AB087559 | this study | - | JP (Taishaku-kyo Gorge) | subaerial (epixylic; tree bark) |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus</i> sp. | LH08SW8044 | this study | this study | N53°11'30.470" E13°55'49.216" | DE (Schorfheide-Chorin) | soil (forest) |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus</i> sp. | WB69 | KF144234 | this study | - | DE (Westerhöfer Bach) | semi-terrestrial (creek biofilm) |
| <i>Pseudostichococcus</i> | <i>Stichococcus mirabilis</i> | CCAP 379/3 | AJ311638 | AJ431679 | - | n/a | n/a |
| <i>Pseudostichococcus</i> | Trebouxiophyceae sp. | EO7-4 | FJ946882 | n/a | - | AQ (East Ongul) | freshwater |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus monalantoides</i> | SAG 380-1 | JX185690 | KM020066 | n/a | DE | freshwater (culture; <i>Enteromorpha</i>) |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus</i> sp. | 3GA1830SK9 | this study | n/a | N50°52'0.000" E10°50'0.000" | DE (Castle Gleichen) | subaerial (epilithic; sandstone) |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus</i> sp. | B3-2 | this study | n/a | N54°11'0.000" E7°53'0.000" | DE (Helgoland) | subaerial (epilithic; bunker wall) |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus</i> sp. | LH10HG3045 | this study | this study | N50°59'53.129" E10°25'58.616" | DE (Hainich-Dün) | soil (grassland) |
| <i>Pseudostichococcus</i> | <i>Stichococcus fragilis</i> | SAG 379-4 | n/a | this study | N41°35'58.300" W70°34'51.900" | US (Massachusetts) | freshwater (aquarium) |
| <i>Pseudostichococcus</i> | <i>Desmococcus spinocystis</i> | SAG 2067 | this study | this study | N43°45'07.700" E15°22'10.300" | HR (Island of Lavsa) | soil |
| <i>Pseudostichococcus</i> | <i>Diplosphaera</i> sp. | Lsn | JN573894 | n/a | - | - | - |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus</i> sp. | AEG6B K4796 | this study | n/a | N48°24'4.600" E9°26'30.100" | DE (Schwäbische Alb) | soil (grassland) |
| <i>Pseudostichococcus</i> | <i>Pseudostichococcus</i> sp. | HEG7MW B K4277 | this study | n/a | N51°16'24.897" E10°24'37.485" | DE (Hainich-Dün) | soil (grassland) |
| <i>Pseudostichococcus</i> | <i>Stichococcus minutus</i> | NJ-17 | JN400256 | n/a | - | AQ (Zhongshan Station) | terrestrial (wet rocks) |
| <i>Stichococcus bacillaris</i> | <i>Chlorosphaera klebsii</i> | CCAP 215/1 | FR865689 | FR865689 | - | - | - |
| <i>Stichococcus bacillaris</i> | <i>Gloeotila</i> cf. <i>protogenita</i> | SAG 56.91 | AM412752 | AM412752 | - | - | - |
| <i>Stichococcus bacillaris</i> | <i>Gloeotila scopulina</i> | SAG 335-8 | AM412753 | AM412753 | - | - | - |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | HEW1B K3375 | this study | n/a | N51°11'7.278" E10°19'25.036" | DE (Hainich-Dün) | soil (forest) |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | NIES-529 | AB488606 | n/a | - | - | - |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | RCC 1054 | KF899844 | n/a | - | - | - |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | SAG 335-3 | EU434029 | n/a | - | - | - |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | SAG 379-1b | AJ311637 | AJ431678 | - | CH (Basel) | freshwater (water dish) |

Table S2. (continuation)

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|--------------------------------|--------------------------------|------------------------|------------|------------|-------------------------------|-------------------------|----------------------------------|
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | SAG 379-1d | n/a | this study | n/a | US (Alaska) | n/a |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | SAG 379-2 | HE610125 | HE610125 | - | DE (Bernburg) | n/a |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | siva2011 | JN168788 | n/a | - | - | - |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | UTEX 315 | AM412751 | AM412751 | - | DE (Bernburg) | n/a |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | WB13 | KF144231 | this study | - | DE (Westerhöfer Bach) | semi-terrestrial (creek biofilm) |
| <i>Stichococcus bacillaris</i> | <i>Stichococcus bacillaris</i> | WB74 | KF144232 | this study | - | DE (Westerhöfer Bach) | semi-terrestrial (creek biofilm) |
| <i>Stichococcus</i> clade1 | <i>Stichococcus bacillaris</i> | K4-4 | AB055866 | n/a | - | - | - |
| <i>Stichococcus</i> clade1 | <i>Stichococcus</i> sp. | DB6-27 | KF144235 | n/a | - | - | - |
| <i>Stichococcus</i> clade1 | <i>Stichococcus</i> sp. | HEG8B K4544 | this study | n/a | N51°16'16.527" E10°25'4.6" | DE (Hainich-Dün) | soil (grassland) |
| <i>Stichococcus</i> clade1 | <i>Stichococcus</i> sp. | HEW1B K3359 | this study | n/a | N51°11'7.278" E10°19'25.036" | DE (Hainich-Dün) | soil (forest) |
| <i>Stichococcus</i> clade1 | <i>Stichococcus</i> sp. | KP09AW1004 | KP081407 | this study | - | DE (Schwäbische Alb) | subaerial (epixylic; tree bark) |
| <i>Stichococcus</i> clade1 | <i>Stichococcus</i> sp. | KS075SM6T | this study | this study | S3°58'31.436" W79°4'17.341" | EC (San Francisco) | subaerial (epixylic; tree bark) |
| <i>Stichococcus</i> clade1 | <i>Stichococcus</i> sp. | SAG 2481 (=LH08AW8002) | KP081394 | n/a | - | DE (Schwäbische Alb) | soil (forest) |
| <i>Stichococcus</i> clade1 | <i>Stichococcus</i> sp. | WB65 | KF144236 | n/a | - | - | - |
| <i>Stichococcus</i> clade2 | <i>Stichococcus</i> sp. | 594-GA18 | EU282451 | n/a | - | RU (Siberia, Kolyma) | soil (permafrost) |
| <i>Stichococcus</i> clade2 | <i>Stichococcus</i> sp. | D4-2A | KF144238 | this study | - | DE (Deinschwanger Bach) | semi-terrestrial (creek biofilm) |
| <i>Stichococcus</i> clade2 | <i>Stichococcus</i> sp. | ITS1A C11 | n/a | HM490287 | - | AQ (Miers Valley) | subaerial (hypolithic) |
| <i>Stichococcus</i> clade2 | <i>Stichococcus</i> sp. | ITS1A H8 | n/a | HM490288 | - | AQ (Miers Valley) | subaerial (hypolithic) |
| <i>Stichococcus</i> clade2 | <i>Stichococcus</i> sp. | LH08AW8104 | KP081396 | this study | - | DE (Schwäbische Alb) | soil (forest) |
| <i>Stichococcus</i> clade2 | <i>Stichococcus</i> sp. | SAG 2482 (=LH08AW8023) | KP081395 | this study | - | DE (Schwäbische Alb) | soil (forest) |
| <i>Stichococcus</i> clade3 | <i>Stichococcus</i> sp. | A43 | HQ418417 | n/a | - | - | soil; Yellowstone NP |
| <i>Stichococcus</i> clade3 | <i>Stichococcus</i> sp. | clone: DA-04 | AB257661 | n/a | - | - | rock surface, Swiss Alps |
| <i>Stichococcus</i> clade3 | <i>Stichococcus</i> sp. | KS106CL6T | this study | this study | S4°6'52.681" W79°10'30.940" | EC (Cajanuma) | subaerial (epixylic; tree bark) |
| <i>Stichococcus</i> clade3 | <i>Stichococcus</i> sp. | KS108CL6T | n/a | this study | S4°6'52.681" W79°10'30.940" | EC (Cajanuma) | subaerial (epixylic; tree bark) |
| <i>Stichococcus</i> clade3 | <i>Stichococcus</i> sp. | SAG 2408 (=WB8) | KF144239 | this study | - | DE (Westerhöfer Bach) | semi-terrestrial (creek biofilm) |
| <i>Stichococcus</i> clade4 | <i>Stichococcus</i> sp. | clone B1_3_1E_88 | JQ627437 | n/a | - | - | - |
| <i>Stichococcus</i> clade4 | <i>Stichococcus</i> sp. | HEG7SWWB K4387 | this study | n/a | N51°16'24.897" E10°24'37.485" | DE (Hainich-Dün) | soil (grassland) |
| <i>Stichococcus</i> clade4 | <i>Stichococcus</i> sp. | LH08SG1073 | this study | this study | N53°5'14.712" E13°58'10.717" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Stichococcus</i> clade4 | <i>Stichococcus</i> sp. | SAG 2406 (=WB47) | KF144240 | this study | - | DE (Westerhöfer Bach) | semi-terrestrial (creek biofilm) |
| <i>Stichococcus</i> clade5 | <i>Stichococcus</i> sp. | DZ K4g | this study | n/a | N51°33'29.700" E9°56'20.300" | DE (Göttingen) | subaerial (epilithic; roof tile) |

Table S2. (continuation)

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|--------------------------------|--------------------------------|----------------|------------|------------|----------------------------------|-------------------------|--|
| <i>Stichococcus</i> clade5 | <i>Stichococcus</i> sp. | KP09HW3004 | this study | this study | N51°16'17.900" E10°18'38.700" | DE (Hainich-Dün) | subaerial (epixylic; tree bark) |
| <i>Stichococcus</i> clade5 | <i>Stichococcus</i> sp. | KS090BU1T | n/a | this study | S4°6'49.852" W78°58'1.012" | EC (Bombuscaro) | subaerial (epixylic; tree bark) |
| <i>Stichococcus</i> clade5 | <i>Stichococcus</i> sp. | KS305SM6L | this study | this study | S3°58'31.436" W79°4'17.341" | EC (San Francisco) | subaerial (epiphytic; leaves) |
| <i>Stichococcus</i> clade5 | <i>Stichococcus</i> sp. | LH08SW1099 | this study | this study | N52°54'3.050" E13°50'46.921" | DE (Schorfheide-Chorin) | soil (forest) |
| <i>Stichococcus</i> clade6 | <i>Stichococcus</i> sp. | SAG 107.80 | n/a | this study | N45°15'33.900" W62°56'0.100" | CA (Nova Scotia) | n/a |
| <i>Stichococcus</i> clade6 | <i>Stichococcus</i> sp. | SAG 108.80 | KM020175 | n/a | - | - | - |
| <i>Stichococcus</i> clade6 | <i>Stichococcus</i> sp. | WB2 | n/a | this study | N51°45'0.000" E10°5'0.000" | DE (Westerhöfer Bach) | semi-terrestrial (creek biofilm) |
| <i>Stichococcus</i> clade6 | <i>Stichococcus</i> sp. | WB66 | this study | n/a | N51°45'0.000" E10°5'0.000" | DE (Westerhöfer Bach) | semi-terrestrial (creek biofilm) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | W1118 | JN573887 | n/a | - | - | - |
| <i>Stichococcus</i> clade7 | <i>Stichococcus bacillaris</i> | NJ-10 | JN400255 | n/a | - | AQ (Zhongshan Station) | terrestrial (wet rocks) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus bacillaris</i> | s3 | AY380557 | n/a | - | AQ | soil |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | FG2/4.2 | KM020048 | KM020048 | - | - | - |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | HEG9B K2590 | this study | n/a | N51°13'26.031" E10°22'50.834" | DE (Hainich-Dün) | soil (grassland) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | LH08AW8025 | KP081397 | n/a | - | DE (Schwäbische Alb) | soil (forest) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | LH08SG5057 | this study | this study | N53°6'26.830" E14°0'1.885" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | LH08SG5079 | this study | this study | N53°6'26.830" E14°0'1.885" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | LH08SG5090 | this study | this study | N53°6'26.830" E14°0'1.885" | DE (Schorfheide-Chorin) | soil (grassland) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | LH10HG2063 | this study | this study | N51°0'2.696" E10°25'48.036" | DE (Hainich-Dün) | soil (grassland) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | LH10HG7072 | this study | this study | N51°16'24.897" E10°24'37.485" | DE (Hainich-Dün) | soil (grassland) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | LH10HG7073 | this study | this study | N51°16'24.897" E10°24'37.485" | DE (Hainich-Dün) | soil (grassland) |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | SAG 2059 | AY762604 | n/a | - | - | - |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | SAG 2060 | AY762606 | n/a | - | - | - |
| <i>Stichococcus</i> clade7 | <i>Stichococcus</i> sp. | SAG 2119 | KM116460 | n/a | - | - | - |
| <i>Stichococcus deasonii</i> | <i>Stichococcus deasonii</i> | SAG 2139 | DQ275460 | this study | - | US (Dauphin Island) | soil |
| <i>Stichococcus deasonii</i> | <i>Stichococcus</i> sp. | ciidir1003 A11 | n/a | JN660592 | - | MX | soil (rhizosphere from commercial field) |
| <i>Stichococcus deasonii</i> | <i>Stichococcus</i> sp. | KSK870SM6T | n/a | this study | S3°58'31.436" W79°4'17.341" | EC (San Francisco) | subaerial (epixylic; tree bark) |
| <i>Stichococcus deasonii</i> | <i>Stichococcus</i> sp. | KSK937BU1L | n/a | this study | S4°6'49.852" W78°58'1.012" | EC (Bombuscaro) | subaerial (epiphytic; leaves) |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus bacillaris</i> | CCAP 379/7 | AB055864 | n/a | - | - | acid springs |

Table S2. (continuation)

| Clade | Species | Identifier | 18S | ITS2 | GPS | Land (region) | Habitat |
|--------------------------------|--------------------------------|-------------|------------|------------|----------------------------------|---------------------------|---------------------------------|
| <i>Stichococcus jenerensis</i> | <i>Stichococcus bacillaris</i> | D10-1 | AB055865 | n/a | - | - | - |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus jenerensis</i> | CCAP 379/5 | KJ756841 | n/a | - | - | marine |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus jenerensis</i> | LU1 | KF569726 | n/a | - | - | - |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus jenerensis</i> | SAG 2138 | DQ275461 | this study | - | MY (Kampong Kuala Jenera) | soil (crust on tree base) |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus</i> sp. | FACHB753 | EU045358 | n/a | - | - | - |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus</i> sp. | HEW2B K3412 | this study | n/a | N51°12'36.000" E10°22'11.800" | DE (Hainich-Dün) | soil (forest) |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus</i> sp. | KP09HW3001 | this study | n/a | N51°16'17.900" E10°18'38.700" | DE (Hainich-Dün) | subaerial (epixylic; tree bark) |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus</i> sp. | KS126SM6L | this study | this study | S3°58'31.436" W79°4'17.341" | EC (San Francisco) | subaerial (epiphytic; leaves) |
| <i>Stichococcus jenerensis</i> | <i>Stichococcus</i> sp. | MBIC10457 | AB183599 | n/a | - | - | - |

Legend. AQ=Antarctica, AT=Austria, CA=Canada, CH=Switzerland, CN=China, DE=Germany, EC=Ecuador, HR=Croatia, ID=Indonesia, JP=Japan, MX=Mexico, MY=Malaysia, NZ=New Zealand, RU=Russia, US=United States of America.

Table S3. List of the analyzed *Stichococcus*-like clades and their representative strains.

| OTU inferred from 18S rDNA similarity $\geq 99.5\%$; $\geq 99.0\%$ * | Representative strain | Author of denomination | Isolator/year | Synonymic GenBank accessions 18S/ITS2* |
|---|------------------------|------------------------|---------------|--|
| OTU8* <i>Diplosphaera</i> | SAG 48.86 | Broady 1983 | Broady/1979 | <i>Stichococcus chodati</i> * <i>Stichococcus chlorelloides</i> <i>Chlorella sphaerica</i> * <i>Stichococcus bacillaris</i> <i>Diplosphaera</i> sp. |
| OTU2* <i>Stichococcus deasonii</i> | SAG 2139 | Neustupa et al. 2007 | Deason/1969 | <i>Stichococcus bacillaris</i> <i>Diplosphaera</i> sp. |
| OTU7* <i>Stichococcus jenerensis</i> * | SAG 2138 | Neustupa et al. 2007 | Neustupa/2000 | <i>Stichococcus bacillaris</i> |
| OTU9* <i>Pseudostichococcus</i> * | SAG 380-1 | Moewus 1951 | Moewus/1951 | <i>Stichococcus mirabilis</i> * <i>Stichococcus minutus</i> <i>Stichococcus fragilis</i> * <i>Desmococcus spinocystis</i> * <i>Diplosphaera</i> sp. |
| OTU6 <i>Stichococcus bacillaris</i> | SAG 379-1b | Vischer(?) | Vischer/1923 | <i>Stichococcus chloranthus</i> * <i>Chlorosphaera klebsii</i> <i>Gloeotila</i> cf. <i>protogenita</i> <i>Gloeotila scopulina</i> <i>Stichococcus bacillaris</i> |
| OTU1 <i>Stichococcus</i> clade1 | KS075SM6T | Hodač 2015 | Spitzer/2013 | |
| OTU2* <i>Stichococcus</i> clade2 | SAG 2482 (=LH08AW8023) | Hodač 2015 | Hodač/2008 | n/a |
| OTU3 <i>Stichococcus</i> clade3 | SAG 2408 (=WB8) | Hodač 2015 | Mohr/2005 | n/a |
| OTU4 <i>Stichococcus</i> clade4 | SAG 2406 (=WB47) | Hodač 2015 | Mohr/2005 | n/a |
| OTU5 <i>Stichococcus</i> clade5 | LH08SW1099 | Hodač 2015 | Hodač/2008 | n/a |
| OTU2* <i>Stichococcus</i> clade6 | SAG 107.80 | Lewin(?) | Lewin/1952 | n/a |
| OTU2* <i>Stichococcus</i> clade7 | LH10HG2063 | Hodač 2015 | Hodač/2010 | n/a |

Appendix | Chapter 4

Supporting Figures

Figure S1. ML phylogeny based on 18S rDNA sequences showing the phylogenetic position of *Jenufa* (**Fig. 2**).

Figure S2. ML phylogeny based on 18S rDNA sequences showing the phylogenetic position *Xylochloris* (**Fig. 3**).

Figure S1. Maximum likelihood phylogeny based on 18S rDNA sequences showing the phylogenetic position of *Jenufa* (Fig. 2).





Figure S2. Maximum likelihood phylogeny based on 18S rDNA sequences showing the phylogenetic position *Xylochloris* (Fig. 3).



Publications

Contributions to the papers and manuscript drafts included in this thesis

Chapter 1 | Molecular diversity of microscopic Green algae isolated from German soils

(Manuscript draft)

Ladislav Hodač, Christine Hallmann, Karolin Spitzer and Thomas Friedl

University of Göttingen, Experimental Phycology and Culture Collection of Algae (SAG),
Göttingen, Germany

Conceived and designed the experiments: LH CH TF

Performed the experiments: LH KS

Analyzed the data: LH, KS

Wrote the paper: LH

Chapter 2 | Diversity of microscopic green algae (Chlorophyta) in calcifying biofilms of two karstic streams in Germany

(Published in Geomicrobiology Journal; doi: 10.1080/01490451.2013.878418)

Ladislav Hodač¹, Nicole Brinkmann², Kathrin I. Mohr³, Gernot Arp², Christine Hallmann¹,
Jessica Ramm⁴, Karolin Spitzer¹, Thomas Friedl¹

¹University of Göttingen, Experimental Phycology and Culture Collection of Algae (SAG),
Göttingen, Germany, ²University of Göttingen, Geoscience Centre, Göttingen, Germany,
³Helmholtz Centre for Infection Research, Braunschweig, Germany, ⁴Brandenburg University of
Technology, Department of Freshwater Conservation, Bad Saarow, Germany

Conceived and designed the experiments: KM NB TF LH

Performed the experiments: KM NB LH JR KS CH

Analyzed the data: LH KM TF

Wrote the paper: LH TF GA KI NB

Chapter 3 | Phylogenetic analysis of polar *Chlorella* and *Stichococcus* suggests biogeography of airborne microalgae

(Manuscript draft submitted to FEMS Microbiology Ecology; FEMSEC-15-11-0606)

Ladislav Hodač¹, Christine Hallmann¹, Karolin Spitzer¹, Josef Elster^{2,3}, Fabian Faßhauer¹, Nicole Brinkmann⁴, Daniela Lepka¹, Vaibhav Diwan¹ and Thomas Friedl¹

¹Experimental Phycology and Culture Collection of Algae (SAG), University of Göttingen, Germany, ²Centre for Polar Ecology, University of South Bohemia, České Budějovice, Czech Republic, ³Institute of Botany, Phycology Centrum, Academy of Sciences of the Czech Republic, Třeboň, Czech Republic, ⁴Department of Forest Botany, University of Göttingen, Germany

Conceived and designed the experiments: LH TF JE

Performed the experiments: LH KS CH NB JE FF DL VD

Analyzed the data: LH, KS, TF

Wrote the paper: LH

Chapter 4 | Molecular evidence for the wide distribution of two lineages of terrestrial green algae (Chlorophyta) from tropics to temperate zone

(Published in ISRN Ecology; doi: 10.5402/2012/795924)

Ladislav Hodač¹, Christine Hallmann¹, Helen Rosenkranz², Fabian Faßhauer¹ and Thomas Friedl¹

¹University of Göttingen, Experimental Phycology and Culture Collection of Algae (SAG), Göttingen, Germany, ²School of Biological Sciences, University of Bristol, Bristol, United Kingdom

Conceived and designed the experiments: LH TF

Performed the experiments: LH CH FF HR

Analyzed the data: LH

Wrote the paper: LH TF CH HR FF

Comprehensive publication list (2009-2015)

- Hodač L**, Brinkmann N, Mohr K, Arp G, Hallmann C, Ramm J, Spitzer K, Friedl T. 2015. Diversity of microscopic green algae (Chlorophyta) in calcifying biofilms of two karstic streams in Germany. *Geomicrobiology Journal* 32(3-4): doi:10.1080/01490451.2013.878418.
- Brinkmann N, **Hodač L**, Mohr KI, Hodačová A, Jahn R, Ramm J, Hallmann C, Arp G, Friedl T. 2015. Cyanobacteria and diatoms in biofilms of two karstic streams in Germany and changes of their communities along calcite saturation gradients. *Geomicrobiology Journal* 32(3-4): doi:10.1080/01490451.2014.901438.
- Hodač L**, Scheben, AP, Hojsgaard D, Paun O, Hörandl E. 2014. ITS polymorphisms shed light on hybrid evolution in apomictic plants: a case study from the *Ranunculus auricomus* complex. *PLoS ONE* 9(7): doi:10.1371/journal.pone.0103003.
- Patzelt DJ, **Hodač L**, Friedl T, Pietrasiak N, Johansen JR. 2014. Biodiversity of soil cyanobacteria in the hyper-arid Atacama Desert, Chile, assessed by culture dependent and independent approaches. *Journal of Phycology* 50: doi:10.1111/jpy.12196.
- Allan E, Bossdorf O, Dormann CF, Prati D, Gossner M, Tschardt T, Blüthgen N, Barto K, Bellach M, Birkhofer K, Boch S, Böhm S, Börschig C, Chatzinotas A, Christ S, Daniel R, Diekoetter T, Fischer C, Friedl T, Glaser K, Hallmann C, **Hodač L**, et al. 2013. Interannual variation in land-use intensity enhances grassland multidiversity. *PNAS* 111(1): doi:10.1073/pnas.1312213111.
- Hodač L**, Hallmann C, Rosenkranz H, Faßhauer F, Friedl T. 2012. Molecular evidence for the wide distribution of two lineages of terrestrial Green algae (Chlorophyta) over tropics to temperate zone. *ISRN Ecology* 2012: doi:10.5402/2012/795924.
- Němcová Y, Eliáš M, Škaloud P, **Hodač L**, Neustupa J. 2011. *Jenufa* gen. nov.: a new genus of coccoid green algae (Chlorophyceae, incertae sedis) previously recorded by environmental sequencing. *Journal of Phycology* 47(4): doi:10.1111/j.1529-8817.2011.01009.x.
- Nacke H, Thürmer A, Wollherr A, Will C, **Hodač L**, Herold N, Schöning I, Schrumpf M, Daniel R. 2011. Pyrosequencing-based assessment of bacterial community structure along different management types in german forest and grassland soils. *PLoS ONE* 6(2): doi:10.1371/journal.pone.0017000.
- Lang I, **Hodač L**, Friedl T, Feussner I. 2011. Fatty acid profiles and their distribution patterns in microalgae: a comprehensive analysis of more than 2000 strains from the SAG culture collection. *BMC Plant Biology* 11(124): doi:10.1186/1471-2229-11-124.

Danksagung

Meinem Betreuer, Herrn Prof. Dr. Thomas Friedl, verdanke ich die Möglichkeit, meine Doktorarbeit in der Sammlung von Algenkulturen Göttingen, einem ganz besonderen und traditionsreichen Standort der Phykologie, abfassen zu dürfen. Anfänglich konnte er mich für die terrestrischen Mikroalgen vollkommen begeistern und seine anregenden Denkanstöße waren für meine Arbeit wegweisend. Herzlichen Dank! Bei meinem Korreferenten, Herrn Prof. Dr. Rolf Daniel, bedanke ich mich für die kritische Begutachtung der vorliegenden Arbeit!

Mein ganz besonderer Dank gilt Frau Dr. Maike Lorenz; ihr kompetenter Rat und ihre Hilfe kamen mir in vielen Angelegenheiten immer sehr zugute. An dieser Stelle möchte ich mich auch bei Frau Prof. Dr. Elvira Hörandl herzlichst für ihr unschätzbare Entgegenkommen bedanken! Für die vielseitige Unterstützung, den regen Austausch und unzählige Diskussionen bedanke ich mich recht herzlich bei meiner Mitdotorandin Christine Hallmann. Dankbar bin ich auch für die immer freundliche und geduldige Hilfsbereitschaft meines Mitdotoranden Fabian Faßhauer sowie aller Mitarbeiterinnen und Mitarbeiter der EPSAG, insbesondere Ilse Kunkel, Marlis Heinemann, Gabrielle Curdt-Hollmann, Hella Timmermann, Reiner Freter, Kathrin I. Mohr, Nicole Brinkmann, Anastasija Kryvenda und Nataliya Rybalka. Mein Dank geht ebenso an die Mitglieder der Verwaltung des Instituts, insbesondere Herrn Dr. Hans Depta, der mir mehrfach hilfreich zur Seite stand.

Großer Dank gebührt meiner Familie, insbesondere meinen Eltern, die mich während des gesamten Promotionsstudiums hilfsbereit und unterstützend aufgemuntert haben. Auch meinem werten Freund David danke ich dafür, dass er mir stets zur Seite stand. Meiner Lebensgefährtin Karolin danke ich aus ganzem Herzen für alles. Ihre stetige und unermüdliche sowohl fachliche als auch private Unterstützung, oder kurz—ihr Dasein in meinem Leben, haben wesentlich zum Gelingen der vorliegenden Arbeit beigetragen.